



# MATH205 Final EXAM AND Solutions August 2018

Differential & Integral Calculus II (Concordia University)

**CONCORDIA UNIVERSITY**  
**Department of Mathematics & Statistics**

<b>Course</b> Mathematics	<b>Number</b> 205	<b>Section</b> CA
<b>Examination</b> Final	<b>Date</b> August 2018	<b>Pages</b> 2
<b>Instructors:</b>	<b>T. Hughes</b>	<b>Course Examiners</b> A. Atoyan & H. Proppe
<b>Special Instructions:</b>	Only calculators approved by the Department are allowed. For full marks show all your work.	

1. a) [5 marks] Sketch the graph of the function

$$f(x) = \begin{cases} x + 6 & -7 \leq x < -3 \\ 3 - \sqrt{9 - x^2} & -3 \leq x \leq 3 \end{cases}$$

on the interval  $-6 \leq x \leq 3$  and calculate the definite integral  $\int_{-6}^3 f(x) dx$  in terms of signed area (do not antidifferentiate).

- b) [5 marks] Use the Fundamental Theorem of Calculus, Part 1, to find  $F'(x)$ , given that

$$F(x) = \int_{x^2}^{x^3} \frac{\sin t}{t} dt$$

2. [10 marks] Evaluate the following indefinite integrals.

a)  $\int \frac{\sin^{-1} x}{\sqrt{1-x^2}} dx$

b)  $\int \frac{3x^2 + x + 1}{x(x^2 + 1)} dx$

3. [10 marks] Evaluate the following definite integrals (do not approximate):

a)  $\int_0^{\frac{\pi}{4}} \sec^4 x dx$

b)  $\int_0^2 x^2 \sqrt{4 - x^2} dx$

4. [10 marks] Evaluate the given improper integral, or show that it diverges.

a)  $\int_2^{\infty} \frac{1}{x \ln^2 x} dx$

b)  $\int_2^3 \frac{dx}{\sqrt{3-x}}$

5. a) [6 marks] Find the area bounded by the curves  $x = y$  and  $4x + y^2 = 12$

b) [6 marks] Find the volume of the solid obtained by rotating the region bounded by the curves  $y = x$  and  $y = \sqrt{x}$ , about the line  $y = 1$ .

c) [6 marks] Find the average value of the function  $f(x) = \frac{\ln x}{x^2}$  from  $x = 1$  to  $x = 2$ .

6. [9 marks] Find the limit of the sequence  $\{a_n\}$  as  $n \rightarrow \infty$  or prove that it does not exist:

a)  $a_n = \cos \frac{2n\pi}{5+6n}$

b)  $a_n = \frac{(\ln n)^2}{n}$

c)  $a_n = \left(1 + \frac{3}{n}\right)^n$

7. [8 marks] Determine whether the series is divergent or convergent. If it is convergent, then determine whether it is conditionally or absolutely convergent:

a)  $\sum_{n=1}^{\infty} \frac{\sin(\frac{1}{n})}{\sqrt{n}}$

b)  $\sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$

8. [6 marks] Find (a) the radius and (b) the interval of convergence of the series

$$\sum_{n=0}^{\infty} \frac{(x-2)^n}{n}$$

9. a) [4 marks] Find the Maclaurin series for  $f(x) = x^2 e^{x^3}$

b) [3 marks] i) Express the series as a function of  $t$ .  $\sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!}$

[2 marks] ii) Find the exact numerical sum of the series  $\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n+1}}{6^{2n+1} (2n+1)!}$


10. a) [5 marks] Use differentiation to find a power series representation for the function  $f(x) = \frac{1}{(1+x)^2}$

b) [5 marks] Find the (numerical) sum of the series  $\sum_{n=1}^{\infty} \frac{2^{n+1} + 3^{n-1}}{5^n}$

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.

MATH 205  
August 2018

1. (a)



$$A_1 = \frac{1}{2} \cdot b \cdot h = \frac{1}{2} (3)(3) = \frac{9}{2}$$

$$A_2 = (6)(3) - \frac{1}{2} \pi (3)^2 = 18 - \frac{9\pi}{2}$$

$$\begin{aligned} \therefore \int_{-6}^3 f(x) dx &= A_1 + A_2 \\ &= \left(\frac{9}{2}\right) + \left(18 - \frac{9\pi}{2}\right) \\ &= 22.5 - \frac{9\pi}{2} \\ &= \frac{45}{2} - \frac{9\pi}{2} \end{aligned}$$

$$1. (b) \quad F(x) = \int_{x^2}^{x^3} \frac{\sin t}{t} dt$$

$$= \int_{x^2}^0 \frac{\sin t}{t} dt + \int_0^{x^3} \frac{\sin t}{t} dt$$

$$F(x) = \int_0^{x^3} \frac{\sin t}{t} dt - \int_0^{x^2} \frac{\sin t}{t} dt$$

$$\begin{aligned} \therefore F'(x) &= \frac{\sin(x^3)}{x^3} \cdot 3x^2 - \frac{\sin(x^2)}{x^2} \cdot 2x \\ &= \frac{3 \sin(x^3)}{x} - \frac{2 \sin(x^2)}{x} \end{aligned}$$

$$2(a) \quad \int \frac{\sin^{-1}x}{\sqrt{1-x^2}} dx$$

$$\text{Let } u = \sin^{-1}x$$

$$\therefore du = \frac{1}{\sqrt{1-x^2}} dx$$

$$\therefore I = \int u du = \frac{1}{2} u^2 + C$$

$$= \frac{1}{2} (\sin^{-1}x)^2 + C$$

$$2 (b) \quad \int \frac{3x^2 + x + 1}{x(x^2 + 1)} dx$$

$$\frac{3x^2 + x + 1}{x(x^2 + 1)} = \frac{A}{x} + \frac{Bx + C}{x^2 + 1}$$

$$\begin{aligned} 3x^2 + x + 1 &= A(x^2 + 1) + (Bx + C)(x) \\ &= Ax^2 + A + Bx^2 + Cx \end{aligned}$$

$$3x^2 + x + 1 = x^2(A + B) + Cx + A$$

$$\therefore \quad \begin{aligned} A &= 1 \\ C &= 1 \end{aligned}$$

$$\begin{aligned} A + B &= 3 \\ \therefore B &= 2 \end{aligned}$$

$$\therefore I = \int \frac{1}{x} dx + \int \frac{2x + 1}{x^2 + 1} dx$$

$$= \int \frac{1}{x} dx + \int \frac{2x}{x^2 + 1} dx + \int \frac{dx}{x^2 + 1}$$

$$= \ln|x| + \ln|x^2 + 1| + \tan^{-1}x + C$$

$$3(a) \int_0^{\pi/4} \sec^4 x \, dx$$

$$\begin{aligned} \int \sec^4 x \, dx &= \int \sec^2 x \sec^2 x \, dx \\ &= \int (\tan^2 x + 1) \sec^2 x \, dx \\ &= \int (\tan^2 x \sec^2 x) \, dx + \int \sec^2 x \, dx \\ &= \frac{1}{3} \tan^3 x + \tan x + C \end{aligned}$$

$$\therefore I = \left( \frac{1}{3} \tan^3 x + \tan x \right) \Big|_0^{\pi/4}$$

$$= \left( \frac{1}{3} \tan^3 \left( \frac{\pi}{4} \right) + \tan \frac{\pi}{4} \right) - \left( \frac{1}{3} \tan^3 0 + \tan 0 \right)$$

$$= \frac{1}{3} (1)^3 + 1 - 0 - 0$$

$$= \frac{4}{3}$$



$$3(b) \int_0^2 x^2 \sqrt{4-x^2} \, dx$$

$$\text{Let } x = 2 \sin \theta$$

$$dx = 2 \cos \theta \, d\theta$$

$$\text{When } x=0, \theta=0$$

$$x=2, \theta = \frac{\pi}{2}$$

$$\therefore I = \int_0^{\frac{\pi}{2}} 4 \sin^2 \theta \sqrt{4-4 \sin^2 \theta} \, 2 \cos \theta \, d\theta$$

$$= 16 \int_0^{\frac{\pi}{2}} \sin^2 \theta \cos^2 \theta \, d\theta$$

$$= 16 \int_0^{\frac{\pi}{2}} \left( \frac{\sin 2\theta}{2} \right)^2 d\theta$$

$$\text{since,} \\ \sin 2\theta = 2 \sin \theta \cos \theta$$

$$= 4 \int_0^{\frac{\pi}{2}} \frac{\sin^2 2\theta}{1} d\theta$$

$$= 4 \int_0^{\pi/2} \left( \frac{1 - \cos 4\theta}{2} \right) d\theta$$

$$= 2 \int_0^{\pi/2} (1 - \cos 4\theta) d\theta$$

$$= 2 \left[ \theta - \frac{1}{4} \sin 4\theta \right] \Big|_0^{\pi/2}$$

$$= 2 \left[ \left( \frac{\pi}{2} - \frac{1}{4} \sin 2\pi \right) - \frac{1}{4} (0 - 0) \right]$$

$$= 2 \left[ \frac{\pi}{2} - 0 \right]$$

$$= \pi$$

$$4(a) \int_2^{\infty} \frac{1}{x \ln^2 x} dx$$

$$\text{let } u = \ln x \\ du = \frac{1}{x} dx$$

$$\left. \begin{array}{l} \text{when } x=2, \quad u = \ln 2 \\ x=t, \quad u = \ln t \end{array} \right\}$$

$$\therefore I = \lim_{t \rightarrow \infty} \int_{\ln 2}^{\ln t} \frac{du}{u^2}$$

$$= \lim_{t \rightarrow \infty} \left( -\frac{1}{u} \right) \Big|_{\ln 2}^{\ln t}$$

$$= \lim_{t \rightarrow \infty} \left[ \left( \frac{-1}{\ln t} \right) - \left( \frac{-1}{\ln 2} \right) \right]$$

$$= 0 + \frac{1}{\ln 2}$$

$$= \frac{1}{\ln 2}$$

$$4(b) \int_2^3 \frac{dx}{\sqrt{3-x}} \quad dx$$

$$= \lim_{t \rightarrow 3^-} \int_2^t \frac{dx}{\sqrt{3-x}}$$

$$= \lim_{t \rightarrow 3^-} \left. -2(3-x)^{1/2} \right|_2^t$$

$$= \lim_{t \rightarrow 3^-} \left[ -2(3-t)^{1/2} + 2(3-2)^{1/2} \right]$$

$$= 0 + 2(1)$$

$$= 2$$

$$5(a) \quad x = y$$

$$4x + y^2 = 12$$

$$x = \frac{12 - y^2}{4}$$

At intersection,

$$x = 3 - \frac{1}{4}y^2$$

$$y = \frac{12 - y^2}{4}$$

$$4y = 12 - y^2$$

$$y^2 + 4y - 12 = 0$$


$$(y+6)(y-2) = 0$$

$$y = 2$$

$$(2, 2)$$

$$\text{or } y = -6$$

$$(-6, -6)$$




$$\begin{aligned} \text{Area} &= \int_{-6}^2 \left[ \left( 3 - \frac{1}{4}y^2 \right) - y \right] dy \\ &= \left( 3y - \frac{y^3}{12} - \frac{y^2}{2} \right) \bigg|_{-6}^2 \end{aligned}$$

$$= (6 - \frac{2}{3} - 2) - (-18 + 18 - 18)$$

$$= 4 - \frac{2}{3} + 18 = 21 \frac{1}{3} = \frac{64}{3}$$

5(b)



$$V = \int_0^1 A(x) dx$$

$$= \int_0^1 \pi [R^2 - r^2] dx$$

$$= \pi \int_0^1 [(1-x)^2 - (1-\sqrt{x})^2] dx$$

$$= \pi \int_0^1 [(1-2x+x^2) - (1-2\sqrt{x}+x)] dx$$

$$= \pi \int_0^1 [-3x + x^2 + 2\sqrt{x}] dx$$

$$= \pi \left[ -\frac{3x^2}{2} + \frac{x^3}{3} + 2 \cdot \frac{x^{3/2}}{3} \right]_0^1$$

$$= \pi \left[ \left( -\frac{3}{2} + \frac{1}{3} + \frac{4}{3} \right) - (0) \right]$$

$$= \pi \left( \frac{5}{3} - \frac{3}{2} \right) = \frac{\pi}{6}$$



$$5(c) \quad f_{avg} = \frac{1}{2-1} \int_1^2 \frac{\ln x}{x^2} dx$$

$$= \int_1^2 \frac{\ln x}{x^2} dx$$

$$\text{let } u = \ln(x)$$

$$du = \frac{1}{x} dx$$

$$dv = \frac{1}{x^2} dx$$

$$v = -\frac{1}{x}$$

$$\therefore I = -\frac{\ln x}{x} \Big|_1^2 + \int_1^2 \frac{1}{x^2} dx$$

$$= \left( -\frac{\ln 2}{2} - \left( -\frac{\ln 1}{1} \right) \right) + \left( -\frac{1}{x} \right) \Big|_1^2$$

$$= -\frac{\ln 2}{2} + 0 + \left[ -\frac{1}{2} - \left( -\frac{1}{1} \right) \right]$$

$$= -\frac{\ln 2}{2} - \frac{1}{2} + 1$$

$$= \frac{1 - \ln 2}{2}$$

$$6) (a) \lim_{n \rightarrow \infty} \cos \left( \frac{2n\pi}{5+6n} \right)$$

$$= \cos \left[ \lim_{n \rightarrow \infty} \frac{2n\pi}{5+6n} \right]$$

$$= \cos \left( \frac{\pi}{3} \right) = \frac{1}{2}$$


---

$$(b) \lim_{n \rightarrow \infty} \frac{(\ln n)^2}{n} = \lim_{n \rightarrow \infty} \frac{2 \ln n \cdot \frac{1}{n}}{1}$$

$$= \lim_{n \rightarrow \infty} 2 \cdot \frac{\ln n}{n}$$

$$= 2 \cdot \lim_{n \rightarrow \infty} \frac{\frac{1}{n}}{1} = 0$$


---

$$c) \lim_{n \rightarrow \infty} \left( 1 + \frac{3}{n} \right)^n$$

$$= \lim_{n \rightarrow \infty} \left( \left( 1 + \frac{3}{n} \right)^{n/3} \right)^3$$

$$= e^3$$

$$7(a) \quad \sum_{n=1}^{\infty} \frac{\sin(1/n)}{\sqrt{n}}$$

$$\text{Let } a_n = \frac{\sin(1/n)}{\sqrt{n}}$$

$$b_n = \frac{1}{n^{3/2}}$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \left( \frac{\sin 1/n}{\sqrt{n}} \right) \cdot \left( \frac{n^{3/2}}{1} \right)$$

$$= \lim_{n \rightarrow \infty} n \cdot \sin(1/n)$$

$$= \lim_{n \rightarrow \infty} \frac{\sin(1/n)}{(1/n)} = 1$$

$\therefore$  By the limit comparison test,  $\sum a_n$  and  $\sum b_n$  both converge or both diverge.

But  $\sum \frac{1}{n^{3/2}}$  converges,

( $p$ -series,  $p = 3/2 > 1$ )

$\therefore \sum_{n=1}^{\infty} a_n$  converges (and converges absolutely)

$$7(b) \quad \sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$$

$$\text{let } b_n = \frac{1}{\ln n}$$

$$\lim_{n \rightarrow \infty} b_n = 0$$

$$\text{and } \frac{1}{\ln(n+1)} < \frac{1}{\ln n}$$

$\therefore$  By the Alternating Series Test,

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{\ln n} \text{ is convergent.}$$

$$\text{Next, consider } \sum_{n=2}^{\infty} \frac{1}{\ln n}$$

$$\ln n < n$$

$$\therefore \frac{1}{\ln n} > \frac{1}{n}$$

$$\sum_{n=2}^{\infty} \frac{1}{n} \text{ is divergent} \\ (\text{p-series, } p=1)$$

Hence,  $\sum_{n=2}^{\infty} \frac{1}{\ln n}$  is divergent by the Comparison Test

$\therefore$  The series is conditionally convergent.

$$8. \sum_{n=0}^{\infty} \frac{(x-2)^n}{n}$$

Ratio Test :

$$\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{(x-2)^{n+1}}{(n+1)} \cdot \frac{n}{(x-2)^n} \right|$$

$$= \left( \frac{n}{n+1} \right) |x-2|$$

$$\therefore \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = |x-2|$$

Converges if  $|x-2| < 1$

$$-1 < x-2 < 1$$

$$1 < x < 3$$

Radius of convergence = 1

Check endpoints :

(a) If  $x=1$  : we have

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{n}, \text{ convergent by the A.S.T}$$

(b) If  $x=3$  : we have  $\sum_{n=0}^{\infty} \frac{1}{n}$ , divergent (Harmonic series)

$\therefore$  Interval of Convergence =  $[1, 3)$

$$9(a) \quad e^u = \sum_{n=0}^{\infty} \frac{u^n}{n!}$$

$$\therefore e^{x^3} = \sum_{n=0}^{\infty} \frac{(x^3)^n}{n!}$$

$$= \sum_{n=0}^{\infty} \frac{x^{3n}}{n!}$$

$$\therefore x^2 e^{x^3} = \sum_{n=0}^{\infty} \frac{x^{3n+2}}{n!}$$

$$9. (b) \sum_{n=0}^{\infty} \frac{(-1)^n \cdot t^{2n+1}}{(2n+1)!} = \sin t$$

$$\sum_{n=0}^{\infty} \frac{(-1)^n \cdot \left(\frac{\pi}{6}\right)^{2n+1}}{(2n+1)!} = \sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$10. (a) f(x) = \frac{1}{1+x}$$

$$\begin{aligned} \frac{1}{1+x} &= \frac{1}{1-(-x)} \\ &= 1 - x + x^2 - x^3 + x^4 - \dots \end{aligned}$$

$$\therefore \frac{-1}{(1+x)} = -1 + x - x^2 + x^3 - x^4 - \dots$$

Differentiate both sides,

$$\therefore \frac{1}{(1+x)^2} = 1 - 2x + 3x^2 - 4x^3 - \dots$$

$$= \sum_{n=0}^{\infty} (-1)^n (n+1) x^n$$



$$10 \text{ (b)} \quad \sum_{n=1}^{\infty} \frac{2^{n+1} + 3^{n-1}}{5^n}$$

$$= 2 \sum_{n=1}^{\infty} \left(\frac{2}{5}\right)^n + \frac{1}{5} \sum_{n=1}^{\infty} \left(\frac{3}{5}\right)^{n-1}$$

$$= 2 \left[ \frac{2/5}{1 - 2/5} \right] + \frac{1}{5} \left[ \frac{1}{1 - 3/5} \right]$$

$$= 2 \left[ \frac{2/5}{3/5} \right] + \frac{1}{5} \left[ \frac{1}{2/5} \right]$$

$$= 2 \cdot \frac{2}{\cancel{5}} \cdot \frac{\cancel{5}}{3} + \frac{1}{\cancel{5}} \cdot \frac{\cancel{5}}{2}$$

$$= \frac{4}{3} + \frac{1}{2}$$

$$= \frac{8+3}{6}$$

$$= \frac{11}{6}$$