



Final 2018 math 205

MATH (Concordia University)

CONCORDIA UNIVERSITY
Department of Mathematics & Statistics

Course	Number	Sections
Mathematics	205	All
Examination	Date	Pages
Final	April 2018	2
Instructors:	J. de Ronde, H. Greenspan, H. Harb S. Mathieu-Soucy, N. Rossokhata	Course Examiners A. Atoyan & H. Proppe
Special Instructions:	Only calculators approved by the Department are allowed. For full marks show all your work.	

[12] 1. (a) Sketch a graph of the function

$$f(x) = \begin{cases} 4 + x & \text{for } -4 \leq x < -2 \\ 2 - \sqrt{4 - x^2} & \text{for } -2 \leq x \leq 2 \end{cases}$$

on the interval $-4 \leq x \leq 2$ and calculate the definite integral $\int_{-4}^2 f(x) dx$ in terms of signed area (*do not* antidifferentiate).

(b) Now graph $g(x) = f(x) - 1$. What is the value of $\int_{-4}^2 g(x) dx$?

(c) Find the derivative $F'(x)$ of $F(x) = \sin x + \int_0^{3x} \sqrt{a^2 + u^2} \cos u \, du$, where a is a parameter, and use it to determine whether $F(x)$ is increasing or decreasing at $x = \pi$.

[10] 2. Find the following indefinite integrals:

$$(a) \int x \arcsin(x^2) dx \qquad (b) \int \frac{x}{x^2 - 3x - 4} dx$$

[12] 3. Evaluate the following definite integrals (give the exact answers):

$$(a) \int_e^{e^4} \frac{1}{x \sqrt{\ln x}} dx \qquad (b) \int_0^{\pi/6} \sqrt{1 + \cos(2x)} dx$$

[6] 4. Find $F(t)$ such that $F'(t) = \left(\frac{\sec t}{\tan t}\right)^4$ and $F\left(\frac{\pi}{4}\right) = -1$

[8] 5. Evaluate the given improper integral or show that it diverges:

$$(a) \int_1^{\infty} \frac{\sqrt{x}}{x + \sqrt{x}} dx \qquad (b) \int_0^{\pi/2} \frac{\cos x}{\sqrt{1 - \sin x}} dx$$

- [17] 6. (a) Sketch the curves $y = 6x - x^2$ and $y = x^2 - 2x$, and find the area enclosed by these curves.
- (b) Find the volume of the solid obtained by rotating the region bounded by the curve $y = x^3$ and the lines $x = 0$ and $y = 8$ about the axis $x = 2$.
- (c) A cup of coffee cools from 95°C to about 61°C in 30 minutes according to the equation $T(t) = 20 + 75e^{-0.02t}$. Find the average temperature T_{av} of the coffee during the interval $[0, 30]$.
- [9] 7. Find the limit of the sequence $\{a_n\}$ at $n \rightarrow \infty$ or prove that it does not exist:

(a) $a_n = \frac{n \cos(n\pi)}{\sqrt{9 + n^3}}$ (b) $a_n = n \sin\left(\frac{\pi}{n}\right)$ (c) $a_n = 2 \ln(1 + 3n) - \ln(4 + n^2)$

- [8] 8. Determine whether the series is divergent or convergent, and if convergent, then absolutely or conditionally :

(a) $\sum_{n=1}^{\infty} \frac{(-1)^n n!}{(n + 100)!}$ (b) $\sum_{n=1}^{\infty} (-1)^n \frac{\ln n}{3n + 1}$

- [10] 9. Find the radius and the interval of convergence of the following series

(a) $\sum_{n=1}^{\infty} \frac{(3x - 2)^n}{5^n}$ (b) $\sum_{n=1}^{\infty} \frac{(x - 4)^n}{n^2}$

- [8] 10. (a) Find the Maclaurin series for $f(x) = x e^{x^2}$
(Hint: start with the series for e^t , then replace t with x^2 . . .)
- (b) Use differentiability of the power series to find the sum

$F(x) = \sum_{n=0}^{\infty} \frac{(x + 1)^{n+1}}{n + 1}$ within its radius of convergence.

- [5] **Bonus Question.** Find all continuous functions $f(x)$ other than $f(x) = 0$ satisfying the equation

$$\int_0^x f(t) dt = (f(x))^2 - (f(0))^2$$

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.