

CONCORDIA UNIVERSITY
Department of Mathematics and Statistics

Course	Number	Section(s)
Mathematics	204	All
Examination	Date	Pages
Final	April 2022	3

Instructors

Course Examiner

C. L. Santana, S. Saputra, H. Pezeshk, T. Hughes, M. Khasraghi; A. Kokotov

Special Instructions: Only approved calculators are allowed !

1. [10] Solve the system of equations

$$\begin{cases} x_1 + x_2 + x_3 + x_4 = 10 \\ 2x_1 - x_2 + x_3 - x_4 = -1 \\ 3x_1 - x_3 - 2x_4 = -8 \\ -x_1 + 3x_2 - x_4 = 1 \\ 5x_1 - x_2 - 3x_4 = -9 \end{cases}$$

2. [10] Solve the following equation for (4×4) matrix X:

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 4 \end{pmatrix} X \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

3. [10] For which a the system

$$\begin{cases} x + y + z = 2 \\ x + 2y + 2z = 3 \\ x + a(a-1)z = a + 1 \end{cases}$$

- has infinitely many solutions?
- has no solutions?
- has the unique solution?

4. [10] The plane α contains the line l with parametric equation

$$\begin{cases} x = 1 + t \\ y = 2 - 3t \\ z = 3 - t \end{cases}$$

and passes through the point $P = (1, 2, 1)$. Find the distance $\text{dist}(Q, \alpha)$ from the point $Q = (1, 1, 1)$ to the plane α .

5. [10] The plane α passes through the points $A = (1, 2, 3)$, $B = (2, -1, 4)$ and $C = (1, 4, 1)$.

The plane β is defined via the equation

$$x + 2y + 3z = 1.$$

Write down the parametric equation of the line of intersection of α and β .

6. [10] The vectors

$$\vec{u} = (a/2, 1, 3/2)$$

$$\vec{v} = (a/3, a/3, 4/3)$$

and

$$\vec{w} = (a, 2, a + 6)$$

have the same initial point and lie in the same plane. Find all the possible values of a .

7. [10] Find the value of the determinant

$$\begin{vmatrix} 1 & 2 & 0 & 3 & 4 \\ 2 & 3 & 0 & 4 & 1 \\ 7 & 8 & 5 & 9 & 5 \\ 3 & 4 & 0 & 1 & 2 \\ 4 & 1 & 0 & 2 & 3 \end{vmatrix}$$

8. [10] Find a basis for the solution space of the homogeneous system

$$\begin{cases} x_1 + x_2 + x_3 + x_4 - x_5 = 0 \\ 2x_1 - x_2 + x_3 - x_4 + x_6 = 0 \\ 4x_1 + x_2 + 3x_3 + x_4 - 2x_5 + x_6 = 0 \end{cases}$$

9. [10] The matrix

$$A = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 4 & 1 & 0 & 0 \\ 2 & 2 & 1 & 0 \\ 2 & -1 & 0 & 3 \end{pmatrix}$$

has an eigenvector $(1, 2, 3, 4)$. Find an eigenvalue of the matrix A .

10. [10] Find an invertible 2×2 matrix P such that the matrix

$$P^{-1} \begin{pmatrix} 2 & \sqrt{2} \\ \sqrt{2} & 3 \end{pmatrix} P$$

is diagonal.

¹The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.

MATH.204.

Final Exam Solutions.

Winter. 2022.

1. $\kappa_1 = 1$

$\kappa_2 = 2$

$\kappa_3 = 3$

$\kappa_4 = 4.$

$$2. \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 4 \end{bmatrix} \times \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$B \quad X \quad A \quad = \quad C$$

$$BXA = C.$$

must compute A^{-1}, B^{-1} , if they exist.

$$A^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{bmatrix}; B^{-1} = \begin{bmatrix} 1 & -1 & -1 & -1 \\ 0 & 1/2 & 0 & 0 \\ 0 & 0 & 1/3 & 0 \\ 0 & 0 & 0 & 1/4 \end{bmatrix}.$$

$$BXA = C$$

$$BXAA^{-1} = CA^{-1} = C$$

$$B^{-1}BX = B^{-1}C$$

$$X = B^{-1}C = \begin{bmatrix} 1 & -1 & -1 & 0 \\ 0 & 1/2 & 0 & 0 \\ 0 & 0 & 1/3 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

3.

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 1 & 2 & 2 & 3 \\ 1 & 0 & a(a-1) & a+1 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & (a^2-a) & a \end{array} \right].$$

No. sol.: $a=1$.

one sol.: $a \neq 0, 1$.

Infinite sols.: $a=0$.

$$4. \quad p_0 = (1, 2, 3); \quad \vec{n} = (1, -3, -1)$$

$$p = (1, 2, 1); \quad Q = (1, 1, 1).$$

$$\vec{p_0 p} = (0, 0, -2).$$

$$\vec{n} \times \vec{p_0 p} = (6, 2, 0).$$

Equation of plane.

$$6(x-1) + 2(y-2) = 0$$

$$3x + y = 5.$$

Distance from Q to plane.

$$D = \frac{|3(1) + (1) - 5|}{\sqrt{(3)^2 + (1)^2}} = \frac{1}{\sqrt{10}} = \frac{\sqrt{10}}{10}.$$

$$5. A(1, 2, 3), B(2, -1, 4), C(1, 4, 1).$$

$$\vec{AB} = (1, -3, 1).$$

$$\vec{AC} = (0, 2, -2).$$

$$\vec{AB} \times \vec{AC} = (4, 2, 2).$$

$$\text{Plane } \alpha: 4(x-1) + 2(y-2) + 2(z-3) = 0$$

$$2x + y + z = 7.$$

Computing line of intersection of the two planes.

$$\left[\begin{array}{ccc|c} 2 & 1 & 1 & 7 \\ 1 & 2 & 3 & 1 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|c} 1 & 0 & -1/3 & 13/3 \\ 0 & 1 & 5/3 & -5/3 \end{array} \right] \Rightarrow$$

$$\text{S.S.: } X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{13}{3} \\ -\frac{5}{3} \\ 0 \end{bmatrix} + t \begin{bmatrix} \frac{1}{3} \\ -\frac{5}{3} \\ 1 \end{bmatrix}.$$

$$x = \frac{13}{3} + \frac{t}{3}$$

$$y = -\frac{5}{3} - \frac{5}{3}t \quad z = t.$$

$$6. \vec{u} = \left(\frac{a}{2}, 1, \frac{3}{2}\right)$$

$$\vec{v} = \left(\frac{a}{3}, \frac{a}{3}, \frac{4}{3}\right)$$

$$\vec{w} = (a, 2, a+6).$$

For the three vectors to lie on the same plane, $\vec{w} \cdot (\vec{u} \times \vec{v})$ must equal 0.

$$\vec{u} \times \vec{v} = (8-3a, -a, a^2-2a).$$

$$(a, 2, a+6) \cdot (8-3a, -a, a^2-2a) = 0$$

$$a^3 + a^2 - 6a = 0$$

$$a(a^2 + a - 6) = 0$$

$$a(a+3)(a-2) = 0$$

$$a = 0, -3, 2.$$

$$7. \det(A) = 800.$$

$$\begin{array}{c}
 8. \\
 \begin{matrix} x & y & z & u & v & w \\
 \begin{bmatrix} 1 & 1 & 1 & 1 & -1 & 0 \\
 2 & -1 & 1 & -1 & 0 & 1 \\
 4 & 1 & 3 & 1 & -2 & 1 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 0 & 2/3 & 0 & -1/3 & 1/3 \\
 0 & 1 & 1/3 & 1 & -2/3 & -1/3 \\
 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}
 \end{matrix}
 \end{array}$$

\Rightarrow Infinite Solutions.

$$\text{S.S.: } X = \begin{bmatrix} x \\ y \\ z \\ u \\ v \\ w \end{bmatrix} = q \begin{bmatrix} -2/3 \\ -1/3 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + r \begin{bmatrix} 0 \\ -1 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} 1/3 \\ 2/3 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1/3 \\ 1/3 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}.$$

$\bar{\alpha} \qquad \qquad \bar{\beta} \qquad \qquad \bar{\gamma} \qquad \qquad \bar{\delta}$

The basis vectors are: $\bar{\alpha}, \bar{\beta}, \bar{\gamma}, \bar{\delta}$.

$$9. \quad A\bar{x} = \lambda\bar{x}$$

$$\begin{bmatrix} 0 & 0 & 1 & 0 \\ 4 & 1 & 0 & 0 \\ 2 & 2 & 1 & 0 \\ 2 & -1 & 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix} = \lambda \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}$$

$$\begin{bmatrix} 3 \\ 6 \\ 9 \\ 12 \end{bmatrix} = \lambda \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix} \Rightarrow \lambda = 3.$$

$$10. \det(\lambda I - A) = \det \begin{pmatrix} (\lambda-2) & -\sqrt{2} \\ -\sqrt{2} & (\lambda-3) \end{pmatrix} = 0$$

$$(\lambda-2)(\lambda-3) - 2 = 0$$

$$\lambda^2 - 5\lambda + 4 = 0$$

$$(\lambda-4)(\lambda-1) = 0$$

$$\lambda = 1; \lambda = 4.$$

$$\lambda = 1:$$

$$\left[\begin{array}{cc|c} -1 & -\sqrt{2} & 0 \\ -\sqrt{2} & -2 & 0 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & \sqrt{2} & 0 \\ 0 & 0 & 0 \end{array} \right] \Rightarrow x = -\sqrt{2}y$$

Let $y = t$

$$\text{S.S: } X = \begin{bmatrix} x \\ y \end{bmatrix} = t \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix}$$

OR

$$x = -\sqrt{2}t$$

$$y = t.$$

$$\lambda = 4:$$

$$\left[\begin{array}{cc|c} 2 & -\sqrt{2} & 0 \\ -\sqrt{2} & 1 & 0 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & -\sqrt{2}/2 & 0 \\ 0 & 0 & 0 \end{array} \right] \Rightarrow$$

$$x = \frac{\sqrt{2}}{2} y$$

OR

$$x = \frac{\sqrt{2}}{2} t$$

$$\text{Let } y = t$$

$$y = t$$

$$X = \begin{bmatrix} x \\ y \end{bmatrix} = t \begin{bmatrix} \frac{\sqrt{2}}{2} \\ 1 \end{bmatrix} = t \begin{bmatrix} \sqrt{2} \\ 2 \end{bmatrix}.$$

$$P = \begin{bmatrix} -\sqrt{2} & \sqrt{2} \\ 1 & 2 \end{bmatrix}.$$

$$P^{-1} = \frac{-1}{3\sqrt{2}} \begin{bmatrix} 2 & -\sqrt{2} \\ -1 & -\sqrt{2} \end{bmatrix}.$$

$$P^{-1}AP = D = \begin{bmatrix} 1 & 0 \\ 0 & 4 \end{bmatrix}.$$