



MATH 205 Final Exam - December 2018

Differential & Integral Calculus II (Concordia University)

CONCORDIA UNIVERSITY
Department of Mathematics & Statistics

Course	Number	Sections
Mathematics	205	All
Examination	Date	Pages
Final	December 2018	2
Instructors:	A. Iovita, U. Mgbemena, I. Pelczer, MD. Rahman	Course Examiners A. Atoyan & H. Proppe
Special	Only approved calculators are allowed	
Instructions:	Show all your work for full marks.	

MARKS

[10] **1. a.** Sketch the graph of the function

$$f(x) = \begin{cases} 3x & x \leq 1 \\ x \frac{|x-5|}{x-5} & 1 < x < 3 \\ -3 & x \geq 3 \end{cases}$$

and find the definite integral $\int_0^5 f(x) dx$ in terms of area
(do not antidifferentiate).

b. Use the Fundamental Theorem of Calculus to calculate the derivative of

$$F(x) = \int_0^{1-x^2} (1-t) e^{-t^2} dt,$$

and determine whether F is increasing or decreasing at $x = 1$.

[15] **2.** Find the following indefinite integrals:

$$(a) \int \frac{\sin^3(x)}{\cos^5(x)} dx \quad (b) \int (2x + x^2) \cos(2x) dx \quad (c) \int \frac{x^2 - 8}{x^2 - 16} dx$$

[18] **3.** Evaluate the following definite integrals (give the exact answers):

$$(a) \int_0^{\ln 2} \frac{e^x}{e^{2x} + 4} dx \quad (b) \int_0^{\pi/4} \frac{\sec^2(x)}{\sqrt{1 + 8 \tan(x)}} dx \quad (c) \int_1^{e^2} x \ln x dx$$

[8] **4.** Evaluate the given improper integral or show that it diverges:

$$(a) \int_e^{\infty} \frac{dx}{x [\ln(x)]^{3/2}} \quad (b) \int_0^1 \frac{dx}{(1-x)^{5/4}}$$

- [16] 5. a. Sketch the curves $y = x^3 - x$ and $y = 3x$, and find the area enclosed.
b. Find the volume of a solid obtained by rotating the region bounded by the curve $y = \sin(x)$ and the x -axis on the interval $0 \leq x \leq \pi$ about the line $y = 2$.
c. Find the exact average value of $f(x) = \sqrt{9 - x^2}$ on the interval $[-3, 3]$.
- [6] 6. Find the limit of the sequence $\{a_n\}$ at $n \rightarrow \infty$ or prove that it does not exist:
(a) $a_n = \frac{3^n + (-3)^n}{4^n}$ (b) $a_n = \ln(1 + 3n + 4n^2) - \ln(8 + 6n + 2n^2)$
- [12] 7. Determine whether the series is divergent or convergent, and if convergent, whether absolutely or conditionally :
(a) $\sum_{n=2}^{\infty} \frac{1}{n \ln(n)}$ (b) $\sum_{n=0}^{\infty} \frac{(-2 + 1/10)^n}{(2 - 1/10)^n}$ (c) $\sum_{n=1}^{\infty} \frac{(-1)^n \sqrt{n^3 + n}}{n^2}$
- [6] 8. Find (a) the radius of convergence, and (b) the interval convergence of the series $\sum_{n=1}^{\infty} \frac{(x - 2)^n}{4^n n^2}$.
- [9] 9. (a) Use the integrability of the power series to express the function $F(x) = \int_0^x \left(\sum_{n=1}^{\infty} n t^{n-1} \right) dt$ as an elementary function (i.e. sum the series for $F(x)$ within the radius of its convergence).
(b) Find the MacLaurin series for the function $f(x) = x^3 \sin(x^2)$. (Hint: start with the series for $\sin z$ then replace z by x^2)
- [5] **Bonus question.** If we know that $\sum_{n=1}^{\infty} a_n$ converges and each $a_n \neq 0$, can anything be said about the series $\sum_{n=1}^{\infty} 1/a_n$ - i.e. does it converge or diverge? Explain your answer.

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.

Final. solutions. Dec. 2018

*Problem 1 :

(1a) sketch the graph


$$f(x) = \begin{cases} 3x & x \leq 1 \\ x \frac{|x-5|}{x-5} & 1 < x < 3 \\ -3 & x \geq 3 \end{cases}$$

we have

$$|x-5| = \begin{cases} x-5 & x \geq 5 \\ -(x-5) & x \leq 5 \end{cases}$$

So

$$f(x) = \begin{cases} 3x & x \leq 1 \\ -x \frac{(x-5)}{(x-5)} & 1 < x < 3 \\ -3 & x \geq 3 \end{cases} \Rightarrow \begin{cases} 3x & \text{if } x \leq 1 \\ -x & \text{if } 1 < x < 3 \\ -3 & \text{if } x \geq 3 \end{cases}$$



Evaluate $\int_0^5 f(x) dx$

$$\int_0^5 f(x) dx = A_1 + A_2 + A_3$$

$$A_1 = \frac{1}{2} \cdot 3 \cdot 1 = \frac{3}{2}$$

$$A_2 = \frac{1}{2} \cdot (1+3) \cdot 2 = 4$$

$$A_3 = 2 \times 3 = 6$$

$$\int_0^5 f(x) dx = \frac{3}{2} - 4 - 6 = -\frac{17}{2}$$

$$\int_0^5 f(x) dx = -\frac{17}{2}$$

(1b) Use FTC to calculate the derivative of $F(x)$

$$F(x) = \int_0^{1-x^2} (1-t) e^{-t^2} dt$$

$$\underline{\text{FTC}} : \left(\int_0^x f(x) \right)' = f(x)$$

So apply FTC on $F(x)$

$$F'(x) = \left(\int_0^{1-x^2} (1-t) e^{-t^2} dt \right)'$$

$$= [1 - (1-x^2)] e^{-(1-x^2)^2} \cdot (1-x^2)'$$

$$= \cancel{x^2} \cancel{e^{-(x^2+1)^2}} \cdot \cancel{(-2x)} \quad x^2 e^{-(1-x^2)^2} (-2x)$$

$$F'(x) = -2x^3 e^{-(x^2+1)^2}$$

$$F'(1) = -2 \cdot 1 e^0 = -2 < 0$$

So $F(x)$ is decreasing at $x=1$

(2)

* Problem 2 : Find the indefinite integrals

$$\begin{aligned} \textcircled{2a} \quad I &= \int \frac{\sin^3(x)}{\cos^5(x)} dx \\ &= \int \frac{\sin^2 x}{\cos^3 x} \cdot \frac{1}{\cos^2(x)} dx \\ &= \int \tan^2(x) \sec^2(x) dx \end{aligned}$$

$$\text{let } u = \tan x \Rightarrow du = \sec^2(x) dx$$

Using substitution u

$$I = \int u^2 du = \frac{1}{3} u^3 + C$$

$$I = \frac{1}{3} \tan^3(x) + C$$

$$\textcircled{2b} \quad I = \int (2x + x^2) \cos(2x) dx$$

Using integration by parts

$$\text{let } u = 2x + x^2 \Rightarrow du = (2 + 2x) dx$$

$$dv = \cos(2x) \Rightarrow v = \frac{\sin(2x)}{2}$$

$$I = uv - \int v du$$

$$= (2x + x^2) \frac{\sin(2x)}{2} - \int \frac{\sin(2x)}{2} (2 + 2x) dx$$

$$= (2x + x^2) \frac{\sin(2x)}{2} - \underbrace{\int (1 + x) \sin(2x) dx}_{I_1}$$

$$\text{let } I_1 = \int (1+x) \sin(2x) dx$$

using integration by parts again

$$\text{let } u = 1+x \Rightarrow du = dx$$

$$dv = \sin(2x) \Rightarrow v = -\frac{\cos(2x)}{2}$$

$$\begin{aligned} I_1 &= -(1+x) \frac{\cos(2x)}{2} - \int -\frac{\cos(2x)}{2} dx \\ &= -\frac{(1+x)}{2} \cos(2x) + \frac{1}{2} \int \cos(2x) dx \\ &= -\frac{(1+x)}{2} \cos(2x) + \frac{1}{4} \sin(2x) + C \end{aligned}$$

So

$$I = (2x+x^2) \frac{\sin(2x)}{2} - \left[-\frac{(x+1)}{2} \cos(2x) + \frac{1}{4} \sin(2x) + C \right]$$

$$I = \frac{(2x+x^2)}{2} \sin(2x) + \frac{(x+1)}{2} \cos(2x) - \frac{1}{4} \sin(2x) + C$$

✓

$$(1c) \quad I = \int \frac{x^2 - 8}{x^2 - 16} dx$$

$$= \int \frac{x^2 - 16 + 8}{x^2 - 16} dx = \int \left[1 + \frac{8}{x^2 - 16} \right] dx$$

$$= x + 8 \int \frac{1}{x^2 - 16} dx$$

$$\text{let } I_1 = \int \frac{1}{x^2 - 16} dx = \int \frac{1}{(x-4)(x+4)} dx$$

$$\frac{1}{(x-4)(x+4)} = \frac{A}{x-4} + \frac{B}{x+4} = \frac{A(x+4) + B(x-4)}{(x-4)(x+4)}$$

$$\Rightarrow 1 = (A+B)x + 4A - 4B$$

$$\Rightarrow \begin{cases} A+B=0 \\ 4A-4B=1 \end{cases} \Rightarrow A = \frac{1}{8}, B = -\frac{1}{8}$$

$$I_1 = \int \frac{1}{(x-4)(x+4)} dx = \int \left[\frac{1}{8(x-4)} - \frac{1}{8(x+4)} \right] dx$$

$$= \frac{1}{8} \ln|x-4| - \frac{1}{8} \ln|x+4|$$

$$\text{So } I = x + 8 \left[\frac{1}{8} \ln|x-4| - \frac{1}{8} \ln|x+4| \right] + C$$

$$I = x + \ln|x-4| - \ln|x+4| + C$$

$$I = x + \ln \frac{|x-4|}{|x+4|} + C$$

~~X~~

* Problem 3: Evaluate the definite integrals

(3a) $I = \int_0^{\ln 2} \frac{e^x}{e^{2x} + 4} dx$

Using substitution

let $u = e^x \Rightarrow du = e^x dx$

$$\int \frac{e^x}{e^{2x} + 4} dx = \int \frac{du}{u^2 + 4} = \int \frac{du}{4 \left[\left(\frac{u}{2} \right)^2 + 1 \right]}$$
$$= \frac{1}{4} \int \frac{du}{\left(\frac{u}{2} \right)^2 + 1}$$

$$= \frac{2}{4} \cdot \tan^{-1} \left(\frac{u}{2} \right) = \frac{1}{2} \tan^{-1} \left(\frac{e^x}{2} \right) + C$$

$$\Rightarrow I = \left. \frac{1}{2} \tan^{-1} \left(\frac{e^x}{2} \right) \right|_0^{\ln 2}$$

$$= \frac{1}{2} \left[\tan^{-1} \left(\frac{e^{\ln 2}}{2} \right) - \tan^{-1} \left(\frac{e^0}{2} \right) \right]$$

$$= \frac{1}{2} \left(\tan^{-1}(1) - \tan^{-1} \left(\frac{1}{2} \right) \right)$$

$$I = \frac{1}{2} \left[\frac{\pi}{4} - \tan^{-1} \left(\frac{1}{2} \right) \right]$$

✗

$$(36) \quad I = \int_0^{\pi/4} \frac{\sec^2(x)}{\sqrt{1+8\tan x}} dx$$

let $u = 1 + 8\tan(x) \Rightarrow du = 8\sec^2(x) dx$

substitute u in to the integral

$$\begin{aligned} \int \frac{\sec^2(x) dx}{\sqrt{1+8\tan x}} &= \int \frac{du}{8\sqrt{u}} = \frac{1}{8} \int u^{-\frac{1}{2}} du \\ &= \frac{2}{8} u^{\frac{1}{2}} = \frac{1}{4} \sqrt{1+8\tan x} \end{aligned}$$

$$\begin{aligned} \text{So } I &= \left[\frac{1}{4} \sqrt{1+8\tan x} \right]_0^{\pi/4} \\ &= \frac{1}{4} \left[\sqrt{1+8\tan \frac{\pi}{4}} - \sqrt{1+8\tan 0} \right] \\ &= \frac{1}{4} (\sqrt{9} - \sqrt{1}) = \frac{1}{2} \end{aligned}$$

$$I = \frac{1}{2}$$

$$(30) \quad I = \int_1^e x \ln(x) dx$$

using integration by parts

let $u = \ln x \Rightarrow du = \frac{1}{x} dx$

$dv = x \Rightarrow v = \frac{x^2}{2}$

$$\begin{aligned} \text{So } \int x \ln(x) &= \frac{x^2}{2} \cdot \ln x - \int \frac{x^2}{2} \cdot \frac{1}{x} dx \\ &= \frac{x^2}{2} \ln x - \frac{1}{2} \int x dx \end{aligned}$$

(7)

$$\int x \ln x = \frac{x^2}{2} \ln(x) - \frac{1}{4} x^2 + C$$

$$I = \left[\frac{x^2}{2} \ln(x) - \frac{1}{4} x^2 \right]_1^{e^2}$$

$$= \left(\frac{e^2}{2} \ln(e) - \frac{1}{4} (e^2)^2 \right) - \left(\frac{1^2}{2} \ln(1) - \frac{1}{4} 1^2 \right)$$

$$I = \frac{e^2}{2} - \frac{e^4}{4} + \frac{1}{4}$$

~~✗~~

* Problem 4 Evaluate the improper integrals

(4a) $I = \int_e^{\infty} \frac{dx}{x(\ln x)^{3/2}}$

since the upper bound is ∞
it is the improper integral type I

$$I = \lim_{b \rightarrow \infty} \int_e^b \frac{dx}{x(\ln x)^{3/2}}$$

Evaluate $\int_e^b \frac{dx}{x(\ln x)^{3/2}}$ by substitution

let $u = \ln x \Rightarrow du = \frac{dx}{x}$

$$\int \frac{dx}{x(\ln x)^{3/2}} = \int \frac{du}{u^{3/2}} = \int u^{-3/2} du$$
$$= -2 u^{-1/2} = -2(\ln x)^{-1/2}$$

$$I = \lim_{b \rightarrow \infty} \left[-\frac{2}{\sqrt{\ln x}} \right]_e^b$$

$$= -2 \lim_{b \rightarrow \infty} \left(\frac{1}{\sqrt{\ln b}} - \frac{1}{\sqrt{\ln e}} \right)$$

$$= -2 \lim_{b \rightarrow \infty} \left(\frac{1}{\sqrt{\ln b}} - 1 \right) = -2(0-1)$$

$I = 2$ conv



$$(46) \quad I = \int_0^1 \frac{dx}{(1-x)^{5/4}}$$

The integrand has a dscont! at $x=1$

$$\text{So } I = \lim_{c \rightarrow 1} \int_0^c \frac{dx}{(1-x)^{5/4}}$$

$$I_1 = \int \frac{dx}{(1-x)^{5/4}}$$

$$\text{let } u = 1-x \Rightarrow du = -dx$$

$$\begin{aligned} I_1 &= - \int \frac{du}{u^{5/4}} = - \int u^{-5/4} du \\ &= 4u^{-1/4} = 4(1-x)^{-1/4} \end{aligned}$$

$$I = \lim_{c \rightarrow 1} \left[\frac{4}{(1-x)^{1/4}} \right]_0^c$$

$$= 4 \lim_{c \rightarrow 1} \left(\frac{1}{(1-c)^{1/4}} - \frac{1}{(1-0)^{1/4}} \right) = 4(\infty - 1)$$

$$I = \infty \quad \text{so } I \text{ div}$$

#

* Problem 5

(5a) Sketch & Find the area enclosed by
 $y = x^3 - x$ and $y = 3x$


* The intersection points

$$x^3 - x = 3x$$

$$x^3 - 4x = 0$$

$$x(x^2 - 4) = x(x-2)(x+2) = 0$$

$$\text{So } x=0, x=2, x=-2$$



$$y = x^3 - x = x(x^2 - 1) = x(x-1)(x+1)$$

the zero of $y = x^3 - x$ is at $x=0, x=1, x=-1$

Since the solid is symmetric with respect to x -axis



$$A = 2A_1 = 2 \int_0^6 [3x - (x^3 - x)] dx$$

$$= 2 \int_0^6 3x - x^3 + x dx = 2 \left(\frac{3}{2}x^2 - \frac{x^4}{4} + \frac{x^2}{2} \right) \Big|_0^6$$

$$A = 8$$

~~X~~

56 Find the volume of the solid obtained by $y = \sin(x)$, x -axis, $0 \leq x \leq \pi$ about the line $y = 2$



Using washer method

$$r(x) = [2 - \sin(x)]$$

$$R(x) = 2$$

$$V = \pi \int_0^\pi [R(x)^2 - r(x)^2] dx$$

$$= \pi \int_0^\pi [2^2 - (2 - \sin(x))^2] dx$$

$$\begin{aligned}
V &= \pi \int_0^{\pi} \left[4 - (4 - 4\sin x + \sin^2 x) \right] dx \\
&= \pi \int_0^{\pi} 4\sin x - \sin^2 x \, dx \\
&= \pi \left[-4\cos(x) \right]_0^{\pi} - \int_0^{\pi} \sin^2 x \, dx \\
&= \pi \left[-4(-1-1) - \int_0^{\pi} \frac{1-\cos 2x}{2} dx \right] \\
&= 8\pi - \frac{\pi}{2} \cdot \int_0^{\pi} (1-\cos 2x) dx \\
&= 8\pi - \frac{\pi}{2} \left(x - \frac{1}{2}\sin 2x \right) \Big|_0^{\pi} \\
&= 8\pi - \frac{\pi}{2} \left(\pi - \frac{1}{2}\sin(2\pi) - 0 + 0 \right) \\
&= 8\pi - \frac{\pi}{2} (\pi)
\end{aligned}$$

$$V = 8\pi - \frac{\pi^2}{2}$$



(5c) Find the ave. of $f(x) = \sqrt{9-x^2}$ on $[-3, 3]$

$$f_{\text{ave}} = \frac{1}{b-a} \int_a^b f(x) dx = \frac{1}{3+3} \int_{-3}^3 \sqrt{9-x^2} dx = \frac{1}{6} \int_{-3}^3 \sqrt{9-x^2} dx$$

$\sqrt{9-x^2}$ is an even function

$$\Rightarrow f_{\text{ave}} = \frac{1}{6} \cdot 2 \int_0^3 \sqrt{9-x^2} dx = \frac{1}{3} \int_0^3 \sqrt{9-x^2} dx$$

$$\text{let } I = \int_0^3 \sqrt{9-x^2} dx$$

using trig. substitution


$$\text{let } x = 3 \sin \theta \Rightarrow dx = 3 \cos \theta d\theta$$

$$\int \sqrt{9-x^2} dx = \int \sqrt{9-9 \sin^2 \theta} \cdot 3 \cos \theta d\theta$$

$$= 9 \int |\cos \theta| \cdot \cos \theta d\theta$$

$$= 9 \int \cos^2 \theta d\theta = 9 \int \frac{1 + \cos 2\theta}{2} d\theta$$

$$= \frac{9}{2} \left(\theta + \frac{1}{2} \sin 2\theta \right) = \frac{9}{2} (\theta + \sin \theta \cos \theta)$$



$$x = 3 \sin \theta$$

$$\Rightarrow \sin \theta = \frac{x}{3}$$

$$\cos \theta = \frac{\sqrt{9-x^2}}{3}$$

$$I = \frac{9}{2} \left(\sin^{-1} \left(\frac{x}{3} \right) + \frac{x}{3} \cdot \frac{\sqrt{9-x^2}}{3} \right) \Bigg|_0^3$$

$$= \frac{9}{2} (\sin^{-1}(1) + 0 - \sin^{-1}(0) + 0)$$

$$= \frac{9}{2} \cdot \frac{\pi}{2} = \frac{9\pi}{4}$$

$$f_{ave} = \frac{1}{3} I = \frac{1}{3} \cdot \frac{9\pi}{4} = \frac{3\pi}{4}$$

$$f_{ave} = \frac{3\pi}{4}$$



*Problem 6 Find the limit


(6a) $a_n = \frac{3^n + (-3)^n}{4^n}$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \left(\frac{3}{4} \right)^n + \left(\frac{-3}{4} \right)^n = 0 + 0 = \boxed{0}$$

(6b) $a_n = \ln(1 + 3n + 4n^2) - \ln(8 + 6n + 2n^2)$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \ln \frac{(1 + 3n + 4n^2)}{(8 + 6n + 2n^2)}$$

$$= \ln \lim_{n \rightarrow \infty} \frac{1 + 3n + 4n^2}{8 + 6n + 2n^2} = \ln \frac{4}{2} = \boxed{\ln 2}$$



*Problem 7: Determine whether the series
conv/div?

7a $\sum_{n=2}^{\infty} \frac{1}{n \ln(n)}$

Using the integral test

let $f(x) = \frac{1}{x \ln(x)}$

$$f'(x) = - \frac{(x \ln(x))'}{(x \ln(x))^2}$$

$$= - \frac{\ln(x) + x \cdot \frac{1}{x}}{(x \ln(x))^2} = - \frac{\ln(x) + 1}{[x \ln(x)]^2}$$

$$\ln(x) > 0 \quad \forall x \geq 2$$

$$\Rightarrow f'(x) < 0 \quad \forall x \geq 2$$

$\Rightarrow f(x)$ is decreasing $\forall x \geq 2$
 $f(x)$ also continuous $\forall x \geq 2$

\Rightarrow consider

$$\int_2^{\infty} \frac{1}{x \ln x} dx = \lim_{b \rightarrow \infty} \int_2^b \frac{1}{x \ln(x)} dx$$

$$\text{let } u = \ln x \Rightarrow du = \frac{dx}{x}$$

$$\int \frac{1}{x \ln x} dx = \int \frac{du}{u} = \ln u = \ln(\ln(x))$$

$$\lim_{b \rightarrow \infty} \int_2^b \frac{1}{x \ln x} dx = \lim_{b \rightarrow \infty} \ln(\ln(x)) \Big|_2^b$$

$$= \lim_{b \rightarrow \infty} [\ln(\ln(b)) - \ln(\ln(2))] = \infty$$

So $\int_2^{\infty} \frac{1}{x \ln x} dx$ div.

by the integral test

$$\sum \frac{1}{n \ln(n)} \text{ div as well}$$

(7b) $\sum_{n=1}^{\infty} \frac{(-2 + \frac{1}{10})^n}{(2 - \frac{1}{10})^n}$

$$= \sum \frac{(-1)^n (2 - \frac{1}{10})^n}{(2 - \frac{1}{10})^n} = \sum (-1)^n$$

$\sum (-1)^n$ div. by the n^{th} term test
for div. series since

$\lim_{n \rightarrow \infty} (-1)^n$ do not exist

$$\Rightarrow \sum \frac{(2 + \frac{1}{10})^n}{(2 - \frac{1}{10})^n} \text{ div}$$

$$(7C) \sum_{n=1}^{\infty} \frac{(-1)^n \sqrt{n^3+n}}{n^2}$$

Using alternating series test

$$\text{let } a_n = \frac{\sqrt{n^3+n}}{n^2}$$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{\sqrt{n^3+n}}{n^2}$$

$$\begin{aligned} &= \lim_{n \rightarrow \infty} \frac{n \sqrt{n + \frac{1}{n}}}{n^2} = \lim_{n \rightarrow \infty} \frac{\sqrt{n + \frac{1}{n}}}{n} \\ &= \lim_{n \rightarrow \infty} \sqrt{\frac{n + \frac{1}{n}}{n^2}} = \lim_{n \rightarrow \infty} \sqrt{\frac{1}{n} + \frac{1}{n^3}} = 0 \end{aligned}$$

$$\lim_{n \rightarrow \infty} a_n = 0$$

$a_n = \frac{\sqrt{n^3+n}}{n^2}$ is a decreasing sequence

$$\text{let } f(x) = \frac{\sqrt{x^3+x}}{x^2} = \sqrt{\frac{1}{x} + \frac{1}{x^3}} = \left(\frac{1}{x} + \frac{1}{x^3} \right)^{\frac{1}{2}}$$

$$\begin{aligned} f'(x) &= \frac{1}{2} \left(\frac{1}{x} + \frac{1}{x^3} \right)^{-\frac{1}{2}} \left(\frac{1}{x} + \frac{1}{x^3} \right)' \\ &= \frac{1}{2 \left(\frac{1}{x} + \frac{1}{x^3} \right)^{\frac{1}{2}}} \left(-\frac{1}{x^2} - 3x^{-4} \right) \end{aligned}$$

$$f'(x) = \frac{1}{2 \left(\frac{1}{x} + \frac{1}{x^3} \right)^{\frac{1}{2}}} \left(-\frac{1}{x^2} - \frac{1}{x^4} \right) \leq 0 \quad \forall x$$

So $f(x)$ is decreasing function

\Rightarrow sequence a_n decreasing as well

By the alternating series test

$$\sum (-1)^n \frac{\sqrt{n^3+n}}{n^2} \text{ conv.}$$

Is it ab. conv. or conditionally conv.

consider $\sum \left| (-1)^n \frac{\sqrt{n^3+n}}{n^2} \right| = \sum \frac{\sqrt{n^3+n}}{n^2}$

Using direct comparison test
We have

$$\forall n, \frac{\sqrt{n^3+n}}{n^2} \geq \frac{\sqrt{n^3}}{n^2} = \frac{n^{\frac{3}{2}}}{n^2} = \frac{1}{n^{2-\frac{3}{2}}} = \frac{1}{n^{\frac{1}{2}}}$$

$\sum \frac{1}{n^{\frac{1}{2}}}$ div by p-series

$\Rightarrow \sum \frac{\sqrt{n^3+n}}{n^2}$ div as well

So.

$$\sum (-1)^n \frac{\sqrt{n^3+n}}{n^2} \text{ conditionally conv}$$

Since the series of the absolute value of the term div



* Problem 8 Find R and I (interval of conv) of

$$\sum_{n=1}^{\infty} \frac{(x-2)^n}{4^n n^2}$$

using ratio test

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{(x-2)^{n+1}}{4^{n+1} (n+1)^2} \cdot \frac{4^n n^2}{(x-2)^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{x-2}{4} \cdot \frac{n^2}{(n+1)^2} \right| = \frac{|x-2|}{4} \end{aligned}$$

the serie conv if $\frac{|x-2|}{4} < 1$

$$\Rightarrow |x-2| < 4$$

$$\text{So } R = 4$$

$$|x-2| < 4$$

$$-4 < x-2 < 4$$

$-2 < x < 6$ is the interval of conv.

• Check endpoints

$$x = -2 \Rightarrow \sum \frac{(x-2)^n}{4^n n^2} = \sum \frac{(-1)^n 4^n}{4^n n^2} = \sum \frac{(-1)^n}{n^2} \text{ conv by alternating series test}$$

$$x = 4 \Rightarrow \sum \frac{(x-2)^n}{4^n n^2} = \sum \frac{4^n}{4^n n^2} = \sum \frac{1}{n^2} \text{ conv. by P-series}$$

So the interval of conv. is

$$-2 \leq x \leq 6$$

~~XX~~

20

* Problem 9: Use integrability of power series

9a

$$F(x) = \int_0^x \left(\sum_{n=1}^{\infty} n t^{n-1} \right) dt$$

By the integrability of power series

$$\Rightarrow F(x) = \sum_{n=1}^{\infty} \int_0^x n t^{n-1} dt$$

$$= \sum_{n=1}^{\infty} \left[n \cdot \frac{1}{n} t^n \right]_0^x$$

$$= \sum_{n=1}^{\infty} x^n = \frac{x}{1-x} \quad \text{where } |x| < 1$$

$$F(x) = \frac{x}{1-x} \quad |x| < 1$$

9b Find the Maclaurin series of $f(x)$
 $f(x) = x^3 \sin(x^2)$

consider $g(t) = \sin t$

$$g(t) = \sin(t) \Rightarrow g(0) = 0$$

$$g'(t) = \cos(t) \Rightarrow g'(0) = 1$$

$$g''(t) = -\sin(t) \Rightarrow g''(0) = 0$$

$$g^{(3)}(t) = -\cos(t) \Rightarrow g^{(3)}(0) = -1$$

$$g^{(4)}(t) = \sin(t) \Rightarrow g^{(4)}(0) = 0$$

The patterns repeat here

the Maclaurin series is

$$\begin{aligned}
 \sin(t) &= f(0) + \frac{f'(0)}{1!}t + \frac{f''(0)}{2!}t^2 + \frac{f'''(0)}{3!}t^3 + \dots + \frac{f^{(n)}(0)}{n!}t^n \\
 &= 0 + t + 0 - \frac{t^3}{3!} + 0 + \frac{t^5}{5!} + 0 - \frac{t^7}{7!} + \dots + \frac{(-1)^n t^{2n+1}}{(2n+1)!} \\
 &= t - \frac{t^3}{3!} + \frac{t^5}{5!} - \frac{t^7}{7!} + \dots + \frac{(-1)^n t^{2n+1}}{(2n+1)!}
 \end{aligned}$$

$$\sin t = \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!}$$

$$\Rightarrow \sin(x^2) = \sum_{n=0}^{\infty} \frac{(-1)^n (x^2)^{2n+1}}{(2n+1)!}$$

$$\begin{aligned}
 \Rightarrow x^3 \sin(x^2) &= x^3 \sum_{n=0}^{\infty} \frac{(-1)^n (x^2)^{2n+1}}{(2n+1)!} \\
 &= \sum_{n=0}^{\infty} \frac{(-1)^n x^{4n+2} \cdot x^3}{(2n+1)!}
 \end{aligned}$$

$$x^3 \sin(x^2) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{4n+5}}{(2n+1)!}$$

#