



Solved Final Exam April 2017

Vectors and Matrices (Concordia University)

CONCORDIA UNIVERSITY
Department of Mathematics and Statistics

Course	Number	Section(s)
Mathematics	204	All - Except EC

Examination	Date	Pages
Final	April 2017	3

Instructors	Course Examiner
C. L. Santana, F. Mobasheramini, T. Hughes	A. Kokotov

Special Instructions: Only approved calculators are allowed !

1. [10] Solve the system of equations

$$\begin{cases} x_1 + 2x_2 - 3x_3 + x_4 = 1 \\ 3x_1 - x_2 - x_3 + 2x_4 = 3 \\ 2x_1 + 6x_2 + 4x_3 - x_4 = 11 \\ -5x_1 + x_2 - x_3 + 7x_4 = 2 \\ 2x_1 - 4x_2 + x_3 + 5x_4 = 4 \end{cases}$$

2. [10] Solve the following equation for (3×3) matrix X:

$$X - \begin{pmatrix} 2 & 1 & 1 \\ 0 & 3 & 1 \\ 1 & 0 & 4 \end{pmatrix} X = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$$

3. [10] For which a the following matrix

$$\begin{pmatrix} a+1 & 1 & 1 & 1 \\ a+2 & a & 2 & 2 \\ a+2 & 1 & a+2 & 2 \\ 2a+3 & 1 & 1 & a+3 \end{pmatrix}$$

is not invertible?

4. [10]

Let the planes α and β be given by equations

$$(\alpha) \quad x + y - z = 1$$

and

$$(\beta) \quad x - y + 2z = -1$$

and let P be the point $(2, 1, 1)$.

(A) Write down the equation of the plane passing through the point P and the line \mathcal{L} of intersection of α and β .

(B) Find the coordinates of the intersection of the line \mathcal{L} and the plane

$$2x - 5y + z = 4$$

5. [10]

Find a for which all three vectors $(1, 2, 3)$, $(1, 2 + a, 6)$ and $(1, 10, a - 1)$ are parallel to the same plane.

6. [10] Find the value of the determinant

$$\begin{vmatrix} 1 & 1 & 2 & 1 & 4 \\ 2 & 3 & 3 & 2 & 3 \\ 0 & 0 & 1 & 1 & 0 \\ 2 & 2 & 0 & 4 & 5 \\ 7 & 7 & 5 & 6 & 9 \end{vmatrix}$$

7. [10] Find a basis of the subspace of \mathbb{R}^4 generated by the vectors $(1, 1, 1, 1)$, $(1, 2, 3, 4)$, $(2, 3, 4, 5)$, $(3, 5, 7, 9)$ and $(4, 6, 8, 10)$.
8. [10] Find a basis for the solution space of the homogeneous system

$$\begin{cases} x_1 + x_2 + 3x_3 + x_4 - x_5 = 0 \\ 2x_1 + 3x_2 - x_3 + 2x_4 - x_6 = 0 \\ 4x_1 + 5x_2 + 5x_3 + 4x_4 - 2x_5 - x_6 = 0 \end{cases}$$

9. [10] Find the eigenvalues and eigenvectors of the matrix

$$A = \begin{pmatrix} 0 & 1 & 0 \\ -4 & 4 & 0 \\ -2 & 1 & 2 \end{pmatrix}$$

Is A diagonalizable?

10. [10] Find an invertible 2×2 matrix P such that the matrix

$$P^{-1} \begin{pmatrix} 1 & \sqrt{6} \\ \sqrt{6} & 2 \end{pmatrix} P$$

is diagonal.

¹The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.

MATH. 204.

FINAL EXAM SOLUTIONS.

APRIL, 2017.

1. $x = y = z = w = 1.$

2.

$$X = \begin{bmatrix} -1 & -0.6 & -1 \\ 0 & -0.6 & 0 \\ 0 & 0.2 & 0 \end{bmatrix}$$

5. The scalar triple product of the three vectors must be 0.

Let: \bar{u} , \bar{v} , \bar{w} be the three vectors.

$$(\bar{u} \times \bar{v}) \cdot \bar{w} = 0.$$

$$\begin{vmatrix} 1 & 2 & 3 \\ 1 & (2+a) & 6 \end{vmatrix} = (6 - 3a, -3, a).$$

$$(1, 10, a - 1) \cdot (6 - 3a, -3, a) = a^2 - 4a - 24.$$

$$a^2 - 4a - 24 = 0$$

$$a = 2 \pm 2\sqrt{7}.$$

$$3. \left| \begin{array}{cccc} (a+1) & 1 & 1 & 1 \\ (a+2) & a & 2 & 2 \\ (a+2) & 1 & (a+2) & 2 \\ (2a+3) & 1 & 1 & (a+3) \end{array} \right| \begin{array}{l} -aR_1 + R_2 \rightarrow R_2 \\ \hline -R_1 + R_3 \rightarrow R_3 \\ -R_1 + R_4 \rightarrow R_4 \end{array}$$

$$\left| \begin{array}{cccc} (a+1) & 1 & 1 & 1 \\ (2-a^2) & 0 & (2-a) & (2-a) \\ 1 & 0 & (a+1) & 1 \\ (a+2) & 0 & 0 & (a+2) \end{array} \right| =$$

$$(1) \left| \begin{array}{ccc} (2-a^2) & (2-a) & (2-a) \\ 1 & (a+1) & 1 \\ (a+2) & 0 & (a+2) \end{array} \right| =$$

$$-a^4 - 2a^3 + a^2 + 2a = 0$$

$$-a(a^3 + 2a^2 - a - 2) = 0$$

$$-a(a+1)(a-1)(a+2) = 0$$

$$a = 0; -1; 1; -2.$$

Matrix is not invertible when determinant equals 0.

$$4. \quad \alpha: x + y - z = 1 \quad P(2, 1, 1).$$

$$\beta: x - y + 2z = -1$$

A.

$$\left[\begin{array}{ccc|c} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \end{array} \right] \xrightarrow{-R_1 + R_2 \rightarrow R_2} \left[\begin{array}{ccc|c} 1 & 0 & -1 & 1 \\ 0 & -2 & 3 & -2 \end{array} \right]$$

$$\xrightarrow{-\frac{1}{2}R_2 \rightarrow R_2} \left[\begin{array}{ccc|c} 1 & 1 & -1 & 1 \\ 0 & 1 & -\frac{3}{2} & 1 \end{array} \right] \xrightarrow{-R_2 + R_1 \rightarrow R_1}$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 1/2 & 0 \\ 0 & 1 & -3/2 & 1 \end{array} \right] \Rightarrow \begin{aligned} x &= -\frac{1}{2}z \\ y &= \frac{3}{2}z + 1 \end{aligned}$$

$$\text{Let } z = t$$

$$x = -\frac{1}{2}t$$

$$y = 1 + \frac{3}{2}t$$

$$\vec{X} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1/2 \\ 3/2 \\ 1 \end{bmatrix}_x$$

$$\therefore L: \vec{X} = (0, 1, 0) + \left(-\frac{1}{2}, \frac{3}{2}, 1\right)t_s = Q + \vec{a}t.$$

Plane:

$$\vec{QP} = (2-0, 1-1, 1-0) = (2, 0, 1).$$

$$\vec{a} = \left(-\frac{1}{2}, \frac{3}{2}, 1\right) = (-1 \ 3 \ 2).$$

$$\vec{a} \times \vec{QP} = \vec{n} = \begin{vmatrix} -1 & 3 & 2 \\ 2 & 0 & 1 \end{vmatrix} = (3, +5, -6).$$

$$X = (x, y, z).$$

$$\text{Plane: } \vec{n} \cdot (X - P) = 0$$

$$(3, 5, -6) \cdot (x-2, y-1, z-1) = 0$$

$$3(x-2) + 5(y-1) + (-6)(z-1) = 0$$

$$3x + 5y - 6z = 5.$$

$$B. \quad 2x - 5y + z = 4.$$

$$L: x = -\frac{1}{2}t$$

$$y = 1 + \frac{3}{2}t$$

$$z = t.$$

$$2\left(-\frac{1}{2}t\right) - 5\left(1 + \frac{3}{2}t\right) + (t) = 4$$

$$-t - 5 - \frac{15}{2}t + t = 4$$

$$-\frac{15}{2}t = 9$$

$$-15t = 18$$

$$t = -\frac{18}{15} = -\frac{6}{5}$$

\therefore , Point of intersection: $\left(\frac{3}{5}, -\frac{4}{5}, -\frac{6}{5}\right)$.

$$6. \begin{vmatrix} 1 & 1 & 2 & 1 & 4 \\ 2 & 3 & 3 & 2 & 3 \\ 0 & 0 & 1 & 1 & 0 \\ 2 & 2 & 0 & 4 & 5 \\ 7 & 7 & 5 & 6 & 9 \end{vmatrix} \xrightarrow{-C_3 + C_4 \rightarrow C_4}$$

$$\begin{vmatrix} 1 & 1 & 2 & -1 & 4 \\ 2 & 3 & 3 & -1 & 3 \\ 0 & 0 & 1 & 0 & 0 \\ 2 & 2 & 0 & 4 & 5 \\ 7 & 7 & 5 & 1 & 9 \end{vmatrix} = (1) \begin{vmatrix} 1 & 1 & -1 & 4 \\ 2 & 3 & -1 & 3 \\ 2 & 2 & 4 & 5 \\ 7 & 7 & 1 & 9 \end{vmatrix} \begin{array}{l} -2R_1 + R_2 \rightarrow R_2 \\ -2R_1 + R_3 \rightarrow R_3 \\ -7R_1 + R_4 \rightarrow R_4 \end{array}$$

$$\begin{vmatrix} 1 & 1 & -1 & 4 \\ 0 & 1 & 1 & -5 \\ 0 & 0 & 6 & -3 \\ 0 & 0 & 8 & -19 \end{vmatrix} = (1) \begin{vmatrix} 1 & 1 & -5 \\ 0 & 6 & -3 \\ 0 & 8 & -19 \end{vmatrix} =$$

$$(1) \begin{vmatrix} 6 & -3 \\ 8 & -19 \end{vmatrix} = -90.$$

7.

$$\begin{bmatrix} 1 & 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 5 & 6 \\ 1 & 3 & 4 & 7 & 8 \\ 1 & 4 & 5 & 9 & 10 \end{bmatrix} \begin{array}{l} -R_1 + R_2 \rightarrow R_2 \\ -R_1 + R_3 \rightarrow R_3 \\ \hline -R_1 + R_4 \rightarrow R_4 \end{array}$$

$$\begin{bmatrix} 1 & 1 & 2 & 3 & 4 \\ 0 & 1 & 1 & 2 & 2 \\ 0 & 2 & 2 & 4 & 4 \\ 0 & 3 & 3 & 6 & 6 \end{bmatrix} \begin{array}{l} -R_2 + R_1 \rightarrow R_1 \\ -2R_2 + R_3 \rightarrow R_3 \\ \hline -3R_2 + R_4 \rightarrow R_4 \end{array}$$

$$\begin{bmatrix} 1 & 0 & 1 & 1 & 2 \\ 0 & 1 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Basis $\bar{v}_1; \bar{v}_2$.

$$8. \begin{array}{cccccc} x & y & z & u & v & w \\ \left[\begin{array}{cccccc} 1 & 1 & 3 & 1 & -1 & 0 \\ 2 & 3 & -1 & 2 & 0 & -1 \\ 4 & 5 & 5 & 4 & -2 & -1 \end{array} \right] \begin{array}{l} -2R_1 + R_2 \rightarrow R_2 \\ \hline -4R_1 + R_3 \rightarrow R_3 \end{array} \end{array}$$

$$\begin{array}{cccccc} \left[\begin{array}{cccccc} 1 & 1 & 3 & 1 & -1 & 0 \\ 0 & 1 & -7 & 0 & 2 & -1 \\ 0 & 1 & -7 & 0 & 2 & -1 \end{array} \right] \begin{array}{l} -R_2 + R_3 \rightarrow R_3 \\ \hline -R_2 + R_1 \rightarrow R_1 \end{array} \end{array}$$

$$\begin{bmatrix} 1 & 0 & 10 & 1 & -3 & 1 \\ 0 & 1 & -7 & 0 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \Rightarrow$$

$$x = -10z - u + 3v - w$$

$$y = +7z - 2v + w$$

$$\text{Let } z = s$$

$$u = t$$

$$v = a$$

$$w = b$$

$$X = \begin{bmatrix} x \\ y \\ z \\ u \\ v \\ w \end{bmatrix} = s \begin{bmatrix} -10 \\ 7 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + a \begin{bmatrix} 3 \\ -2 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + b \begin{bmatrix} -1 \\ 1 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}.$$

\therefore Basis: $\begin{bmatrix} -10 \\ 7 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}; \begin{bmatrix} -1 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}; \begin{bmatrix} 3 \\ -2 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}; \begin{bmatrix} -1 \\ 1 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}.$

$$9. \det(\lambda I - A) = 0$$

$$\det \left(\begin{bmatrix} \lambda & -1 & 0 \\ 4 & (\lambda-4) & 0 \\ 2 & -1 & (\lambda-2) \end{bmatrix} \right) = 0$$

$$\lambda^3 - 6\lambda^2 + 12\lambda - 8 = 0$$

$$(\lambda - 2)^3 = 0$$

$$\lambda = 2.$$

$$\begin{pmatrix} 2 & -1 & 0 \\ 4 & -2 & 0 \\ 2 & -1 & 0 \end{pmatrix} \longrightarrow \begin{pmatrix} 1 & -\frac{1}{2} & 0 \\ 4 & -2 & 0 \\ 2 & -1 & 0 \end{pmatrix} \longrightarrow$$

$$\begin{pmatrix} 1 & -\frac{1}{2} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \begin{aligned} x &= \frac{1}{2}y \\ \text{Let } y &= s \\ z &= t \end{aligned}$$

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ 1 \\ 0 \end{bmatrix} s + \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} t. \quad \text{Eigenvectors:}$$

$$\begin{bmatrix} \frac{1}{2} \\ 1 \\ 0 \end{bmatrix}; \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

A is not diagonalizable since the three λ 's are not distinct.

10.

$$\det(\lambda I - A) = 0$$

$$\det\left(\begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} - \begin{pmatrix} 1 & \sqrt{6} \\ \sqrt{6} & 2 \end{pmatrix}\right) = 0$$

$$\det\left(\begin{bmatrix} (\lambda-1) & -\sqrt{6} \\ -\sqrt{6} & (\lambda-2) \end{bmatrix}\right) = 0$$

$$\lambda^2 - 3\lambda - 4 = 0$$

$$(\lambda-4)(\lambda+1) = 0$$

$$\lambda = -1; 4.$$

$$\lambda = -1:$$

$$\begin{pmatrix} -2 & -\sqrt{6} \\ -\sqrt{6} & -3 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & \frac{\sqrt{6}}{2} \\ 0 & 0 \end{pmatrix} \Rightarrow x = -\frac{\sqrt{6}}{2} y$$

$$\text{Let } y = t$$

$$X = \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -\frac{\sqrt{6}}{2} \\ 1 \end{bmatrix} t.$$

$$\lambda = 4:$$

$$\begin{pmatrix} 3 & -\sqrt{6} \\ -\sqrt{6} & 2 \end{pmatrix} \longrightarrow \begin{pmatrix} 1 & -\frac{\sqrt{6}}{3} \\ 0 & 0 \end{pmatrix} \Rightarrow x = \frac{\sqrt{6}}{3} y$$

Let $y = t$

$$X = \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{6}}{3} \\ 1 \end{bmatrix} t.$$

$$P = \begin{bmatrix} -\frac{\sqrt{6}}{2} & \frac{\sqrt{6}}{3} \\ 1 & 1 \end{bmatrix}; \quad P^{-1} = \frac{-\sqrt{6}}{5} \begin{bmatrix} 1 & -\frac{\sqrt{6}}{3} \\ -1 & -\frac{\sqrt{6}}{2} \end{bmatrix}.$$

$$P^{-1}AP = \begin{pmatrix} -1 & 0 \\ 0 & 4 \end{pmatrix}.$$