

# Computational Physics II

## FYS-4411

Gullik Vetvik Killie  
Håkon S.B. Mørk  
Jose Emeilio Ruiz Navarro

February 28, 2015

### Abstract

Fill in abstract

## 1 Introduction

## 2 Methods

### 2.1 Monte Carlo of the Helium Atom

In a quantum mechanical system the energy is given by the expectation value of the Hamiltonian.

$$E[H] = \langle \Psi | \quad (1)$$

### 2.2 Derivation of local energies

The local energy of is dependant on the Hamiltonian and the wavefunction describing the system, the Hamiltonian incorporates both a kinetic energy part given by  $\frac{\nabla^2}{2}$  for each particle and a potential energy part given by  $\frac{Z}{r_i}$  and  $\frac{1}{r_{ij}}$ , where  $Z$  is the charge of the center,  $r_i$  is the distance for electron  $i$  to the atom center and  $r_{ij}$  is the distance between electron  $l$  and  $m$ . Then the local energy is given by the following:

$$E_L = \sum_{i,i < j} \frac{1}{\Psi_T(\mathbf{r}_i, \mathbf{r}_{ij})} \hat{H} \Psi_T(\mathbf{r}_i, \mathbf{r}_{ij}) \quad (2)$$

$$= \sum_{i,i < j} \frac{1}{\Psi_T(\mathbf{r}_i, \mathbf{r}_{ij})} \left( -\frac{\nabla_i^2}{2} - \frac{Z}{r_i} - \frac{Z}{r_j} + \frac{1}{r_{ij}} \right) \Psi_T(\mathbf{r}_i, \mathbf{r}_{ij}) \quad (3)$$

$$= \sum_{i,i < j} -\frac{1}{2\Psi_T} (\nabla_i^2 \Psi_T) - \frac{Z}{r_i} - \frac{Z}{r_j} + \frac{1}{r_{ij}} \quad (4)$$

Let us change derivation variables:

$$-\frac{1}{2\Psi_T} (\nabla_i^2 \Psi_T) = \sum_{m=1}^3 -\frac{1}{2\Psi_T} \left( \frac{\partial^2 \Psi_T}{\partial x_m^2} \right)_i \quad (5)$$

$$= \sum_{m=1}^3 -\frac{1}{2\Psi_T} \left( \frac{\partial}{\partial x_m} \left( \frac{\partial \Psi_T}{\partial r_i} \frac{\partial r_i}{\partial x_m} \right) \right)_i \quad (6)$$

Since  $r_i = (x_1^2 + x_2^2 + x_3^2)^{1/2}$  then  $\frac{\partial r_i}{\partial x_m} = \frac{\partial (x_1^2 + x_2^2 + x_3^2)^{1/2}}{\partial x_m} = \frac{x_m}{r_i}$

$$= \sum_{m=1}^3 -\frac{1}{2\Psi_T} \left( \frac{\partial}{\partial x_m} \left( \frac{\partial \Psi_T}{\partial r_i} \frac{x_m}{r_i} \right) \right)_i \quad (7)$$

$$= \sum_{m=1}^3 -\frac{1}{2\Psi_T} \left( \frac{\partial^2 \Psi_T}{\partial x_m \partial r_i} \frac{x_m}{r_i} + \frac{\partial \Psi_T}{\partial r_i} \frac{\partial}{\partial x_m} \left( \frac{x_m}{r_i} \right) \right)_i \quad (8)$$

The term  $\frac{\partial}{\partial x_m} \left( \frac{x_m}{r_i} \right)$  becomes for the different values for  $m$ ,  $\frac{\partial}{\partial x_1} \left( \frac{x_1}{(x_1^2 + x_2^2 + x_3^2)^{1/2}} \right) = \frac{x_2^2 + x_3^2}{r_i^3}$  so all the values for  $m$  term it should sum up to  $\frac{2(x_1^2 + x_2^2 + x_3^2)}{r_i^3}$

$$= -\frac{1}{2\Psi_T} \left( \frac{\partial^2 \Psi_T}{\partial r_i^2} \frac{x_1^2 + x_2^2 + x_3^2}{r_i^2} + \frac{\partial \Psi_T}{\partial r_i} \frac{2(x_1^2 + x_2^2 + x_3^2)}{r_i^3} \right)_i \quad (9)$$

$$= -\frac{1}{2\Psi_T} \left( \frac{\partial^2 \Psi_T}{\partial r_i^2} + \frac{\partial \Psi_T}{\partial r_i} \frac{2}{r_i} \right) \quad (10)$$

Then the local energy becomes:

$$E_L = \sum_{i,i < j} -\frac{1}{2\Psi_T} \left( \frac{\partial^2 \Psi_T}{\partial r_i^2} + \frac{\partial \Psi_T}{\partial r_i} \frac{2}{r_i} \right) - \frac{Z}{r_i} - \frac{Z}{r_j} + \frac{1}{r_{ij}} \quad (11)$$

### 2.2.1 Helium: Simple trialfunction

The simple version of the trial function is only dependant on one parameter  $\alpha$  and does not take into account interaction between the two electrons, it is of the form

$$\Psi_T(\mathbf{r}_1, \mathbf{r}_2) = \exp\{-\alpha(r_1 + r_2)\}$$

Let us set this trialfunction into the equation for the local energy (11).

$$E_L = \sum_{i,i < j} -\frac{1}{2\Psi_T} \left( \frac{\partial^2 e^{-\alpha(r_i + r_j)}}{\partial r_i^2} + \frac{\partial e^{-\alpha(r_i + r_j)}}{\partial r_i} \frac{2}{r_i} \right) - \frac{Z}{r_i} - \frac{Z}{r_j} + \frac{1}{r_{ij}} \quad (12)$$

$$E_L = -\frac{1}{2\Psi_T} \sum_{i=1}^2 \left( \alpha^2 - \alpha \frac{2}{r_i} \right) \Psi_T - \frac{Z}{r_i} + \frac{1}{r_{ij}} \quad (13)$$

$$E_L = -\alpha^2 + (\alpha - Z) \left( \frac{1}{r_1} + \frac{1}{r_2} \right) + \frac{1}{r_{12}} \quad (14)$$

## 3 Results and discussion

## 4 Conclusions and perspectives

The local energy for the simple trialfunction

$$Z\left(-\frac{1}{r_2}-\frac{1}{r_1}\right)-\alpha^2+\alpha\left(\frac{1}{r_2}+\frac{1}{r_1}\right)+\frac{1}{r_{12}}$$