

# Fourier series: basic concepts

Øyvind Ryan

Jan 13, 2017

## Square-integrable functions, Definition 1.6

The set of continuous, real functions defined on an interval  $[0, T]$  is denoted  $C[0, T]$ .

A real function  $f$  defined on  $[0, T]$  is said to be square integrable if  $f^2$  is Riemann-integrable, i.e., if the Riemann integral of  $f^2$  on  $[0, T]$  exists,

$$\int_0^T f(t)^2 dt < \infty.$$

The set of all square integrable functions on  $[0, T]$  is denoted  $L^2[0, T]$ .

Both  $L^2[0, T]$  and  $C[0, T]$  are vector spaces. Moreover, if the two functions  $f$  and  $g$  lie in  $L^2[0, T]$  (or in  $C[0, T]$ ), then the product  $fg$  is Riemann-integrable (or in  $C[0, T]$ ). Moreover, both spaces are inner product spaces with inner product defined by

$$\langle f, g \rangle = \frac{1}{T} \int_0^T f(t)g(t) dt,$$

and associated norm

$$\|f\| = \sqrt{\frac{1}{T} \int_0^T f(t)^2 dt}.$$

Let  $V_{N,T}$  be the subspace of  $C[0, T]$  spanned by the set of functions given by

$$\mathcal{D}_{N,T} = \{1, \cos(2\pi t/T), \cos(2\pi 2t/T), \dots, \cos(2\pi Nt/T), \\ \sin(2\pi t/T), \sin(2\pi 2t/T), \dots, \sin(2\pi Nt/T)\}.$$

The space  $V_{N,T}$  is called the  *$N$ 'th order Fourier space*. The  $N$ th-order Fourier series approximation of  $f$ , denoted  $f_N$ , is defined as the best approximation of  $f$  from  $V_{N,T}$  with respect to the inner product defined by (1.3).

## Fourier coefficients, Theorem 1.9

The set  $\mathcal{D}_{N,T}$  is an orthogonal basis for  $V_{N,T}$ . In particular, the dimension of  $V_{N,T}$  is  $2N + 1$ , and if  $f$  is a function in  $L^2[0, T]$ , we denote by  $a_0, \dots, a_N$  and  $b_1, \dots, b_N$  the coordinates of  $f_N$  in the basis  $\mathcal{D}_{N,T}$ , i.e.

$$f_N(t) = a_0 + \sum_{n=1}^N (a_n \cos(2\pi nt/T) + b_n \sin(2\pi nt/T)).$$

The  $a_0, \dots, a_N$  and  $b_1, \dots, b_N$  are called the (real) Fourier coefficients of  $f$ , and they are given by

$$a_0 = \langle f, 1 \rangle = \frac{1}{T} \int_0^T f(t) dt,$$

$$a_n = 2 \langle f, \cos(2\pi nt/T) \rangle = \frac{2}{T} \int_0^T f(t) \cos(2\pi nt/T) dt \quad \text{for } n \geq 1,$$

$$b_n = 2 \langle f, \sin(2\pi nt/T) \rangle = \frac{2}{T} \int_0^T f(t) \sin(2\pi nt/T) dt \quad \text{for } n \geq 1.$$

## Convergence of Fourier series Theorem 1.26



Suppose that  $f$  is periodic with period  $T$ , and that

- $f$  has a finite set of discontinuities in each period.
- $f$  contains a finite set of maxima and minima in each period.
- $\int_0^T |f(t)| dt < \infty$ .

Then we have that  $\lim_{N \rightarrow \infty} f_N(t) = f(t)$  for all  $t$ , except at those points  $t$  where  $f$  is not continuous.


# Fourier series of the square wave

```
t = linspace(0, T, 100);  
y = zeros(size(t));  
for n = 1:2:19  
    y = y + (4/(n*pi))*sin(2*pi*n*t/T);  
end  
plot(t,y)
```



**Figure:** The Fourier series  $(f_s)_{20}$  and the values for the first 100 Fourier coefficients  $b_n$ .

# Fourier series of the triangle wave



**Figure:** The Fourier series  $(f_t)_{20}$  and the values for the first 100 Fourier coefficients  $a_n$ .



- With  $N = 1$ , the Fourier series of both the square and the triangle wave are pure tones with frequency  $\nu = 1/T$ . Largest term.
- $f_s$  is anti-symmetric about 0, and its Fourier series was a sine series.
- $f_t$  is symmetric about 0, and its Fourier series was a cosine series.
- $f_t$  is continuous, and  $f_s$  is not.
- The Fourier series of  $f_t$  converged faster than that of  $f_s$ .

## Symmetry and antisymmetry, Theorem 1.10

- If  $f$  is antisymmetric about 0 ( $f(-t) = -f(t)$  for all  $t$ ), then  $a_n = 0$ , so the Fourier series is actually a sine-series.
- If  $f$  is symmetric about 0 ( $f(-t) = f(t)$  for all  $t$ ), then  $b_n = 0$ , so the Fourier series is actually a cosine-series.

We define the set of functions

$$\mathcal{F}_{N,T} = \{e^{-2\pi i Nt/T}, e^{-2\pi i(N-1)t/T}, \dots, e^{-2\pi it/T}, \\ 1, e^{2\pi it/T}, \dots, e^{2\pi i(N-1)t/T}, e^{2\pi i Nt/T}\},$$

and call this the order  $N$  *complex Fourier basis* for  $V_{N,T}$ .

# Complex vector spaces and inner products

For general complex functions we extend the definition of the inner product as

$$\langle f, g \rangle = \frac{1}{T} \int_0^T f \bar{g} dt.$$

The associated norm is

$$\|f\| = \sqrt{\frac{1}{T} \int_0^T |f(t)|^2 dt}. \quad (1)$$

# Complex Fourier coefficients, Theorem 1.12


We denote by  $y_{-N}, \dots, y_0, \dots, y_N$  the coordinates of  $f_N$  in the basis  $\mathcal{F}_{N,T}$ , i.e.

$$f_N(t) = \sum_{n=-N}^N y_n e^{2\pi i n t / T}.$$

The  $y_n$  are called the complex Fourier coefficients of  $f$ , and they are given by.

$$y_n = \langle f, e^{2\pi i n t / T} \rangle = \frac{1}{T} \int_0^T f(t) e^{-2\pi i n t / T} dt.$$

## Distribution of Fourier coefficients in Example 1.24



**Figure:** Plot of  $|y_n|$  when  $f(t) = e^{2\pi it/T_2}$ , and  $T_2 > T$ . Left:  $T/T_2 = 0.5$ . Right:  $T/T_2 = 0.9$ .

we use the notation  $f \rightarrow y_n$  to indicate that  $y_n$  is the  $n$ 'th (complex) Fourier coefficient of  $f(t)$ . The functions  $1$ ,  $e^{2\pi int/T}$ , and  $\chi_{-a,a}$  have the Fourier coefficients

$$\begin{aligned}1 &\rightarrow \mathbf{e}_0 = (1, 0, 0, 0, \dots,) \\ e^{2\pi int/T} &\rightarrow \mathbf{e}_n = (0, 0, \dots, 1, 0, 0, \dots) \\ \chi_{-a,a} &\rightarrow \frac{\sin(2\pi na/T)}{\pi n}.\end{aligned}$$

## Fourier series properties, Theorem 1.15

The mapping  $f \rightarrow y_n$  is linear: if  $f \rightarrow x_n$ ,  $g \rightarrow y_n$ , then

$$af + bg \rightarrow ax_n + by_n$$

For all  $n$ . Moreover, if  $f$  is real and periodic with period  $T$ , the following properties hold:

- ❶  $y_n = \overline{y_{-n}}$  for all  $n$ .
- ❷ If  $f(t) = f(-t)$  (i.e.  $f$  is symmetric), then all  $y_n$  are real, so that  $b_n$  are zero and the Fourier series is a cosine series.
- ❸ If  $f(t) = -f(-t)$  (i.e.  $f$  is antisymmetric), then all  $y_n$  are purely imaginary, so that the  $a_n$  are zero and the Fourier series is a sine series.
- ❹ If  $g(t) = f(t - d)$  (i.e.  $g$  is the function  $f$  delayed by  $d$ ) and  $f \rightarrow y_n$ , then  $g \rightarrow e^{-2\pi ind/T} y_n$ .
- ❺ If  $g(t) = e^{2\pi idt/T} f(t)$  with  $d$  an integer, and  $f \rightarrow y_n$ , then  $g \rightarrow y_{n-d}$ .
- ❻ Let  $d$  be a number. If  $f \rightarrow y_n$ , then  $f(d + t) = f(d - t)$  for all  $t$  if and only if the argument of  $y_n$  is  $-2\pi nd/T$  for all  $n$ .



# Convergence of Fourier series 1

**Corollary 1.17** If the complex Fourier coefficients of  $f$  are  $y_n$  and  $f$  is differentiable, then the Fourier coefficients of  $f'(t)$  are  $\frac{2\pi in}{T} y_n$ .

Turning this around: the Fourier coefficients of  $f(t)$  are  $T/(2\pi in)$  times those of  $f'(t)$ , when  $f$  is differentiable. In other words, the Fourier coefficients of a function which is many times differentiable decay to zero very fast.

**Observation 1.18** The Fourier series converges quickly when the function is many times differentiable.

## Convergence of Fourier series 2

**Idea 1.19** Assume that  $f$  is continuous on  $[0, T)$ . Can we construct another periodic function which agrees with  $f$  on  $[0, T]$ , and which is both continuous and periodic (maybe with period different from  $T$ )?

If this is possible the Fourier series of the new function could produce better approximations for  $f$ . It turns out that the following extension strategy does the job:



**Definition 1.20:** Let  $f$  be a function defined on  $[0, T]$ . By the *symmetric extension* of  $f$ , denoted  $\check{f}$ , we mean the function defined on  $[0, 2T]$  by

$$\check{f}(t) = \begin{cases} f(t), & \text{if } 0 \leq t \leq T; \\ f(2T - t), & \text{if } T < t \leq 2T. \end{cases}$$

## Convergence of Fourier series 3

The following holds:

If  $f$  is continuous on  $[0, T]$ , then  $\check{f}$  is continuous on  $[0, 2T]$ , and  $\check{f}(0) = \check{f}(2T)$ . If we extend  $\check{f}$  to a periodic function on the whole real line (which we also will denote by  $\check{f}$ ), this function is continuous, agrees with  $f$  on  $[0, T]$ , and is a symmetric function.



**Figure:** Two different extensions of  $f$  to a periodic function on the whole real line. Periodic extension (left) and symmetric extension (right).