

# Dynamic Programming Chain Matrix Multiply

(Class 20)

- Let  $m[i, j]$  denote the minimum number of multiplications needed to compute  $A_{i...j}$ .
- Where  $1 \leq i \leq j \leq n$  ( $i$  and  $j$  are number of matrices).
- The optimum can be described by the following recursive formulation.
  - If  $i = j$ , there is only one matrix and thus  $m[i, i] = 0$  (the diagonal entries).
  - If  $i < j$ , we are asking for the product  $A_{i...j}$ .
  - This can be split by considering each  $k$ ,  $i \leq k < j$ , as  $A_{i...k}$  times  $A_{k+1...j}$ .

- The optimum time to compute  $A_{i\dots k}$  is  $m[i, k]$  and optimum time for  $A_{k+1\dots j}$  is in  $m[k + 1, j]$ .
- Since  $A_{i\dots k}$  is a  $p_{i-1} \times p_k$  matrix and  $A_{k+1\dots j}$  is  $p_k \times p_j$  matrix, the time to multiply them is  $p_{i-1} \times p_k \times p_j$ .
- This suggests the following recursive rule:

$$m[i, i] = 0$$

$$m[i, j] = \min_{i \leq k < j} (m[i, k] + m[k + 1, j] + p_{i-1}p_kp_j)$$

- We do not want to calculate  $m$  entries recursively.
- So how should we proceed? We will fill  $m$  along the diagonals.
- Set all  $m[i, i] = 0$  using the base condition.
- Compute cost for multiplication of a sequence of 2 matrices.

- These are  $m[1,2], m[2,3], m[3,4], \dots, m[n-1, n]$ .
- For example,  $m[1,2]$  is:

$$m[1,2] = m[1,1] + m[2,2] + p_0 \cdot p_1 \cdot p_2$$

- For example, the  $m$  for product of 5 matrices at this stage would be:

$m[1, 1]$	$\leftarrow m[1, 2]$ $\downarrow$			
	$m[2, 2]$	$\leftarrow m[2, 3]$ $\downarrow$		
		$m[3, 3]$	$\leftarrow m[3, 4]$ $\downarrow$	
			$m[4, 4]$	$\leftarrow m[4, 5]$ $\downarrow$
				$m[5, 5]$

- Next, we compute cost of multiplication for sequences of three matrices.
- These are  $m[1, 3], m[2, 4], m[3, 5], \dots, m[n - 2, n]$ .
- For example,  $m[1, 3]$  is:

$$m[1, 3] = \min \begin{cases} m[1, 1] + m[2, 3] + p_0 \cdot p_1 \cdot p_3 \\ m[1, 2] + m[3, 3] + p_0 \cdot p_2 \cdot p_3 \end{cases}$$

- We repeat the process for sequences of four, five and higher number of matrices.
- The final result will end up in  $m[1, n]$ .

- **Example:** We want to find the optimal multiplication order for:

$$\begin{matrix} A_1 & \cdot & A_2 & \cdot & A_3 & \cdot & A_4 & \cdot & A_5 \\ (5 \times 4) & & (4 \times 6) & & (6 \times 2) & & (2 \times 7) & & (7 \times 3) \end{matrix}$$

- We will compute the entries of the  $m$  matrix starting with the base condition.
- We first fill that main diagonal using base case:



0				
	0			
		0		
			0	
				0

- Next, we compute the entries in the first super diagonal, i.e., the diagonal above the main diagonal:

$$m[1,2] = m[1,1] + m[2,2] + p_0 \cdot p_1 \cdot p_2 = 0 + 0 + (5 \cdot 4 \cdot 6) = 120$$

$$m[2,3] = m[2,2] + m[3,3] + p_1 \cdot p_2 \cdot p_3 = 0 + 0 + (4 \cdot 6 \cdot 2) = 48$$

$$m[3,4] = m[3,3] + m[4,4] + p_2 \cdot p_3 \cdot p_4 = 0 + 0 + (6 \cdot 2 \cdot 7) = 84$$

$$m[4,5] = m[4,4] + m[5,5] + p_3 \cdot p_4 \cdot p_5 = 0 + 0 + (2 \cdot 7 \cdot 3) = 42$$

- The matrix  $m$  now looks as follows after the product of two matrices (first super diagonal):

0	120			
	0	48		
		0	84	
			0	42
				0

- We now proceed to the second super diagonal.
- This time, however, we will need to try two possible values for  $k$ .
- For example, there are two possible splits for computing  $m[1, 3]$ .

- We will choose the split that yields the minimum:

$$m[1,3] = m[1,1] + m[2,3] + p_0 \cdot p_1 \cdot p_3 = 0 + 48 + (5 \cdot 4 \cdot 2) = 88$$

$$m[1,3] = m[1,2] + m[3,3] + p_0 \cdot p_2 \cdot p_3 = 120 + 0 + (5 \cdot 6 \cdot 2) = 180$$

*the minimum  $m[1,3] = 88$  occurs with  $k = 1$*

- Similarly, for  $m[2, 4]$  and  $m[3, 5]$ :

$$m[2,4] = m[2,2] + m[3,4] + p_1 \cdot p_2 \cdot p_4 = 0 + 48 + (4 \cdot 6 \cdot 7) = 252$$

$$m[2,4] = m[2,3] + m[4,4] + p_1 \cdot p_3 \cdot p_4 = 48 + 0 + (4 \cdot 2 \cdot 7) = 104$$

*the minimum  $m[2,4] = 104$  at  $k = 3$*

$$m[3,5] = m[3,3] + m[4,5] + p_2 \cdot p_3 \cdot p_5 = 0 + 42 + (6 \cdot 2 \cdot 3) = 78$$

$$m[3,5] = m[3,4] + m[5,5] + p_2 \cdot p_4 \cdot p_5 = 84 + 0 + (6 \cdot 7 \cdot 3) = 210$$

*the minimum  $m[3,5] = 78$  at  $k = 3$*

- With the second super diagonal computed, the  $m$  matrix looks as follow:

0	120	88		
	0	48	104	
		0	84	78
			0	42
				0

- We repeat the process for the remaining diagonals.
- However, the number of possible splits (values of  $k$ ) increases:

$$m[1,4] = m[1,1] + m[2,4] + p_0 \cdot p_1 \cdot p_4 = 0 + 104 + (5 \cdot 4 \cdot 7) = 244$$

$$m[1,4] = m[1,2] + m[3,4] + p_0 \cdot p_2 \cdot p_4 = 120 + 84 + (5 \cdot 6 \cdot 7) = 414$$

$$m[1,4] = m[1,3] + m[4,4] + p_0 \cdot p_3 \cdot p_4 = 88 + 0 + (5 \cdot 2 \cdot 7) = 158$$

*the minimum  $m[1,4] = 158$  at  $k = 3$*



$$m[2,5] = m[2,2] + m[3,5] + p_1 \cdot p_2 \cdot p_5 = 0 + 78 + (4 \cdot 6 \cdot 3) = 150$$

$$m[2,5] = m[2,3] + m[4,5] + p_1 \cdot p_3 \cdot p_5 = 48 + 42 + (4 \cdot 2 \cdot 3) = 114$$

$$m[2,5] = m[2,4] + m[5,5] + p_1 \cdot p_4 \cdot p_5 = 104 + 0 + (4 \cdot 7 \cdot 3) = 188$$

*the minimum  $m[2,5] = 114$  at  $k = 3$*

- The matrix  $m$  at this stage is:

0	120	88	158	
	0	48	104	114
		0	84	78
			0	42
				0

That leaves the  $m[1, 5]$  which can now be computed:

$$m[1,5] = m[1,1] + m[2,5] + p_0 \cdot p_1 \cdot p_5 = 0 + 114 + (5 \cdot 4 \cdot 3) = 174$$

$$m[1,5] = m[1,2] + m[3,5] + p_0 \cdot p_2 \cdot p_5 = 120 + 78 + (5 \cdot 6 \cdot 3) = 288$$

$$m[1,5] = m[1,3] + m[4,5] + p_0 \cdot p_3 \cdot p_5 = 88 + 42 + (5 \cdot 2 \cdot 3) = 160$$

$$m[1,5] = m[1,4] + m[5,5] + p_0 \cdot p_4 \cdot p_5 = 158 + 0 + (5 \cdot 7 \cdot 3) = 263$$

*the minimum  $m[1,5] = 160$  at  $k = 3$*

- We thus have the final cost matrix  $m$  as:

0	120	88	158	<i>160</i>
0	0	48	104	114
0	0	0	84	78
0	0	0	0	42
0	0	0	0	0

- Here is the order in which  $m$  entries are calculated:

0	1	5	8	10
0	0	2	6	9
0	0	0	3	7
0	0	0	0	4
0	0	0	0	0


- And the split  $k$  values that led to a minimum  $m[i, j]$  value:

0	1	1	3	3
	0	2	3	3
		0	3	3
			0	4
				0

- Based on the computation, the minimum cost for multiplying the five matrices is 160 and the optimal.
- So, the optimal order for multiplication is:

$$((A_1(A_2A_3))(A_4A_5))$$

- This can be represented as a binary tree:



Optimum matrix multiplication order for the five matrices example