

Antenatal fetal wellbeing

Hamdy Bakry, MD

2026-07-16

1 According to Modality

- Ultrasound
- Doppler studies
- Non-stress test (NST)
- Biophysical profile (BPP)
- Maternal serum screening
- Invasive testing (CVS, amniocentesis)
- Cell-free fetal DNA testing

2 According to GA

2.1 First Trimester (up to 13⁺⁶ weeks)

- Ultrasound: Nuchal translucency, crown-rump length, early anatomy
- Biochemical markers: PAPP-A, free beta-hCG
- Cell-free fetal DNA testing
- Chorionic villus sampling (CVS) if indicated
- Fetal heartbeat detection
- Dating scan
- Early fetal anomaly screening
- Assessment of multiple pregnancies
- Early placental location assessment

2.2 Second Trimester (14 to 27⁺⁶ weeks)

- Ultrasound: Detailed anatomy scan, growth assessment
- Maternal serum screening: Quadruple test
- Amniocentesis if indicated
- Fetal echocardiography if indicated
- Assessment of fetal movements
- Placental location and function assessment
- Cervical length measurement if indicated

2.3 Third Trimester (28 Weeks to delivery)

- Ultrasound: Growth scans, amniotic fluid assessment, Doppler studies

Antenatal fetal wellbeing

- Non-stress test (NST)
- Biophysical profile (BPP)
- Assessment of fetal position and presentation
- Cervical assessment for labor readiness
- Fetal movement counting
- Assessment of placental function and aging
- Monitoring for signs of fetal distress
- Preparation for delivery planning
- Assessment of multiple pregnancies
- Evaluation of fetal well-being in high-risk pregnancies
- Assessment of fetal growth restriction (FGR) or macrosomia
- Screening for gestational diabetes effects on fetus
- Monitoring for preeclampsia effects on fetus
- Evaluation of amniotic fluid volume (polyhydramnios/oligohydramnios)
- Fetal lung maturity assessment if preterm delivery is anticipated