

RELATIVE HITCHIN FIBRATIONS AND ENDOSCOPY FOR UNITARY FRIEDBERG–JACQUET VARIETIES

THOMAS HAMEISTER AND SPENCER LESLIE

ABSTRACT. We prove a relative fundamental lemma for the unitary symmetric variety $\mathrm{GL}_n(k') \backslash \mathrm{U}_{2n}$, proving a conjecture of [Les24, LXZ25a]. Our proof reduces to the Lie algebra and argues over function fields in positive characteristic, using methods from [LN08, Ngo10]. In particular, we reinterpret orbital integrals in terms of the cohomology of relative affine Springer fibers and globalize this using unitary versions of Hitchin moduli spaces for symmetric pairs via a product formula. As the main global input, we prove a matching of perverse sheaves between relative Hitchin moduli spaces and their corresponding endoscopic variants.

CONTENTS

1.	Introduction	1
2.	Invariant theory for symmetric varieties	6
3.	Unitary Higgs Bundles	15
4.	The relative Hitchin moduli space	17
5.	Symmetries of relative Hitchin fibers	23
6.	Relative affine Springer fibers	33
7.	Point Counting	50
8.	The support theorem and computation of cohomology	59
9.	Proof of the Main Theorems	62
	References	66

1. INTRODUCTION

Fix an unramified quadratic extension of nonarchimedean local fields F'/F of positive characteristic. In the body of this paper, we prove an endoscopic fundamental lemma for the Lie algebra of the symmetric space $\mathrm{GL}_{n,F'} \backslash \mathrm{U}_{2n}$ (Theorem 1.1 below). In the appendix, we show that our statement implies the fundamental lemma for $\mathrm{GL}_{n,F'} \backslash \mathrm{U}_{2n}$ at the group level with no assumption on characteristic. This result confirms a conjecture in [LXZ25a] and fits naturally into a program towards relative endoscopy initiated by the second author [Les24]. Its resolution is an essential step towards the stabilization of the relative trace formula associated to $\mathrm{GL}_{n,F'}$ -periods of automorphic forms on U_{2n} .

This is the first example of an endoscopic relative fundamental lemma to be proved utilizing geometric techniques pioneered by Ngô and Ngô–Laumon [LN08, Ngo06, Ngo10]. The moduli spaces arising naturally in the proof are expected to be important objects of independent interest in the relative Langlands program of Ben-Zvi, Sakellaridis, and Venkatesh [BZSV23], see for example [HLM24]. We also expect the methods used in this paper can be used to re-prove the fundamental lemma associated to the Lie algebra of $\mathrm{U}_n \times \mathrm{U}_n \backslash \mathrm{U}_{2n}$, which was proved in [Les25] using methods from p -adic harmonic analysis.

1.1. Background and motivation. The study of relative fundamental lemmas arises in part from the stabilization of the relative trace formula, an important tool used to study questions of distinction for automorphic representations. Namely, let $H \subset G$ be reductive, quasi-split groups over a number field F with $X = H \backslash G$ spherical, and let $\mathbb{A}_F$ denote the ring of adeles of F . A cuspidal automorphic representation π of G is H -*distinguished* if the period integral

$$\int_{[H]} \varphi(h) dh$$

is nonzero for some $\varphi \in \pi$, where $[H] = H(F) \backslash H(\mathbb{A}_F)$. A general philosophy arising from the relative Langlands program [SV17, BZSV23] is that properties of period integrals can be understood through L functions determined by X . For example, for linear periods of $\mathrm{GL}_n \times \mathrm{GL}_n \subset \mathrm{GL}_{2n}$, Friedberg and Jacquet show that H distinction of π is given by a condition that π be of “symplectic type” and that $L(\frac{1}{2}, \pi, \mathrm{std}) \neq 0$ [FJ93]. This paper concerns a unitary version of Friedberg and Jacquet’s linear periods given by $\mathrm{GL}_{n,F'} \subset \mathrm{U}_{2n}$. This was one of several cases considered in recent work of Leslie, Xiao, and Zhang, which in particular proposes conjectures detecting distinction in this case in terms of base change L functions [LXZ25a, LXZ25b].

To establish distinction properties, an essential tool is the relative trace formula, introduced by Jacquet to study a formula of Waldspurger [JL85]. The relative trace formula relates a spectral side, written in terms of period integrals, with a geometric side, in terms of orbital integrals. To be useful towards properties of distinction, it is important that the geometric side of the trace formula be written in terms of geometric data. In practice, this involves grouping orbital integrals by stable (geometric) conjugacy, not just rational conjugacy. This process is known as *stabilization*. The relative fundamental lemma relates these groupings of orbital integrals by stable conjugacy to stable orbital integrals for endoscopic spaces. This process should be thought of as an instance of functoriality in the relative Langlands program.

It is a stunning fact that all known proofs of the classical fundamental lemma come from a reduction to positive characteristic, where powerful geometric techniques in either perverse sheaves or p -adic integration can be applied [Ngo10, GWZ20]. Subsequent work has established related formulas for the Jacquet-Rallis fundamental lemma [Yun16, WZ24] and the classical fundamental lemma without reduction to Lie algebras [Wan26] using generalizations of affine Springer fibers and moduli spaces of Higgs bundles. This paper uses novel variations of such methods to study relative endoscopy, representing the first such step for a case with nontrivial endoscopic terms.

1.2. Main Theorem. Let F be a non-archimedean local field with ring of integers $\mathcal{O}$ and with residue field $k = \mathbb{F}_q$ a finite field of characteristic greater than n . Let F' be the fixed nonsplit, unramified extension of F with residue field k' . Let $\sigma \in \mathrm{Gal}(k'/k)$ be the order 2 generator, and denote $x^* = {}^t\sigma(x)$. We denote by

$$\mathrm{U}_n = \{g \in \mathrm{GL}_n(F') \mid g \cdot g^* = 1\}$$

the rank n unitary group over F and by

$$\mathrm{Herm}_n = \{x \in \mathfrak{gl}_n \mid x = x^*\}$$

the space of $n \times n$ Hermitian matrices. Consider the following two representations:

- The “Siegel” case: $H^{\mathbf{S}} = \mathrm{Res}_{F'/F} \mathrm{GL}_n$ acting on $V^{\mathbf{S}} = \mathrm{Herm}_n \times \mathrm{Herm}_n$ by

$$h \cdot (x, y) = (h \cdot x \cdot h^*, (h^*)^{-1} x h^{-1})$$

- The “unitary” case: $H^{\mathbf{U}} = \mathrm{U}_n \times \mathrm{U}_n$ acting on $V^{\mathbf{U}} = \mathfrak{gl}_n(F')$ by

$$(h_1, h_2) \cdot x = h_1 x h_2^*$$

We have an identification $V^{\mathbf{S}}//H^{\mathbf{S}} \simeq V^{\mathbf{U}}//H^{\mathbf{U}}$ of GIT quotients, and we denote this common quotient by $\mathfrak{c}$. Let $\mathfrak{c}^{\text{rs}}$ denote the image of regular, semisimple matrices. For any $a \in \mathfrak{c}(\mathcal{O}) \cap \mathfrak{c}^{\text{rs}}(F)$ we have a Kostant–Rallis section $[\epsilon](a) \in V^{\mathbf{S}}(F)$, and we consider the orbital integral

$$\text{Orb}_a^{\mathbf{S}, \kappa} = \sum_{\gamma \sim [\epsilon](a)} \langle \kappa, \text{inv}(\gamma, \gamma_0) \rangle \int_{I_\gamma \backslash H^{\mathbf{S}}(F)} \mathbb{1}_{V^{\mathbf{S}}(\mathcal{O})}(h^{-1} \cdot \gamma) \frac{dh}{dt}$$

where the sum is over all F conjugacy classes of γ which are conjugate to $[\epsilon](a)$ over $\check{F} = F \otimes_k \bar{k}$; I_γ is the centralizer of γ in $H^{\mathbf{S}}(F)$; and dh and dt are Haar measures on $H(F)$ and I_γ , respectively, normalized so that $\text{vol}(H^{\mathbf{S}}(\mathcal{O}), dh) = 1$. Here the weights $\langle \kappa, \text{inv}(\gamma, \gamma_0) \rangle$ depend on a parameter κ , which is a 2-torsion element of the canonical torus of the dual symmetric variety $\hat{\mathbf{X}} = \hat{\mathbf{G}}_{\mathbf{X}} \backslash \hat{\mathbf{G}}$. We may similarly define the stable orbital integral $\text{SO}_a^{\mathbf{U}}$ by summing over all lifts $\gamma \in V^{\mathbf{U}}(F)$ of a (without κ weights). For a product $X^{\mathfrak{c}} = X_r^{\mathbf{S}} \times X_s^{\mathbf{U}}$, with subscripts indicating rank, the stable orbital integral is the product $\text{SO}_{(a^{\mathbf{S}}, a^{\mathbf{U}})}^{\mathfrak{c}} = \text{SO}_{a^{\mathbf{S}}}^{\mathbf{S}} \times \text{SO}_{a^{\mathbf{U}}}^{\mathbf{U}}$.

Theorem 1.1. *For every elliptic endoscopic datum $\mathfrak{c} = (\kappa, \rho)$ over F (see §2.3), there is an endoscopic symmetric variety $X^{\mathfrak{c}}$ defined over F , which is a product of Siegel and unitary symmetric varieties of complementary rank, and a transfer map*

$$\nu_{\mathfrak{c}}: \mathfrak{c}_{\mathfrak{c}} \rightarrow \mathfrak{c}$$

such that for any $a_{\mathfrak{c}} \in \mathfrak{c}_{\mathfrak{c}}(\mathcal{O})$ with image $a = \nu_{\mathfrak{c}}(a_{\mathfrak{c}}) \in \mathfrak{c}(\mathcal{O}) \cap \mathfrak{c}(F)^{\text{rs}}$, we have

$$\text{Orb}_a^{\mathbf{S}, \kappa} = q^{r_{\mathfrak{c}}(a)} \text{SO}_{a_{\mathfrak{c}}}^{\mathfrak{c}}$$

where $r_{\mathfrak{c}}(a)$ is a nonnegative integer, defined in §7.4.

The construction of the spaces $X^{\mathfrak{c}}$ and transfer maps $\nu_{\mathfrak{c}}$ in the above theorem was done in [Les24] for a much broader class of symmetric varieties.

1.3. Reformulation as Lattice Counts. We reformulate Theorem 1.1 as a point count of a moduli space of lattices which we call the *relative affine Springer fiber*. In particular, to an n -tuple $a = (a_2, \dots, a_{2n})$ in $\mathcal{O}^n$, we may form the F' algebra

$$R_a(F') = F'[x]/(P_a(x)), \quad \text{for } P_a(x) := x^{2n} + a_2 x^{2n-2} + \dots + a_{2n-2} x^2 + a_{2n}$$

with the obvious sub- $\mathcal{O}'$ algebra $R_a(\mathcal{O}')$. Then $R_a(F')$ carries natural involutions $\iota(x) = -x$ and σ from the Galois action of F'/F .

We say $\Lambda \subset R_a(F')$ is an $R_a(\mathcal{O}')$ lattice if it is stable under the $R_a(\mathcal{O}')$ action and it is free of rank $2n$ as an $\mathcal{O}'$ module. There are natural non-degenerate Hermitian forms $h^{\mathbf{S}}$ and $h^{\mathbf{U}}$ on $R_a(F')$ constructed via the trace pairing by

$$h^{\mathbf{S}}(f, g) = \text{Tr}_{R_a(F')/F'} \left(\frac{f \cdot \sigma(g)}{P'_a} \right), \quad h^{\mathbf{U}}(f, g) = h^{\mathbf{S}}(xf, g).$$

For $\star \in \{\mathbf{S}, \mathbf{U}\}$, we define $\mathcal{M}^{\star}(a)$ as the moduli of $R_a(\mathcal{O}')$ lattices Λ such that $\iota(\Lambda) = \Lambda$ and Λ is self dual with respect to $h^{\star}$. We call $\mathcal{M}^{\star}(a)$ the relative affine Springer fiber (RASF) at a . It is a direct analogue of affine Springer fibers, subvarieties of the affine Grassmannian introduced by Kazhdan and Lusztig [KL88] of profound importance to geometric representation theory. In fact, it can be considered as the fixed points of the involution ι acting on the unitary affine Springer fiber studied in [LN08]. We remark on generalizations to other spherical varieties in §6.2.

The RASF $\mathcal{M}^{\star}(a)$ is representable by a locally finite type, projective scheme. It admits an action of a group of symmetries $\mathcal{P}(J_a)$, which is independent of $\star \in \{\mathbf{S}, \mathbf{U}\}$. Together, $\mathcal{M}^{\star}(a)$ and $\mathcal{P}(J_a)$ satisfy the point counting properties of [Ngo10, Hypotheses 8.1.9]; namely, the stabilizers of $\mathcal{P}(J_a)$ acting on $\mathcal{M}^{\star}(a)$ are of finite type, and there is a canonical lattice $\Gamma_a \subset \mathcal{P}(J_a)$ which acts freely on $\mathcal{M}^{\star}(a)$ such that $\mathcal{M}^{\star}(a)/\Gamma_a$ and $\mathcal{P}(J_a)/\Gamma_a$ are finite type. In contrast to the case of affine Springer fibers, however, $\mathcal{M}^{\star}(a)/\Gamma_a$ need not be connected, even for an optimally chosen lattice Γ_a . Instead,

Γ_a only acts transitively on irreducible components in fixed connected components of $\mathcal{M}^*(a)$. This phenomena obstructs standard approaches to proving dimension formulas for RASF's, and so our dimension formula (Proposition 6.28) and the equidimensionality of $\mathcal{M}^*(a)$ (Proposition 6.30) can be viewed as one of our main results. Indeed, inspired by work of Jingren Chi [Chi22], we use global methods to approach these questions, exploiting properness of the relative Hitchin map over a carefully constructed family.

For an endoscopic symmetric space $X^\epsilon = X_r^{\mathbf{S}} \times X_s^{\mathbf{U}}$, the RASF is a product $\mathcal{M}^\epsilon(a^{\mathbf{S}}, a^{\mathbf{U}}) = \mathcal{M}^{\mathbf{S}}(a^{\mathbf{S}}) \times \mathcal{M}^{\mathbf{U}}(a^{\mathbf{U}})$. We consider the endoscopic case with the action of $\mathcal{P}(J_a)$, whose lattice Γ_a still acts simply transitively, but whose stabilizers are usually larger than $\mathcal{P}(J_{\epsilon,a})$. The finiteness hypotheses on $\mathcal{M}^*(a)$ and $\mathcal{P}(J_a)$ imply in particular that there are finitely many $\mathcal{P}(J_a)$ orbits on $\mathcal{M}^*(a)$. Therefore, the sums

$$\#[\mathcal{M}^{\mathbf{S}}(a)/\mathcal{P}(J_a)](k)_\kappa = \sum_O \frac{\langle \kappa, \text{cl}(O) \rangle}{\# \text{Aut}_{\mathcal{P}(J_a)}(O)(k)}, \quad \#[\mathcal{M}^\epsilon(a)/\mathcal{P}(J_a)](k) = \sum_O \frac{1}{\# \text{Aut}_{\mathcal{P}(J_a)}(O)(k)},$$

which sum over points O in the groupoid $[\mathcal{M}^{\mathbf{S}}(a)/\mathcal{P}(J_a)](k)$, respectively $[\mathcal{M}^\epsilon(a)/\mathcal{P}(J_a)](k)$, are well-defined where $\langle \kappa, \text{cl}(O) \rangle$ is a sign defined in §7.2. The orbital integrals appearing in Theorem 1.1 can be written in terms of these weighted point counts of groupoids.

Theorem 1.2. *Let ϵ be an endoscopic datum, and suppose that $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$ and $a_\epsilon \in \mathfrak{c}_\epsilon^\heartsuit(\mathcal{O}_v)$ are such that $\nu_\epsilon(a_\epsilon) = a$. Then, for all finite extensions K/k_v , we have*

$$(1.1) \quad \#[\mathcal{M}_v^{\mathbf{S}}(a)/\mathcal{P}_v(J_a)](K)_\kappa = q^{r_\epsilon(a) \deg(v)} \cdot \#[\mathcal{M}_v^\epsilon(a_\epsilon)/\mathcal{P}_v(J_a)](K)$$

Note that Theorem 1.2 combined with Grothendieck-Lefschetz reduces Theorem 1.1 to a computation of cohomology of RASFs. This was the perspective taken by Goresky, Kottwitz, and MacPherson [GKM04, GKM06, GKM09], who attempted to prove the fundamental lemma by proving purity of affine Springer fibers.

1.4. Global moduli spaces. To study point counts on RASFs, we construct global moduli spaces called *relative Hitchin spaces*, which are related to RASFs via a product formula (Proposition 6.24). These global moduli spaces are unitary analogues of Hitchin systems for symmetric varieties, which have been studied extensively by Schaposnik [Sch13, Sch18b, Sch15, Sch18a], Peón-Nieto and García-Prada [GPPN23], and the first author with Morrissey [HM25].

Fix a smooth, projective, geometrically-connected curve Σ of genus at least 2 and a divisor D of degree at least $2g$. Fix an étale double cover $\vartheta: \Sigma' \rightarrow \Sigma$, with Σ' smooth, projective, and geometrically connected as well. From this, we may form moduli spaces $\mathcal{M}^* = \text{Maps}(\Sigma, [V_D^*/H^*])$, where the subscript D denotes the induced bundle on Σ , and H^* and V^* are taken over k instead of F .

The moduli spaces $\mathcal{M}^*$ have a simple and more explicit description in terms of fixed point stacks of involutions on moduli of unitary Higgs bundles. Namely, we let $\mathcal{N}$ describe the moduli of unitary Higgs bundles of Laumon and Ngô, whose S points parametrize

$$\mathcal{N}(S) = \left\{ (\mathcal{E}', h, \phi') \left| \begin{array}{l} \mathcal{E}' \text{ is a rank } 2n \text{ vector bundle on } \Sigma' \times S, \\ h: \mathcal{E}' \rightarrow \sigma^*(\mathcal{E}')^\vee \text{ is a Hermitian form on } \mathcal{E}', \text{ i.e. } h = \sigma^* h^\vee, \\ \phi': \mathcal{E}' \rightarrow \mathcal{E}' \otimes D, \text{ satisfying } \sigma^*(\phi')^\vee \circ h + h \circ \phi' = 0 \end{array} \right. \right\}.$$

Then, we have involutions ι^* on $\mathcal{N}$ by

$$\iota^*(\mathcal{E}', h, \phi') = \begin{cases} (\mathcal{E}', -h, -\phi') & \text{if } \star = \mathbf{S} \\ (\mathcal{E}', h, -\phi') & \text{if } \star = \mathbf{U} \end{cases}$$

Then, $\mathcal{M}^\star$ is a distinguished component of the fixed point stack $\mathcal{N}^{\iota^\star}$, and is equal when we restrict to a suitable anisotropic locus. We have concrete realizations of $\mathcal{M}^\mathbf{S}$ and $\mathcal{M}^\mathbf{U}$; for example,

$$\mathcal{M}^\mathbf{S}(S) = \left\{ (\mathcal{E}', h, \phi'_{12}, \phi'_{21}) \left| \begin{array}{l} \mathcal{E}' \text{ is a rank } n \text{ vector bundle on } \Sigma' \times S, \phi'_{12}: \mathcal{E}' \rightarrow \sigma^*(\mathcal{E}')^\vee \otimes D \\ \text{and } \phi'_{21}: \sigma^*(\mathcal{E}')^\vee \rightarrow \mathcal{E}' \otimes D \text{ such that } \sigma^* \phi'_{ij} = \phi'_{ij} \\ \text{and } h: \mathcal{E}' \rightarrow \sigma^*(\mathcal{E}')^\vee \text{ is a Hermitian form on } \mathcal{E}' \end{array} \right. \right\}.$$

For $\star \in \{\mathbf{S}, \mathbf{U}\}$, there is a Hitchin-type map

$$f^\star: \mathcal{M}^\star \rightarrow \mathcal{A}$$

where $\mathcal{A} = \bigoplus_{i=1}^n H^0(\Sigma, D^{\otimes 2i})$ is an affine space. These maps essentially compute the characteristic polynomial of the Higgs field ϕ' . As in the case of the Hitchin moduli space, this morphism reveals a remarkable amount of structure in $\mathcal{M}^\star$. For example, there is a family of curves $\tilde{\Sigma}_a$ over $\mathcal{A}$ such that whenever $\tilde{\Sigma}_a$ is reduced, the fiber $f^{\star,-1}(a)$ admits a spectral description as a sort of compactified Prym variety on $\tilde{\Sigma}_a$ (Proposition 3.4). In particular, there is a commutative group stack $\mathcal{P}$, itself a relative Prym variety of a family of curves over $\mathcal{A}$, which acts on the fibers of $f^\star$ with finitely many orbits whenever $\tilde{\Sigma}_a$ is reduced. When $\tilde{\Sigma}_a$ is smooth (a generic condition on $\mathcal{A}$), the fiber $f^{\star,-1}(a)$ is a finite unions of $\mathcal{P}_a$ torsors.

There is a Beauville-Laszlo style gluing map from RASFs to relative Higgs bundles. This produces a product formula

$$\#[\mathcal{M}_a^\star / \mathcal{P}_a](k)_\kappa = \prod_{v \in |\Sigma|} \#[\mathcal{M}_v^\star(a) / \mathcal{P}(J_{a,v})](k_v)_\kappa$$

for a in an elliptic locus $\mathcal{A}^{\text{ani}} \subset \mathcal{A}$, where only finitely many terms on the right-hand side are nontrivial (Proposition 6.24).

After restricting to an anisotropic locus (§5.1) and performing an étale base change on the base $\underline{\mathcal{A}}^{\text{ani}} \rightarrow \mathcal{A}^{\text{ani}}$ (§5.3), there is a surjection $\mathbb{X}_*(\hat{\mathbf{C}}) \rightarrow \pi_0(\mathcal{P}_a)$ which allows us to decompose the cohomology into $\kappa \in \hat{\mathbf{C}}[2]$ constituent pieces

$$\mathbf{R}f_{\underline{*}}^{\text{ani}} \overline{\mathbb{Q}}_\ell = \bigoplus_{\kappa} (\mathbf{R}f_{\underline{*}}^{\text{ani}} \overline{\mathbb{Q}}_\ell)_\kappa$$

where $(\mathbf{R}f_{\underline{*}}^{\text{ani}} \overline{\mathbb{Q}}_\ell)_\kappa \subset \mathbf{R}f_{\underline{*}}^{\text{ani}} \overline{\mathbb{Q}}_\ell$ is the subsheaf on which $\mathbb{X}_*(\hat{\mathbf{C}})$ acts through evaluation at κ . We prove the following global result.

Theorem 1.3. *The transfer maps induce closed embeddings on Hitchin bases*

$$\nu_\epsilon: \underline{\mathcal{A}}_\epsilon \rightarrow \underline{\mathcal{A}}.$$

For any $\kappa \in \hat{\mathbf{C}}[2]$, we have an isomorphism

$$\nu_\epsilon^* (\mathbf{R}f_{\underline{*}}^{\mathbf{S}, \text{ani}} \overline{\mathbb{Q}}_\ell)_\kappa [2r_\epsilon(D)](r_\epsilon(D)) = (\mathbf{R}f_{\underline{*}}^{\epsilon, \text{ani}} \overline{\mathbb{Q}}_\ell)_{\text{st}}$$

over $\bar{k}$ restricted to the locus $\underline{\mathcal{A}}_\epsilon^{\text{ani}}$ which induces an isomorphism of semisimplifications over k .

We note that we do not prove Theorem 1.3 directly, but rather first prove it over a subset of $\mathcal{A}_\epsilon^{\text{ani}}$, then use this to deduce Theorem 1.2 (and hence Theorem 1.1) before reversing the flow of information to deduce the full strength of Theorem 1.3.

We prove Theorem 1.3 roughly as follows: For each κ , the decomposition theorem of Beilinson, Bernstein, Deligne, and Gabber [?] implies that $(\mathbf{R}f_{\underline{*}}^{\text{ani}} \overline{\mathbb{Q}}_\ell)_\kappa$ is a semisimple perverse sheaf over $\bar{k}$. By Ngô's support theorem, to determine the supports of the simple summands of $(\mathbf{R}f_{\underline{*}}^{\text{ani}} \overline{\mathbb{Q}}_\ell)_\kappa$, it is enough to study the top degree ordinary cohomology $\mathbf{R}^{2d} f_{\underline{*}} \overline{\mathbb{Q}}_\ell$ along δ critical subsets. For this, it will be enough to study *regular* Higgs bundles. Using the theory of the regular quotient, developed for symmetric varieties by the first author and B. Morrissey in [HM25], we show there is an open

set $\mathcal{A}^\dagger \subset \mathcal{A}^{\text{ani}}$ where the regular part $\mathcal{M}^{\star, \text{reg}} \subset \mathcal{M}^\star$ is dense, and the cohomology $R^{2d} \underline{f}_*^{\text{ani}} \overline{\mathbb{Q}}_\ell$ can be explicitly computed using an étale cover $\mathcal{A}^{\text{reg}} \rightarrow \mathcal{A}$ and a factorization of the relative Hitchin map

$$\mathcal{M}^{\star, \text{reg}} \rightarrow \mathcal{A}^{\text{reg}} \rightarrow \mathcal{A}.$$

After identifying the top cohomology on $\mathcal{A}^\dagger$, proving Theorem 1.3 reduces to checking some small cases of Theorem 1.1 (see §7.4).

1.5. Structure of the article. In §2, we discuss invariant theory that underlies both the local and global formulations of our main results. Most results in this section are not new, the exception being the k rational results in §2.2. In §3 we review some basic results of Ngô and Laumon on unitary Higgs bundles, which we will use to deduce some basic properties of our global moduli spaces. In §4, we introduce and study the relative Hitchin fibration. In §6, we introduce and study relative affine Springer fibers. The crucial technical result in this section is the dimension formula and equidimensionality (Propositions 6.28 and 6.30), which uses global methods. In §7, we recall the formalism of point counting for groupoids and show that our orbital integrals can be interpreted as point counts on RASFs. We also deduce a few cases of Theorem 1.1, including the case where both sides of the equality are zero. In §8, we outline how to apply Ngô's support theorem to our case, including the computation of top degree cohomology over $\mathcal{A}^\dagger$. Finally, in §9 we finish the proofs of the main theorems. The key new technical piece is the interpolation step: For any local a , we must extend a to a point in the Hitchin base with nonzero point counts at every place of Σ .

1.6. Acknowledgments. The first author thanks Ngô Bao Châu for his role in introducing him to related moduli spaces for his PhD thesis. He also thanks Siddharth Mahendrakar, Xinchun Ma, Zeyu Wang and Griffin Wang for useful discussions.

Spencer's acknowledgements

Prior to the editing process, AI was not used in any significant way in this work, neither in obtaining results nor writing. However, ChatGPT 5.6 Astra was prompted to identify errors in a draft, which led to corrections in the proof of Proposition 7.11 and minor modifications to the interpolation argument in section 9.2 along with numerous typos throughout the paper.

2. INVARIANT THEORY FOR SYMMETRIC VARIETIES

2.1. The representation $\mathbf{H}$ acting on $\mathfrak{h}^\perp$. For this section (and this section only), we assume throughout that the base field k is algebraically closed such that $\text{char}(k) \neq 2$. The results of this section are discussed in greater detail in [HM25], with most of the technical steps in the proofs in characteristic p contained in Levy's thesis [Lev07]. While we will keep this section focused on our example of interest, all constructions can be made for general symmetric varieties with minimal modifications (cf. [HM25]), and much remains true for general spherical varieties (see [HLM24] for a summary).

Let $\mathbf{G} = \text{GL}_{2n}$. While results in this section do not depend on it, we will assume that the characteristic of k is greater than n for Corollary 5.30. We have an algebraic involution $\theta: \mathbf{G} \rightarrow \mathbf{G}$ given by

$$\theta(g) = I_{n,n} g I_{n,n}, \quad \text{for } I_{n,n} = \begin{pmatrix} I_n & \\ & -I_n \end{pmatrix}.$$

The fixed points under this action are given by the Levi subgroup $\mathbf{H} = \text{GL}_n \times \text{GL}_n$ inside $\mathbf{G}$. Moreover, the action of θ induces a decomposition on the level of Lie algebras

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{h}^\perp$$

where $\mathfrak{h} = \text{Lie}(\mathbf{H})$ is the fixed points of the action of θ and $\mathfrak{h}^\perp$, the orthogonal complement of $\mathfrak{h}$ under a nondegenerate G invariant form on $\mathfrak{g}$, is the (-1) -eigenspace of the θ action. The group $\mathbf{H}$ acts on $\mathfrak{h}^\perp$ by restricting the adjoint action.

We also define the following analogue of Cartan subalgebras adapted for the symmetric setting.

Definition 2.1. A θ -Cartan subalgebra $\mathfrak{a}$ is a maximal (with respect to inclusion) abelian Lie algebra consisting of semi-simple elements contained in $\mathfrak{h}^\perp$.

All θ -Cartan subalgebras in $\mathfrak{h}^\perp$ are conjugate under $\mathbf{H}$ ([Lev07, Thm 2.11]); we will usually fix the choice

$$(2.1) \quad \mathfrak{a} = \left\{ \begin{pmatrix} & \delta \\ \delta & \end{pmatrix} \middle| \delta \text{ is a diagonal } n \times n \text{ matrix} \right\}.$$

The theory of spherical varieties endows the θ -Cartan $\mathfrak{a}$ with a root system of type B_n , whose roots on the choice of Cartan (2.1) are described by

$$\Phi_{\mathbf{X}, \mathfrak{a}} = \{\pm\delta_i^* \pm \delta_j^* \mid 1 \leq i < j \leq n\} \cup \{\pm\delta_i^* \mid 1 \leq i \leq n\},$$

where δ_i^* is the coordinate function for the i -th entry of the $n \times n$ diagonal matrix δ . We may see this as a renormalization of the restricted root system of $\mathbf{X}$.

We have a Weyl group $W_{\mathfrak{a}}$ for the root system $\Phi_{\mathbf{X}, \mathfrak{a}}$, which is presented as $W_{\mathfrak{a}} = N_{\mathbf{G}}(\mathfrak{a})/C_{\mathbf{G}}(\mathfrak{a}) = N_{\mathbf{H}}(\mathfrak{a})/C_{\mathbf{H}}(\mathfrak{a})$. For general symmetric varieties $W_{\mathfrak{a}}$ is a subquotient of the usual Weyl group W , but in our case, $W_{\mathfrak{a}}$ is the subgroup of W given by $W_{\mathfrak{a}} = \mathfrak{S}_n \rtimes \{\pm 1\}^n$. The action of $W_{\mathfrak{a}}$ on (2.1) is by letting $\mathfrak{S}_n$ permute the diagonal entries of δ and letting $\{\pm 1\}^n$ change the signs of the entries δ_i of δ .

There is a construction also of the canonical torus of the symmetric variety, which arises as follows. Let $\mathbf{T}$ denote the canonical torus for $\mathbf{G}$. Then, for any choice of Borel B such that $B \cap \theta(B)$ is a torus, $\mathbf{T} = B/[B, B]$ acts on $\mathbf{H} \backslash \mathbf{G}/[B, B]$ through the quotient $\mathbf{A}$, which we call the *canonical torus* for $\mathbf{X} = \mathbf{H} \backslash \mathbf{G}$. The canonical torus $\mathbf{A}$ comes equipped with a root system $\Phi_{\mathbf{X}}$ and an action of a Weyl group $W_{\mathbf{X}}$. The map on Lie algebras $\mathfrak{a} \rightarrow \text{Lie}(\mathbf{A})$ induced by any choice of Borel B as above is an isomorphism which identifies the two respective root systems, and so we do not distinguish between the two Lie algebras.

We also have a notion of regularity and semi-simplicity for elements of $\mathfrak{h}^\perp$.

Definition 2.2. We say $x \in \mathfrak{h}^\perp$ is *semisimple* if it is semisimple as an element of $\mathfrak{g}$.

Let $I \subset \mathfrak{h}^\perp \times \mathbf{H}$ denote the group scheme of centralizers, i.e.

$$I_x = \{h \in \mathbf{H} \mid h \cdot x = x\}.$$

Then, we say that $x \in \mathfrak{h}^\perp$ is *regular* if $\dim I_x$ is minimal. We denote the regular locus of $\mathfrak{h}^\perp$ by $(\mathfrak{h}^\perp)^{\text{reg}}$. We denote the regular semi-simple locus by $(\mathfrak{h}^\perp)^{\text{rs}}$.

While the notion of regularity for $x \in \mathfrak{h}^\perp$ under the action of $\mathbf{H}$ is generally distinct from regularity of $x \in \mathfrak{g}$ under the action of $\mathbf{G}$, the two notions agree for $\mathbf{X} = \text{GL}_n \times \text{GL}_n \backslash \text{GL}_{2n}$.

The following result traces back to Kostant and Rallis in characteristic zero [KR71] and to Paul Levy in characteristic $p \neq 2$ [Lev07].

Theorem 2.3. Let $\mathfrak{c} := \mathfrak{h}^\perp // \mathbf{H} := \text{Spec}(k[\mathfrak{h}^\perp]^\mathbf{H})$ denote the GIT quotient of $\mathfrak{h}^\perp$ by $\mathbf{H}$, and let $\chi: \mathfrak{h}^\perp \rightarrow \mathfrak{c}$ denote the quotient map.

- (1) ([Lev07, Thm 4.9]) There is an isomorphism $\mathfrak{c} \simeq \mathfrak{a} // W_X$.
- (2) ([Lev07, Lem 6.30]) The map χ admits a section ϵ , which is valued in $(\mathfrak{h}^\perp)^{\text{reg}}$. Moreover, this section can be made $\mathbb{G}_m$ -equivariant after base change by $t \mapsto t^2$.

We refer to such a section $\epsilon: \mathfrak{c} \rightarrow (\mathfrak{h}^\perp)^{\text{reg}}$ as a Kostant–Rallis section.

Returning to our centralizer group scheme $I \subset \mathfrak{h}^\perp \times \mathbf{H}$, we call the restriction of I to $(\mathfrak{h}^\perp)^{\text{reg}}$ the regular centralizer group scheme and denote it by I^{reg} . We have the following package describing I^{reg} .

Proposition 2.4. ([HM25, Prop 4.1]) Recall that $\chi: \mathfrak{h}^\perp \rightarrow \mathfrak{c}$ denotes the quotient map; we also denote $\chi^{\text{reg}} = \chi|_{(\mathfrak{h}^\perp)^{\text{reg}}}$.

- (1) The regular centralizer group scheme I^{reg} is a smooth commutative group scheme over $(\mathfrak{h}^\perp)^{\text{reg}}$ which descends to a smooth commutative group scheme J over $\mathfrak{c}$.
- (2) There is a canonical map $\chi^* J \rightarrow I$ extending the isomorphism $(\chi^{\text{reg}})^* J \simeq I^{\text{reg}}$.

The group scheme J admits a useful description in terms of spectral covers. We begin by recalling some facts and definitions from invariant theory (cf. [HM25]). Recall that $\mathfrak{c}_{2n} = \mathfrak{gl}_{2n} // \text{GL}_{2n} \simeq \mathfrak{t}_{\text{GL}_{2n}} // \mathfrak{S}_{2n}$ admits a spectral cover $\mathfrak{s}_{2n} = \mathfrak{t}_{\text{GL}_{2n}} // \mathfrak{S}_{2n-1}$. Explicitly, if

$$A_{2n} := k[\mathfrak{c}_{2n}] = k[a_1, \dots, a_{2n}]$$

for $a_i(X) = \text{tr } \bigwedge^i X$ the i -th elementary symmetric function of the eigenvalues of X , then

$$k[\mathfrak{s}_{2n}] = A_{2n}[x]/(x^{2n} + a_1 x^{2n-1} + \dots + a_{2n})$$

The natural morphism $\mathfrak{s}_{2n} \rightarrow \mathfrak{c}_{2n}$ is finite and flat of degree $2n$.

Lemma 2.5. [LXZ25a, Lemma 5.1] *The GIT quotient $\mathfrak{c} = \mathfrak{h}^\perp // \mathbf{H}$ is an affine space $\mathfrak{c} = \mathbb{A}^n$. More explicitly, for $x = \begin{pmatrix} X \\ Y \end{pmatrix} \in \mathfrak{h}^\perp$,*

$$\chi(x) = (b_1(x), \dots, b_n(x)),$$

where $b_i(x) = (-1)^i \text{tr } \bigwedge^i (XY)$. In particular, there is a commutative diagram

$$\begin{array}{ccc} \mathfrak{h}^\perp & \longrightarrow & \mathfrak{gl}_{2n} \\ \downarrow \chi & & \downarrow \chi_{2n} \\ \mathfrak{c} & \longrightarrow & \mathfrak{c}_{2n}, \end{array}$$

with the natural map $\mathfrak{c} \rightarrow \mathfrak{c}_{2n}$ given by setting $b_i = a_{2i}$ and $a_i = 0$ for all odd i . In particular $\mathfrak{c} \simeq \mathbb{A}^n$ is naturally equipped with a $\mathbb{G}_m$ -action by the exponents $2, 4, \dots, 2n$.

Writing $A = k[\mathfrak{c}] = k[a_2, a_4, \dots, a_{2n}]$, the base change $\mathfrak{s} = \mathfrak{s}_{2n} \times_{\mathfrak{c}_{2n}} \mathfrak{c}$ is thus a finite, flat, degree- $2n$ cover of $\mathfrak{c}$ with ring of functions

$$k[\mathfrak{s}] = A[x]/(x^{2n} + a_2 x^{2n-2} + \dots + a_{2n})$$

The spectral cover $\mathfrak{s}$ therefore carries an involution ι , defined over $\mathfrak{c}$, given by $\iota(x) = -x$, and the quotient $\mathfrak{s}/\iota$ remains a finite, flat cover of degree n . We have the following description of centralizers.

Proposition 2.6. ([HM25, Prop. 6.2]) *We have an isomorphism*

$$J \simeq \text{Res}_{\mathfrak{c}}^{\mathfrak{s}/\iota}(\mathbb{G}_m)$$

of group schemes over $\mathfrak{c}$.

2.2. Rational structures in invariant theory. The group $\mathbf{H} = \text{GL}_n \times \text{GL}_n$ and $\mathbf{H}$ representation $\mathfrak{h}^\perp$ admit involutions, which in turn will allow us to deduce specific rational forms. First, we have two involutions on $\mathbf{H}$:

- The involution $\tau_1: \mathbf{H} \rightarrow \mathbf{H}$ given by the transpose-inverse $h = (h_1, h_2) \mapsto {}^t h^{-1} = ({}^t h_1^{-1}, {}^t h_2^{-1})$;
- The involution $\tau_2: \mathbf{H} \rightarrow \mathbf{H}$ given by the swapping map on the factors $h = (h_1, h_2) \mapsto ({}^t h_2^{-1}, {}^t h_1^{-1})$.

We have involutions on $\mathfrak{h}^\perp$ corresponding to each of the τ_j above. Namely, if we identify $\mathfrak{h}^\perp \simeq \{(A, B) \mid A, B \text{ } n \times n \text{ matrices}\}$, then we have:

- The involution $\tau_1: \mathfrak{h}^\perp \rightarrow \mathfrak{h}^\perp$ given by the swapping-transpose map $(A, B) \mapsto (-{}^t B, -{}^t A)$.
- The involution $\tau_2: \mathfrak{h}^\perp \rightarrow \mathfrak{h}^\perp$ given by the transpose map $(A, B) \mapsto (-{}^t A, -{}^t B)$.

The involutions τ_i on $\mathbf{H}$ and $\mathfrak{h}^\perp$ are compatible in the sense that $\tau_i(h \cdot x) = \tau_i(h) \cdot \tau_i(x)$ for all $h \in \mathbf{H}$, $x \in \mathfrak{h}^\perp$.

We now define k rational forms of the invariant theoretic spaces of the previous section. Let $k = \mathbb{F}_q$ be a finite field, and let k' be a quadratic extension of k . We denote by σ the Frobenius element of k , thought of as the generator in $\text{Gal}(k'/k)$. Consider the Weil restrictions

$$\mathbf{G}' = \text{Res}_{k'/k} \mathbf{G}_{k'} \quad \text{and} \quad \mathbf{H}' = \text{Res}_{k'/k} \mathbf{H}_{k'}$$

The involution σ acts on $\mathbf{G}'$ and $\mathbf{H}'$, and hence also on $(\mathfrak{h}')^\perp \subset \mathfrak{g}' = \text{Lie}(\mathbf{G}')$. The involutions τ_i similarly induce involutions on $\mathbf{H}'$ and $(\mathfrak{h}')^\perp$ which we call by the same name, and which commute with the action of σ . We form the two rational twists:

- The split-inert or unitary case:

$$H^{\mathbf{U}} = (\mathbf{H}')^{\tau_1 \sigma} \quad \text{and} \quad V^{\mathbf{U}} = (\mathfrak{h}'^\perp)^{\tau_1 \sigma}$$

- The inert-inert or Siegel case:

$$H^{\mathbf{S}} = (\mathbf{H}')^{\tau_2 \sigma} \quad \text{and} \quad V^{\mathbf{S}} = (\mathfrak{h}'^\perp)^{\tau_2 \sigma}$$

More precisely, set $g^* = {}^t \sigma(g)$ for any matrix g . We have the unitary group U_n consisting of $g \in \text{GL}_n(k')$ such that $g \cdot g^* = 1$. Recall that $\text{Herm}_n = \{x \in \mathfrak{gl}_n(k') \mid x = x^*\}$ is the space of $n \times n$ Hermitian matrices.

The above construction produces the following spaces.

- the split-inert/Siegel case Here, $H^{\mathbf{S}} = \text{Res}_{k'/k} \text{GL}_n$ acts on $V^{\mathbf{S}} = \text{Herm}_n \times \text{Herm}_n$ by $g \cdot (x, y) = (gxg^*, (g^*)^{-1}yg^{-1})$. This is the infinitesimal model of the symmetric variety $X^{\mathbf{S}} = \text{Res}_{k'/k} \text{GL}_n \backslash U_{2n}$.
- the inert-inert/unitary case Here $H^{\mathbf{U}} = U_n \times U_n$ acts on $V^{\mathbf{U}} = \mathfrak{gl}_n(k')$ by $(g_1, g_2) \cdot x = g_1 x g_2^*$. This is the infinitesimal model of the symmetric variety $X^{\mathbf{U}} = (U_n \times U_n) \backslash U_{2n}$.

Remark 2.7. The terminology of *split-inert* and *inert-inert* comes from [LXZ25a, LXZ25b], and is motivated by a more general bi-quadratic setup. In this paper, we opt for the self-contained terminology of the *Siegel* and *unitary* cases as we do not discuss this more general setting.

We keep the same notation as Section 2.1 with $V^\star$ replacing $\mathfrak{h}^\perp$ (for $\star = \mathbf{S}$ or $\mathbf{U}$). The following is an easy variation on Lemma 2.5.

Lemma 2.8. [LXZ25a, Lemma 5.3] *The GIT quotients $\mathfrak{c}$ for each of $X^{\mathbf{S}}$ and $X^{\mathbf{U}}$ above are isomorphic as k -varieties. Setting $\mathfrak{c} = V^\star // H^\star$ for this quotient, we have $\mathfrak{c} \simeq \mathbf{A}^n$ is an affine space over k equipped with a $\mathbb{G}_m$ -action by the exponents $2, 4, \dots, 2n$.*

Moreover, the following result identifies the regular centralizer group schemes for $X^{\mathbf{U}}$ and $X^{\mathbf{S}}$.

Theorem 2.9. *Consider the k rational form $X^\star$ for $\star \in \{\mathbf{U}, \mathbf{S}\}$. There exists a canonical smooth commutative group k -scheme $J = J^\star$ over $\mathfrak{c}$ such that we have an isomorphism $(\chi^{\text{reg}})^* J = I^{\text{reg}}$ defined over k which extends uniquely to a morphism $\chi^* J \rightarrow I$. Moreover, J satisfies the following:*

- (1) *The group schemes J for $X^{\mathbf{U}}$ and $X^{\mathbf{S}}$ are canonically isomorphic.*
- (2) *J can be explicitly described as the kernel of the norm map*

$$\text{Nm}: \text{Res}_{\mathfrak{c}}^{(\mathfrak{s}/\iota)_{k'}}(\mathbb{G}_m) \rightarrow \text{Res}_{\mathfrak{c}}^{\mathfrak{s}/\iota}(\mathbb{G}_m),$$

which takes an S point $\xi \in \Gamma\left(S \times_{\mathfrak{c}} (\mathfrak{s}/\iota)_{k'}, \mathcal{O}_{S \times_{\mathfrak{c}} (\mathfrak{s}/\iota)_{k'}}^\times\right)$ to $\xi \cdot \sigma(\xi)$.

Proof. By Proposition 2.4, if we write $\check{I}^{\text{reg}} = I^{\text{reg}} \otimes_k \bar{k}$, then there is a unique group scheme $\check{J}$ such that $(\chi^{\text{reg}})^* \check{J} \simeq \check{I}^{\text{reg}}$, with a unique extension to a map $(\chi^{\text{reg}})^* \check{J} \rightarrow \check{I}$. Moreover, by Proposition 2.6, we may write $\check{J} = \text{Res}_{\mathfrak{c}}^{\mathfrak{s}/\iota}(\mathbb{G}_m)$.

Consider the k -model $J^\dagger$ of $\check{J}$ given by the following Frobenius action: Let σ act diagonally on $\mathbb{G}_m \times \mathfrak{s}/\iota$, with the action on $\mathbb{G}_m$ by inversion and the action on $\mathfrak{s}/\iota$ induced by the natural action of σ on $\mathfrak{s}$ such that $\sigma(x) = -x$. We note that $J^\dagger$ is simply the kernel of the norm map as in (2) above. It is clear that $J^\dagger$ is smooth and commutative. We claim that over the regular semisimple locus $(\chi^{\text{rs}})^* J^\dagger \simeq I^{\text{rs}}$. Indeed, for either case if we use our chosen model of $G = \text{U}_{2n}$, one verifies that the centralizer $C \subset H^*$ of $\mathfrak{a} \cap V^*$ for $\mathfrak{a}$ as in (2.1) is a group scheme over k identified with $C(\bar{k}) \simeq \mathbb{G}_m(\bar{k})^n$ with Frobenius acting by inversion composed with the Galois action on coefficients. Since all regular semisimple elements are conjugate, and since $\mathfrak{a} \cap V^* \neq \emptyset$ for $\star \in \{\mathbf{U}, \mathbf{S}\}$, we get our stated identification on the regular semisimple locus.

Now, since $\check{I}^{\text{reg}}$ is smooth and separated, it admits a unique k -model agreeing with I^{rs} over $\mathfrak{c}^{\text{rs}}$. Since $\chi^* J^\dagger$ and I^{reg} are two such k models, we conclude $(\chi^{\text{reg}})^* J^\dagger \simeq I^{\text{reg}}$. Moreover, since $J^\dagger$ is smooth, there exists a canonical extension $\chi^* J^\dagger \rightarrow I$. Hence, the result is proven with the k model $J := J^\dagger$ of $\check{J}$. $\square$

While the GIT quotient is independent of form, the existence of Kostant-Rallis sections is sensitive to which k form of $\mathbf{X}$ we consider.

Theorem 2.10. *The section in Theorem 2.3(2) descends to a section*

$$\mathfrak{c} := V^{\mathbf{S}} // H^{\mathbf{S}} \rightarrow V^{\mathbf{S}, \text{reg}}$$

over k for $X^{\mathbf{S}}$. However, such a section does not exist for $X^{\mathbf{U}}$.

Proof. Suppose that e is a regular, nilpotent element in $V^*(\bar{k})$, and let λ be an associated character for e valued in $H^*(\bar{k})$, i.e. $\text{Ad}(\lambda(t))e = t^2 e$ for all $t \in \bar{k}^\times$. Such an associated character exists by [Lev07, Corollary 5.4]. By [Lev07, Lemma 6.30], a section to the map $\chi: V^* \rightarrow \mathfrak{c}$ over the algebraic closure $\bar{k}$ is given by $e + \mathfrak{v}$ where e is a regular nilpotent element in $\mathfrak{h}^\perp$ and $\mathfrak{v}$ is any choice of λ graded subspace of $\mathfrak{h}^\perp$ such that a slightly enhanced version of the complementary property

$$[e, \mathfrak{h}] \oplus \mathfrak{v} = \mathfrak{h}^\perp$$

holds. (“Enhanced” here means that it remains complementary in a smaller subspace of $\mathfrak{h}^\perp$. See [Lev07, §6.5] for a precise statement.)

Suppose that we can find a k rational point $e \in V^*(k)$ which is regular and nilpotent. Then since k is perfect of characteristic good for G , by [McN04, Theorem 26], we can further assume that the associated cocharacter λ is defined over k , and so takes values in H^* . As $[e, \mathfrak{h}] \subset \mathfrak{h}^\perp$ is stable under the Frobenius action, we can therefore choose $\mathfrak{v}$ to be stable under Frobenius as well. In this case, Levy’s section over $\bar{k}$ is stable under Frobenius and hence descends to a section over k . Note that the existence of a section over k implies the existence of a k rational regular, nilpotent element e , so we have proved that the existence of a section over k is equivalent to the existence of a regular, nilpotent element over k .

It reduces, therefore, to study for which rational forms there exists a k -rational regular nilpotent element $e \in V^*(k)$. Let $\mathcal{N}^* \subset V^*$ denote the nilpotent cone in V^* . Geometrically, there are two H -orbits of regular nilpotent elements $\mathcal{N}^{*, \text{reg}}$, represented by the elements

$$\begin{pmatrix} & e_n \\ I_n & \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} & I_n \\ e_n & \end{pmatrix},$$

where $e_n \in \mathfrak{gl}_n$ is regular nilpotent. It is not hard to see that the Frobenius element permutes these two orbits in the unitary case and stabilizes each them in the Siegel case. Hence in the unitary case $\mathcal{N}^{\mathbf{U}, \text{reg}}(k) = \emptyset$ and in the Siegel case $\mathcal{N}^{\mathbf{S}, \text{reg}}(k) \neq \emptyset$. The result follows. $\square$

We will fix a choice of k -rational section for $X^{\mathbf{S}}$ for the rest of this paper, denoting it by ϵ and referring to it as a *Kostant–Rallis section*.

2.3. Endoscopic symmetric varieties. The *endoscopic symmetric varieties* for a rational form $X^\star$ were studied by the second author for $X^\mathbf{U}$ in [Les25], where a fundamental lemma was proven for $X^\mathbf{U}$ over a p -adic field, and for general symmetric varieties satisfying mild assumptions in [Les24]. We summarize the latter theory here for the special case of $X^\mathbf{S}$.

Let $\mathbf{T}$ denote the canonical torus for G . We have a canonical involution θ on the torus $\mathbf{T}$ whose centralizer $\mathbf{C} = \mathbf{T}^\theta$ is identified with the centralizer subgroup of a generic regular, semisimple element of $\mathfrak{h}^\perp$ (see [Les21, Proposition 1.12] and [HM25, Lemma 4.13]). Let $\widehat{\mathbf{G}}$ denote the Langlands dual group of G , considered over $\mathbb{C}$, with dual torus $\widehat{\mathbf{T}}$. Recall that we have a canonical torus $\mathbf{A}$ associated to the symmetric variety $\mathbf{X}$, which fits into an exact sequence

$$1 \rightarrow \mathbf{C} \rightarrow \mathbf{T} \rightarrow \mathbf{A} \rightarrow 1.$$

This exact sequence dualizes to an exact sequence

$$1 \rightarrow \widehat{\mathbf{A}} \rightarrow \widehat{\mathbf{T}} \rightarrow \widehat{\mathbf{C}} \rightarrow 1.$$

Recall the dual group $\widehat{\mathbf{G}}_X$ associated to $\mathbf{X}$, which in our case is the subgroup $\widehat{\mathbf{G}}_X = \mathrm{Sp}_{2n}$ of $\widehat{\mathbf{G}} = \mathrm{GL}_{2n}$. The homogeneous variety $\widehat{\mathbf{X}} = \widehat{\mathbf{G}}_X \backslash \widehat{\mathbf{G}}$ is itself a symmetric variety for an involution $\widehat{\theta}$. Moreover, the torus $\widehat{\mathbf{T}}$ acts on $\widehat{\mathbf{X}}$ freely through the quotient $\widehat{\mathbf{T}} \rightarrow \widehat{\mathbf{C}}$, so that the above sequence extends to a commutative diagram of

$$\begin{array}{ccccc} \widehat{\mathbf{A}} & \longrightarrow & \widehat{\mathbf{T}} & \longrightarrow & \widehat{\mathbf{C}} \\ \downarrow & & \downarrow & & \downarrow \\ \widehat{\mathbf{G}}_X & \longrightarrow & \widehat{\mathbf{G}} & \longrightarrow & \widehat{\mathbf{X}}. \end{array}$$

The variety $\widehat{\mathbf{X}}$ is a quasi-Hamiltonian variety, admitting a group-valued moment map

$$s: \widehat{\mathbf{X}} \rightarrow \widehat{\mathbf{G}}, \quad \widehat{\mathbf{G}}_X \cdot g \mapsto g^{-1} \cdot \widehat{\theta}(g).$$

We will refer to the map s as the *symmetrization map* for $\widehat{\mathbf{X}}$. Note that s is almost a section of the canonical map $\widehat{\mathbf{T}} \rightarrow \widehat{\mathbf{C}}$ in the sense that s is injective and the composition

$$\widehat{\mathbf{C}} \xrightarrow{s} \widehat{\mathbf{T}} \rightarrow \widehat{\mathbf{C}}$$

is the map $x \mapsto x^2$.

We now recall the classification of elliptic endoscopic data for the unitary group U_{2n} . Let $\mathbf{G} = \mathrm{GL}_{2n}$ and ρ_G be the $\mathrm{Out}(\mathbf{G})$ -torsor which defines the unitary group U_{2n} . For $\kappa \in \widehat{\mathbf{T}}$, we may take the neutral component of the centralizer $\widehat{\mathbf{G}}_\kappa$ of κ in $\widehat{\mathbf{G}}$, and similarly for $(\widehat{\mathbf{G}} \rtimes \mathrm{Out}(\mathbf{G}))_\kappa$ in $\widehat{\mathbf{G}} \rtimes \mathrm{Out}(\mathbf{G})$. Let $\pi_0(\kappa)$ be the group of connected components of $(\widehat{\mathbf{G}} \rtimes \mathrm{Out}(\mathbf{G}))_\kappa$, which is a finite group fitting into a short exact sequence

$$1 \rightarrow \widehat{\mathbf{G}}_\kappa \rightarrow (\widehat{\mathbf{G}} \rtimes \mathrm{Out}(\mathbf{G}))_\kappa \rightarrow \pi_0(\kappa) \rightarrow 1.$$

The group $\pi_0(\kappa)$ admits morphisms

$$(2.2) \quad \mathrm{Out}(\widehat{\mathbf{G}}_\kappa) \leftarrow \pi_0(\kappa) \rightarrow \mathrm{Out}(\widehat{\mathbf{G}})$$

where the rightward map is induced by the projection onto the second factor and the leftward map is induced by the action of $(\widehat{\mathbf{G}} \rtimes \mathrm{Out}(\mathbf{G}))_\kappa$ on $\widehat{\mathbf{G}}_\kappa$.

Let F be either a non-archimedean local field with residue field k or a function field $F = k(\Sigma)$ for Σ a smooth, projective, geometrically-connected curve over k of genus at least 2. Consider a quadratic extension F'/F . For example, in the case of a function field, such an extension is determined by an étale, degree 2 cover

$$\vartheta: \Sigma' \rightarrow \Sigma,$$

with Σ' a smooth, projective, geometrically-connected curve over k . Let $\sigma \in \text{Gal}(F'/F)$ be the canonical degree 2 involution, and let $G = \text{U}_{2n}$ be the associated unitary group. That is, if F is a non-archimedean local field,

$$G = \text{U}_{2n} = \{g \in \text{GL}(F') \mid g \cdot {}^t\sigma(g) = 1\},$$

while if $F = k(\Sigma)$ is a function field,

$$G = \vartheta_*(\text{GL}_{2n} \times \Sigma')^{\tau\sigma}$$

where τ is the transpose map on GL_{2n} and σ acts through the automorphism group of the cover. This group G is determined by an $\text{Out}(\mathbf{G})$ -torsor ρ_G over F .

Definition 2.11. Fix $x \in \Sigma(k)$ for the function field case and $x \in \Sigma^\bullet = \text{Spec}(F)$ in the case of a non-archimedean local field. An *endoscopic datum* for the unitary group U_{2n} is a pair $\mathfrak{e} = (\kappa, \rho)$, where $\kappa \in \hat{\mathbf{T}}$ and ρ is a homomorphism

$$\rho: \pi_1(\Sigma, x) \rightarrow \pi_0(\kappa)$$

which induces ρ_G via the rightward map in (2.2).

We call an endoscopic datum (κ, ρ) *elliptic* if $(Z(\hat{\mathbf{G}}_\kappa)^{\pi_1(\Sigma, x)})^\circ = \{1\}$, where $(\bullet)^\circ$ denotes the neutral component and $\pi_1(\Sigma, x)$ acts through ρ .

If the unitary group U_{2n} is defined over the local field $F = k((\varpi))$ for k a finite field, then it is easy to verify that (κ, ρ) is elliptic if and only if $\kappa \in \hat{\mathbf{C}}$ is 2-torsion. Moreover, in this case, the map $\pi_0(\kappa) \rightarrow \text{Out}(\mathbf{G})$ is a bijection of order-2 groups. Hence, the data (κ, ρ) is completely determined by the data of κ in such cases.

An endoscopic datum determines a quasisplit group $G^\mathfrak{e}$ over Σ by considering the induced quasisplit group from the split group $\mathbf{G}_\kappa$ dual to $\hat{\mathbf{G}}_\kappa$ together with the $\text{Out}(\mathbf{G}_\kappa)$ -torsor induced by ρ_κ via the leftward arrow in (2.2). The associated endoscopic groups $G^\mathfrak{e}$ for F a non-archimedean field are the unitary groups $\text{U}_r \times \text{U}_s$ for $r + s = 2n$.

We now add the data of a relative endoscopic datum.

Definition 2.12. An *endoscopic datum* for a symmetric variety $X^\mathbf{S}$, or *relative endoscopic datum*, is a pair (κ, ρ) , where $\kappa \in \hat{\mathbf{C}}$ and $(s(\kappa), \rho)$ is an endoscopic datum for the unitary group $G = \text{U}_{2n}$.

We say that (κ, ρ) is *elliptic* if $(s(\kappa), \rho)$ is elliptic as an endoscopic datum for U_{2n} .

In [Les24], the second author produced a symmetric space $Y^\mathfrak{e}$ for $G^\mathfrak{e}$, which satisfies the following properties.

Theorem 2.13. ([Les24, Theorem 1.4]) *For any elliptic relative endoscopic datum (κ, ρ) for X a “well adapted” symmetric space, there exists a symmetric space*

$$Y^\mathfrak{e} = H_{Y^\mathfrak{e}} \backslash G^\mathfrak{e},$$

unique up to $G^\mathfrak{e}$ inner form, where $G^\mathfrak{e}$ is the endoscopic group of G determined by the endoscopic datum $(s(\kappa), \rho)$, such that

- (1) *The canonical torus for $Y^\mathfrak{e}$ is identified with the canonical torus for X ;*
- (2) *$\hat{\mathbf{G}}_{Y^\mathfrak{e}}$ is the centralizer of κ in $\hat{\mathbf{G}}_X$;*
- (3) *There is a canonical map $\text{Aut}_d(X) \rightarrow \text{Aut}_d(Y^\mathfrak{e})$ between the groups of doubling automorphisms (see §3.4 of loc cit.) such that the geometric 1-cocycle (see §3.4.3 of loc cit.) of $Y^\mathfrak{e}$ is induced from the geometric 1-cocycle of X via this map.*
- (4) ([Les24, Proposition 8.5]) *There is canonical transfer morphism*

$$\nu_\mathfrak{e}: \mathfrak{c}_\mathfrak{e} \rightarrow \mathfrak{c} = T^*X // G,$$

where $\mathfrak{c}_\mathfrak{e} = (\mathfrak{g}^\mathfrak{e})^{\theta^\mathfrak{e}=-1} // H_{Y^\mathfrak{e}}$, which is finite and flat.

Conditions (1) through (3) above are needed to describe Y^ϵ up to G^ϵ inner form, but only conditions (1) and (4) will be needed in this paper.

We will now specialize to the case $X = X^{\mathbf{S}}$ and explicitly describe the transfer morphism ν_ϵ of Theorem 2.13. The reader unfamiliar with [Les24] can, for the purposes of this paper, take the following as a definition.

Consider an elliptic relative endoscopic datum (κ, ρ) for $X^{\mathbf{S}}$. By our earlier remarks, $\kappa \in \widehat{\mathbf{C}}$ must be 2 torsion, and ρ is completely determined by κ . View $\kappa = (\kappa_1, \dots, \kappa_n)$ as an n -tuple with $\kappa_i \in \{\pm 1\}$, and suppose that κ has r indices $+1$ and s indices -1 (with $r + s = n$). Then,

$$Y^\epsilon = X_r^{\mathbf{S}} \times X_s^{\mathbf{S}} := (\mathrm{GL}_r(F') \backslash \mathrm{U}_{2r}) \times (\mathrm{GL}_s(F') \backslash \mathrm{U}_{2s}).$$

For κ having $\kappa_i = 1$ for r indices and $\kappa_i = -1$ for s indices, the transfer map ν_ϵ is the morphism

$$\mathfrak{c}_r \times \mathfrak{c}_s \simeq \mathfrak{a} // ((\mathfrak{S}_r \times \mathfrak{S}_s) \rtimes \{\pm 1\}^n) \rightarrow \mathfrak{a} // W_X$$

with subscripts r and s denoting the rank of the symmetric variety factor, and the final map induced by the embedding $\mathfrak{S}_r \times \mathfrak{S}_s \subset \mathfrak{S}_n$ given by the distribution of $+1$'s and -1 's in κ . It is evident that this morphism is finite and flat.

Let $\mathfrak{s}_\epsilon \rightarrow \mathfrak{c}_\epsilon$ denote the spectral cover for $\mathfrak{c}_\epsilon$, with involution ι_ϵ , which is finite, flat of degree $2n$ over $\mathfrak{c}_\epsilon$. The following characterizes $\mathfrak{s}_\epsilon$ and its relationship to $\mathfrak{s}$.

Proposition 2.14. *There is a canonical partial normalization map $\mathfrak{s}_\epsilon \rightarrow \mathfrak{s} \times_{\mathfrak{c}} \mathfrak{c}_\epsilon$, which induces a canonical morphism $\nu_\epsilon^* J \rightarrow J_\epsilon$. The latter map of regular centralizers is an isomorphism over $\nu_\epsilon^{-1}(\mathfrak{c} \setminus \mathfrak{D}^{\mathrm{cent}})$.*

Proof. If $X^\epsilon = X_r^{\mathbf{S}} \times X_s^{\mathbf{U}}$ for $r + s = n$, then identify $\mathfrak{c}_\epsilon \simeq \mathfrak{c}_r \times \mathfrak{c}_s$, with $\mathfrak{c}_r$ the GIT quotient for $X_r^{\mathbf{S}}$ and $\mathfrak{c}_s$ for $X_s^{\mathbf{U}}$. Let $\mathfrak{s}_r \rightarrow \mathfrak{c}_r$ and $\mathfrak{s}_s \rightarrow \mathfrak{c}_s$ denote the respective spectral covers. Then, we have

$$k[\mathfrak{s}_\epsilon] = k[\mathfrak{s}_r] \times k[\mathfrak{s}_s],$$

so that $\mathfrak{s}_\epsilon = \mathfrak{s}_r \sqcup \mathfrak{s}_s$. Let $P_r(x), P_s(x)$ be the polynomials in x such that $k[\mathfrak{s}_r] = A[x]/P_r(x)$ and $k[\mathfrak{s}_s] = A[x]/P_s(x)$. The coefficients of P_r, P_s are the coordinate functions on $\mathfrak{c}_r, \mathfrak{c}_s$. Therefore, the morphism $\mathfrak{c}_r \times \mathfrak{c}_s \rightarrow \mathfrak{c}_n$, being induced by the inclusion $\mathfrak{S}_r \times \mathfrak{S}_s \subset \mathfrak{S}_n$, induces on any point $a \in \nu_\epsilon(\mathfrak{c}_\epsilon)$ a canonical factorization

$$x^{2n} + a_2 x^{2n-2} + \dots + a_{2n} = P_r(x) \cdot P_s(x)$$

so that $k[\mathfrak{s} \times_{\mathfrak{c}} \mathfrak{c}_\epsilon] = A[x]/(P_r(x) \cdot P_s(x))$. Hence, we obtain a ι equivariant morphism

$$\mathfrak{s}_\epsilon \rightarrow \mathfrak{s} \times_{\mathfrak{c}} \mathfrak{c}_\epsilon$$

over $\mathfrak{c}_\epsilon$ which is the partial normalization $\mathfrak{s}_r \sqcup \mathfrak{s}_s \rightarrow \mathfrak{s}_r \cup \mathfrak{s}_s$ at the intersection of $\mathfrak{s}_r$ and $\mathfrak{s}_s$ in $\mathfrak{s} \times_{\mathfrak{c}} \mathfrak{c}_\epsilon$.

In particular, we also deduce a partial normalization map

$$\mathfrak{s}_\epsilon / \iota_\epsilon \rightarrow \mathfrak{s} / \iota \times_{\mathfrak{c}} \mathfrak{c}_\epsilon$$

Since $\mathrm{Res}_{\mathfrak{c}_\epsilon}^{\mathfrak{s}_\epsilon / \iota_\epsilon}(\mathbb{G}_m) = \mathrm{Res}_{\mathfrak{s} / \iota \times_{\mathfrak{c}} \mathfrak{c}_\epsilon}^{\mathfrak{s}_\epsilon / \iota_\epsilon} \left(\mathrm{Res}_{\mathfrak{c}_\epsilon}^{\mathfrak{s} / \iota \times_{\mathfrak{c}} \mathfrak{c}_\epsilon}(\mathbb{G}_m) \right)$, we obtain the morphism $\nu_\epsilon^* J \rightarrow J_\epsilon$ by the adjunction map for the Weil restriction $\mathrm{Res}_{\mathfrak{s} / \iota \times_{\mathfrak{c}} \mathfrak{c}_\epsilon}^{\mathfrak{s}_\epsilon / \iota_\epsilon}$. In particular, this morphism is an isomorphism over the open set in $\mathfrak{c}_\epsilon$ over which $\mathfrak{s}_r$ and $\mathfrak{s}_s$ are disjoint in $\mathfrak{s} \times_{\mathfrak{c}} \mathfrak{c}_\epsilon$, which includes the preimage of $\mathfrak{c} \setminus \mathfrak{D}^{\mathrm{cent}}$. $\square$

In [LXZ25a], it was noted that Y^ϵ is not the endoscopic space needed to formulate a fundamental lemma for $X^{\mathbf{S}}$. Indeed, the endoscopic variety Y^ϵ does not distinguish between the endoscopic datum for $\kappa = \mathbf{1}_r \times (-\mathbf{1}_s)$ and for $\kappa = (-\mathbf{1}_r) \times \mathbf{1}_s$ (and the corresponding orbital integrals are different and the full endoscopic datum does distinguish between the two).

In [LXZ25a], the authors propose the following endoscopic space, which we give as a definition.

Definition 2.15. For an elliptic relative endoscopic datum $\mathfrak{e} = (\kappa, \rho)$, with $\kappa_i = 1$ for r indices and $\kappa_i = -1$ for s indices, we call the variety

$$X^\mathfrak{e} = H^\mathfrak{e} \backslash G^\mathfrak{e} = (\mathrm{GL}_r(F') \backslash \mathrm{U}_{2r}) \times ((\mathrm{U}_s \times \mathrm{U}_s) \backslash \mathrm{U}_{2s}).$$

the *endoscopic symmetric space* associated to $\mathfrak{e}$.

Since $\mathfrak{c}_\mathfrak{e}$ and $J_\mathfrak{e}$ are the same for $X^\mathfrak{e}$ and $Y^\mathfrak{e}$, there is still a unique transfer morphism $\nu_\mathfrak{e}$ with canonical isomorphism $\nu_\mathfrak{e}^* J \simeq J_\mathfrak{e}$.

Remark 2.16. The definition of $X^\mathfrak{e}$ above is more or less *ad hoc* as it is not clear how to extract the implied comparison of orbital integrals from the formalism of [Les24]. A generalization of *loc. cit.* that accommodates this case more naturally is ongoing work of the second author.

2.4. Divisors in $\mathfrak{c}$. In this section, we introduce several important divisors in the GIT quotient $\mathfrak{c}$ which will control the geometry of our local and global moduli spaces.

We consider first the discriminant divisor of the root system $\Phi_{\mathbf{X}}$. Namely, we consider the function

$$\prod_{\alpha \in \Phi_{\mathbf{X}}} d\alpha \in k[\mathfrak{a}]$$

where we are viewing each root as a function $d\alpha$ on $\mathfrak{a}$. As this function is W_X -invariant, it induces a divisor in $\mathfrak{c}$ which we denote by $\mathfrak{D}_{\mathbf{X}}$ and call the *discriminant divisor*. For the θ -Cartan $\mathfrak{a}$ as in (2.1) we have

$$\mathfrak{D}_{\mathbf{X}} = \left((-1)^{\frac{n^2+n}{2}} \left(\prod_i \delta_i^2 \right) \left(\prod_{i < j} (\delta_i - \delta_j)^2 \right) \right) = \left(a_{2n} \cdot \mathrm{Disc}(x^n + a_2 x^{n-1} + a_4 x^{n-2} + \cdots + a_{2n}) \right)$$

where δ_i is the coordinate function for the i -th diagonal entry in δ in (2.1), and Disc denotes the discriminant of the polynomial in x , which is a polynomial function of the coefficients a_{2i} . We record here the following basic observation about $\mathfrak{D}_{\mathbf{X}}$.

Lemma 2.17. ([Ngo10, Lem 1.10.1]) *The divisor $\mathfrak{D}_{\mathbf{X}}$ is a reduced, effective divisor on $\mathfrak{c}$, and the map $\pi: \mathfrak{a} \rightarrow \mathfrak{c}$ is étale on the complement of $\mathfrak{D}_{\mathbf{X}}$. Moreover, this divisor is stable under the action of $\mathrm{Out}(\mathbf{H})$.*

We have a subdivisor of $\mathfrak{D}_{\mathbf{X}}$ corresponding to the type A_n root system on $\mathfrak{a}$, given by

$$\mathfrak{D}^{\mathrm{cent}} = \left((-1)^{\frac{n^2-n}{2}} \prod_{i < j} (\delta_i - \delta_j)^2 \right) = \left(\mathrm{Disc}(x^n + a_2 x^{n-1} + a_4 x^{n-2} + \cdots + a_{2n}) \right).$$

The centralizer group scheme J is closely related to the type GL_n centralizers (see the spectral description given in Proposition 2.6 and part (2) of Theorem 2.9), and so we call this divisor the *centralizer divisor* associated to $\mathbf{X}$. The following hints at its importance in describing regular centralizers using the spectral cover $\mathfrak{s}$.

Lemma 2.18. *Let $\mathfrak{s}$ denote the spectral cover over $\mathfrak{c}$ described in §2.1. The map $\mathfrak{s}/\iota \rightarrow \mathfrak{c}$ is étale away from the divisor $\mathfrak{D}^{\mathrm{cent}}$.*

Proof. Ramification of the spectral cover $\mathfrak{s} \rightarrow \mathfrak{c}$ can occur over a point $a = (a_i)$ in $\mathfrak{c}$ if and only if the corresponding spectral equation

$$Q_a(x) = x^n + a_2 x^{n-1} + \cdots + a_{2n}$$

has repeated roots. Since $\mathfrak{D}^{\mathrm{cent}}$ is cut out by the discriminant of the polynomial $Q_a(x)$, this is precisely the condition that $a \in \mathfrak{D}^{\mathrm{cent}}$. $\square$

We finally come to the most subtle divisor, the branching divisor for $\mathbf{X}$. In [HM25], it was shown that there exists a divisor $\mathfrak{D}^{\text{br}} \subset \mathfrak{c}$ such that the fibers $\chi^{-1}(x)$ contain exactly 2 regular $\mathbf{H}$ -orbits when $x \in \mathfrak{D}^{\text{br}}$ and exactly 1 regular $\mathbf{H}$ -orbit when $x \in \mathfrak{c} - \mathfrak{D}^{\text{br}}$. We call this divisor the *branching divisor* for $\mathbf{X}$. The following structure result will serve as a model from which the regular parts of our moduli spaces will be understood.

Theorem 2.19. ([HM25, Theorems 3.7 and 3.37]) *Let $(\mathfrak{h}^\perp)^{\text{reg}} // \mathbf{H}$ denote the nonseparated scheme obtained by gluing two copies of $\mathfrak{c}$ away from the branching divisor $\mathfrak{D}^{\text{br}}$, i.e.*

$$(\mathfrak{h}^\perp)^{\text{reg}} // \mathbf{H} = \mathfrak{c} \coprod_{\mathfrak{c} \setminus \mathfrak{D}^{\text{br}}} \mathfrak{c}.$$

Then, the natural map $[\chi^{\text{reg}}]: [(\mathfrak{h}^\perp)^{\text{reg}} / \mathbf{H}] \rightarrow \mathfrak{c}$ factors through

$$[(\mathfrak{h}^\perp)^{\text{reg}} / \mathbf{H}] \rightarrow (\mathfrak{h}^\perp)^{\text{reg}} // \mathbf{H} \rightarrow \mathfrak{c}$$

with the first map being a gerbe banded by the regular centralizer group scheme, trivialized by a choice of Kostant–Rallis section $[\epsilon]$.

The branching divisor $\mathfrak{D}^{\text{br}}$ can be described for more general spherical varieties, and it has a role in the relative Langlands program matching the determinant of the associated symplectic representation for tempered spherical varieties (see [HLM24]). It is a surprising feature of this case that it is complementary to $\mathfrak{D}^{\text{cent}}$ in the full discriminant divisor.

Lemma 2.20. ([HM25, Prop 3.44, 3.45]) *For the spherical variety $\mathbf{X} = (\text{GL}_n \times \text{GL}_n) \backslash \text{GL}_{2n}$, we have $\mathfrak{D}^{\text{br}} = (a_{2n})$ is the divisor given by the determinant. In particular, we have a decomposition of reduced divisors*

$$\mathfrak{D}_{\mathbf{X}} = \mathfrak{D}^{\text{cent}} + \mathfrak{D}^{\text{br}}$$

Finally, we related the divisors above to their endoscopic counterparts via the transfer map. Let X^ϵ be an endoscopic symmetric variety for X , and let $\nu_\epsilon: \mathfrak{c}_\epsilon \rightarrow \mathfrak{c}$ be the corresponding transfer map (defined in §2.3). Choose $\Psi \subset \Phi_{X,\mathfrak{a}} - \Phi_{X^\epsilon,\mathfrak{a}}$ such that for any $\{\pm\alpha\} \subset \Phi_{X,\mathfrak{a}} - \Phi_{X^\epsilon,\mathfrak{a}}$, exactly one of α and $-\alpha$ is in Ψ . Then, by [Ngo10, §1.10.3] the divisor $\nu_\epsilon^* \mathfrak{D}_{\mathbf{X}}$ decomposes as

$$\nu_\epsilon^* \mathfrak{D}_{\mathbf{X}} = \mathfrak{D}_{X^\epsilon} + 2\mathfrak{R}_\epsilon$$

where $\mathfrak{R}_\epsilon = (\prod_{\alpha \in \Psi} \alpha)$ is the (reduced) ramification divisor for the map ν_ϵ . This decomposition admits a further refinement for $\mathfrak{D}^{\text{cent}}$ and $\mathfrak{D}^{\text{br}}$ as follows.

Lemma 2.21. *For $\mathfrak{R}_\epsilon$ as defined above, we have the following decompositions of pullbacks*

$$\nu_\epsilon^* \mathfrak{D}^{\text{cent}} = \mathfrak{D}_\epsilon^{\text{cent}} + 2\mathfrak{R}_\epsilon, \quad \nu_\epsilon^* \mathfrak{D}^{\text{br}} = \mathfrak{D}_\epsilon^{\text{br}}$$

Proof. Write $X^\epsilon = X_m^{\text{U}} \times X_{n-m}^{\text{S}}$. We may explicitly express the map $\nu_\epsilon: \mathfrak{c}_\epsilon \rightarrow \mathfrak{c}$ as

$$\mathfrak{a} // (W_{X_m^{\text{U}}} \times W_{X_{n-m}^{\text{S}}}) = \mathfrak{a} // ((\mathfrak{S}_m \times \mathfrak{S}_{n-m}) \rtimes \{\pm 1\}^n) \rightarrow \mathfrak{a} // W_{\mathfrak{a}}$$

whose ramification is evidently described by a subdivisor of $\mathfrak{D}^{\text{cent}}$. □

3. UNITARY HIGGS BUNDLES

We review here the notation and definitions of the moduli spaces of unitary Higgs bundles of Ngô and Laumon [LN08]. We will use some of their results to deduce properties of the relative Hitchin moduli spaces and their unitary relatives in §4.

First we fix the following data: fix smooth, projective, geometrically-connected curves Σ', Σ over k with an étale degree 2 cover $\vartheta: \Sigma' \rightarrow \Sigma$. Let $\sigma \in \text{Aut}(\Sigma'/\Sigma)$ denote the order-two generator of the automorphism group of the cover. We also fix a divisor D on Σ and let S denote a test k -scheme.

Definition 3.1. The moduli space of U_n Higgs bundles is the functor $\mathcal{N}_n$ which takes values

$$\mathcal{N}_n(S) = \left\{ (\mathcal{E}', h, \Phi') \left| \begin{array}{l} \mathcal{E}' \text{ is a rank } n \text{ vector bundle on } \Sigma' \times S, \\ h: \mathcal{E}' \rightarrow \sigma^*(\mathcal{E}')^\vee \text{ is a Hermitian form on } \mathcal{E}', \text{ i.e. } h = \sigma^* h^\vee, \\ \Phi': \mathcal{E}' \rightarrow \mathcal{E}' \otimes D, \text{ satisfying } \sigma^*(\Phi')^\vee \circ h + h \circ \Phi' = 0 \end{array} \right. \right\}$$

Note that, in the above, $(-)^\vee = \text{Hom}(-, \mathcal{O}_{\Sigma'})$ denotes the usual dual of vector bundles. In [LN08], it is shown that $\mathcal{N}_n$ is representable by a smooth, finite-type algebraic stack.

We now recall the construction of the Hitchin morphism. The involution σ induces a decomposition

$$\vartheta_* \mathcal{O}_{\Sigma'} = \mathcal{O}_\Sigma \oplus \mathcal{L}$$

where $\mathcal{L}$ is the (-1) -eigenspace of σ on $\vartheta_* \mathcal{O}_{\Sigma'}$. Note that $\mathcal{L}^{\otimes 2} \simeq \mathcal{O}_\Sigma$.

Definition 3.2. Let $\mathcal{B}_n$ denote the affine space

$$\mathcal{B}_n = \bigoplus_{i=1}^n H^0(\Sigma, \mathcal{L}(D)^{\otimes i}),$$

where $\mathcal{L}(D) := \mathcal{L} \otimes D$. The Hitchin morphism $f: \mathcal{N}_n \rightarrow \mathcal{B}_n$ is the morphism taking $(\mathcal{E}', h, \Phi')$ to the coefficients of the characteristic polynomial of Φ' , i.e. $\text{tr } \bigwedge^i \Phi' \in H^0(\Sigma, \mathcal{L}(D)^{\otimes i})$.

Fibers of the morphism f have an interpretation via spectral covers similar to those associated to classical groups by Hitchin [Hit87]. To state it, let $\tilde{\Sigma}'$ and $\tilde{\Sigma}$ be the base change of the spectral cover $\mathfrak{s}_n \rightarrow \mathfrak{c}$ along the respective morphisms fitting in the diagram

$$\begin{array}{ccccc} \tilde{\Sigma}' & \longrightarrow & \tilde{\Sigma} & \longrightarrow & (\mathfrak{s}_n)_{\mathcal{L}(D)} \\ \downarrow \pi'_a & & \downarrow \pi_a & & \downarrow \\ \Sigma' \times \mathcal{B}_n & \xrightarrow{\vartheta} & \Sigma \times \mathcal{B}_n & \xrightarrow{\text{ev}} & (\mathfrak{c}_n)_{\mathcal{L}(D)}. \end{array}$$

For every $a \in \mathcal{B}_n(S)$, we denote by $\pi'_a: \tilde{\Sigma}'_a \rightarrow \Sigma'$ the corresponding finite, flat cover of Σ' , and similarly for $\pi_a: \tilde{\Sigma}_a \rightarrow \Sigma$. The automorphism $\sigma \in \text{Aut}(\Sigma'/\Sigma)$ acts naturally on $\tilde{\Sigma}'$, and it is easy to verify that $\tilde{\Sigma}_a = \tilde{\Sigma}'_a/\sigma$ under this action.

Convention 3.3. For $\mathcal{G}$ a sheaf on Σ' , we denote

$$\mathcal{G}^\vee = \text{Hom}_{\mathcal{O}_{\Sigma'}}(\mathcal{G}, \mathcal{O}_{\Sigma'}),$$

while for $\mathcal{F}$ a sheaf on a spectral cover $\tilde{\Sigma}'_a$ for $a \in \mathcal{A}(S)$, we denote

$$\mathcal{F}^\vee = \text{Hom}_{\mathcal{O}_{\tilde{\Sigma}'_a}}(\mathcal{F}, \omega_{\tilde{\Sigma}'_a/\Sigma' \times S}).$$

where $\omega_{\tilde{\Sigma}'_a/\Sigma' \times S}$ is the relative dualizing sheaf for the finite, flat cover $\tilde{\Sigma}'_a \rightarrow \Sigma' \times S$.

Proposition 3.4. Let $\mathcal{N}_a = f^{-1}(a)$. For all $a \in \mathcal{B}_n(S)$ such that $\tilde{\Sigma}'_a$ is reduced, we have

$$\mathcal{N}_a \simeq \left\{ (\mathcal{F}, h) \mid \mathcal{F} \in \overline{\text{Pic}}(\tilde{\Sigma}'_a/S), h: \mathcal{F} \rightarrow \sigma^*(\mathcal{F})^\vee \text{ Hermitian, i.e. } h = \sigma^*(h)^\vee \right\}$$

where the $\overline{\text{Pic}}$ denotes the compactified relative Picard scheme classifying torsion-free, generic rank-1 sheaves on $\tilde{\Sigma}_a$, and the superscript indicates that we are considering the sub-functor classifying $\mathcal{F}'$ such that $\mathcal{F}' \simeq \sigma^*(\mathcal{F}')^\vee$.

Given a torsion-free, generic rank-1 sheaf $\mathcal{F}'$, we construct the corresponding unitary Higgs bundle by taking $\pi'_{a,*}(\mathcal{F}')$, with the isomorphism $\mathcal{F}' \simeq \sigma^*(\mathcal{F}')^\vee$ providing the Hermitian structure h and multiplication by the tautological section providing the section Φ' .

The entire moduli space is acted on by its group of symmetries

$$\mathcal{P}_\mathcal{N} = \text{Pic}(\tilde{\Sigma}'/\mathcal{B}_n)^{\sigma^* = (-)^\vee}$$

in the natural way.

Definition 3.5. The *anisotropic locus* $\mathcal{B}_n^{\text{ani}} \subset \mathcal{B}_n$ is the set of $a \in \mathcal{B}_n$ such that $\pi_0(\mathcal{P}_{\mathcal{N},a})$ is finite; that is, such that the morphism $\tilde{\Sigma}'_a \rightarrow \tilde{\Sigma}_a$ induces a bijection on irreducible components. (See [LN08, Lem 2.8.2].)

It was shown in [LN08] that $\mathcal{B}_n^{\text{ani}}$ is a nonempty, open subset of $\mathcal{B}_n$; the restriction $\mathcal{N}_n^{\text{ani}} = f^{-1}(\mathcal{B}_n^{\text{ani}})$ is a Deligne–Mumford stack; and the Hitchin morphism $f^{\text{ani}} = f|_{\mathcal{N}_n^{\text{ani}}}$ is proper.

4. THE RELATIVE HITCHIN MODULI SPACE

4.1. Global moduli spaces. We keep the same notation as in §3 and formulate the global forms of the rational symmetric spaces in Section 2.2. Namely, let Σ be a smooth, geometrically-connected projective curve of genus $g \geq 2$ over the finite field $k = \mathbb{F}_q$, and let $\vartheta: \Sigma' \rightarrow \Sigma$ be an étale 2-cover over k , with Σ' also smooth, projective, and geometrically connected. Let $\text{Aut}(\Sigma'/\Sigma) = \langle \sigma \rangle$ for σ an involution.

The involution σ acts on the Weil restrictions

$$\mathbf{G}' = \vartheta_* \mathbf{G} \quad \text{and} \quad \mathbf{H}' = \vartheta_* \mathbf{H},$$

and hence also on the bundle $(\mathfrak{h}')^\perp \subset \mathfrak{g}' = \text{Lie}(\mathbf{G}')$. The involutions τ_i induce involutions on $\mathbf{H}'$ and $(\mathfrak{h}')^\perp$ which we call by the same name, and which commute with the action of σ . We form the two twists $(H^{\mathbf{U}}, V^{\mathbf{U}})$ and $(H^{\mathbf{S}}, V^{\mathbf{S}})$ as before.

In each case, $V^\star$ admits a $\mathbb{G}_m$ -action by scaling, which commutes with the action of $H^\star$, and hence, we may form the stack $[V_D^\star/H^\star]$ over Σ , where $V_D^\star = V^\star \times^{\mathbb{G}_m} D^\times$ is the D -twisted form of V' . Here and below, the superscript $\star$ denotes either $\star = \mathbf{U}$ for the split-inert case (using τ_1) or $\star = \mathbf{S}$ for the inert-inert case (using τ_2). Both are rational twists of the case

$$H = (\mathbf{H}')^\sigma = \mathbf{H} \times \Sigma \quad \text{and} \quad V = (\mathfrak{h}'^\perp)^\sigma = \mathfrak{h}^\perp \times \Sigma.$$

Define the moduli stack $\mathcal{M}^\star = \mathcal{M}_D^\star$ as the mapping stack

$$\mathcal{M}_D^\star = \text{Maps}(\Sigma, [V_D^\star/H^\star]).$$

Explicitly, we may describe $\mathcal{M}^\star$ as functors of points classifying certain unitary Higgs bundles with additional data.

Lemma 4.1. *The moduli $\mathcal{M}^\star$ represent the functors sending a scheme S to*

$$(4.1) \quad \mathcal{M}^{\mathbf{U}}(S) = \left\{ (\mathcal{E}'_1, \mathcal{E}'_2, h_1, h_2, \phi') \left| \begin{array}{l} \mathcal{E}'_i \text{ are rank } n \text{ vector bundles on } \Sigma' \times S, \\ h_i: \mathcal{E}'_i \rightarrow \sigma^*(\mathcal{E}'_i)^\vee \text{ is a Hermitian form, i.e. } \sigma^* h_i^\vee = h_i, \\ \text{and } \phi': \mathcal{E}'_1 \rightarrow \mathcal{E}'_2 \otimes D \text{ such that } \sigma^*(\phi')^\vee \circ h_1 + h_2 \circ \phi' = 0 \end{array} \right. \right\}$$

$$(4.2) \quad \mathcal{M}^{\mathbf{S}}(S) = \left\{ (\mathcal{E}', h, \phi'_{12}, \phi'_{21}) \left| \begin{array}{l} \mathcal{E}' \text{ is a rank } n \text{ vector bundle on } \Sigma' \times S, \phi'_{12}: \mathcal{E}' \rightarrow \sigma^*(\mathcal{E}')^\vee \otimes D \\ \text{and } \phi'_{21}: \sigma^*(\mathcal{E}')^\vee \rightarrow \mathcal{E}' \otimes D \text{ such that } \sigma^* \phi'_{ij}^\vee = \phi'_{ij} \\ \text{and } h: \mathcal{E}' \rightarrow \sigma^*(\mathcal{E}')^\vee \text{ is a Hermitian form on } \mathcal{E}' \end{array} \right. \right\}.$$

We will describe these moduli spaces in terms of spectral covers in §4.5. First we show how they arise from taking fixed points under the appropriate involutions on the stack $\mathcal{N}_{2n}$.

For $\star \in \{\mathbf{U}, \mathbf{S}\}$, we define the maps $\mu^\star: \mathcal{M}^\star \rightarrow \mathcal{N}_{2n}$ as follows. In the unitary case, the map $\mu^{\mathbf{U}}$ is given on S -points by

$$(4.3) \quad \mu^{\mathbf{U}}(\mathcal{E}'_1, \mathcal{E}'_2, h_1, h_2, \phi') = (\mathcal{E}'_1 \oplus \mathcal{E}'_2, h_1 \oplus h_2, \Phi')$$

where the Higgs field Φ' is given by the sum $\phi' \oplus \sigma^*(\phi')^\vee$, viewing the latter as a map $\mathcal{E}'_2 \rightarrow \mathcal{E}'_1 \otimes D$ by

$$\mathcal{E}'_2 \xrightarrow{h_2} \sigma^*(\mathcal{E}'_2)^\vee \xrightarrow{\sigma^*(\phi')^\vee} \sigma^*(\mathcal{E}'_1)^\vee \otimes D \xrightarrow{h_1} \mathcal{E}'_1 \otimes D.$$

It is now simple to check that $\mu^{\mathbf{U}}(\mathcal{E}'_1, \mathcal{E}'_2, h_1, h_2, \phi') \in \mathcal{N}_{2n}(S)$.

In the Siegel case, the map $\mu^{\mathbf{S}}$ is given on S -points by

$$(4.4) \quad \mu^{\mathbf{S}}(\mathcal{E}', h, \phi'_{12}, \phi'_{21}) = (\mathcal{E}' \oplus \sigma^*(\mathcal{E}')^\vee, h \oplus \sigma^*(h)^\vee, \phi'_{12} \oplus \phi'_{21}) \in \mathcal{N}_{2n}(S).$$

4.2. Fixed Points on $\mathcal{N}_{2n}$. Denote by $\mathcal{N} = \mathcal{N}_{2n}$ the moduli space of unitary Higgs bundles of rank $2n$ (see §3). To relate the moduli spaces defined in the previous section to fixed point stacks, we first consider involutions at the level of invariant theory. Recall that $G = \mathrm{U}_{2n}$ is the unitary group over k , defined relative to the quadratic extension k' . Let $\theta^* : G \rightarrow G$ denote the involution for which $H^* = G^{\theta^*}$. More precisely, write $G = \mathrm{U}(V)$ where (V, h) is our fixed Hermitian space over k with $h : V \xrightarrow{\sim} \sigma^*(V)^\vee$; we sometimes write $\langle \cdot, \cdot \rangle : V \otimes V \rightarrow k'$ for the induced σ -sesquilinear form

$$\langle v, w \rangle := h(v)(\sigma^*(w)).$$

For any $g \in \mathrm{GL}(V)$ set $g^\dagger \in \mathrm{GL}(\sigma^*(V)^\vee)$ such that

$$g^\dagger(h(v))(\sigma^*(w)) = h(v)(\sigma^*(g^{-1}w)) = \langle v, g^{-1}w \rangle;$$

then $g \in \mathrm{U}(V)$ if and only if the diagram

$$\begin{array}{ccc} V & \xrightarrow{h} & \sigma^*(V)^\vee \\ \downarrow g & & \downarrow g^\dagger \\ V & \xrightarrow{h} & \sigma^*(V)^\vee \end{array}$$

commutes.

Let $J^* : V \rightarrow V$ be the involution inducing an eigenspace decomposition $V = V_+ \oplus V_-$ such that

(1) $(\star = \mathbf{U})$ $\langle J^{\mathbf{U}}v, J^{\mathbf{U}}w \rangle = \langle v, w \rangle$. This implies that

$$J^{\mathbf{U}, \dagger} \circ h = h \circ J^{\mathbf{U}},$$

which in particular forces the restriction of h to $V_\pm$ is a non-degenerate Hermitian form.

(2) $(\star = \mathbf{S})$ $\langle J^{\mathbf{S}}v, w \rangle + \langle v, J^{\mathbf{S}}w \rangle = 0$. Note that this implies $\langle J^{\mathbf{S}}v, J^{\mathbf{S}}w \rangle = -\langle v, w \rangle$ since $(J^{\mathbf{S}})^2 = I$.

This implies that

$$J^{\mathbf{S}, \dagger} \circ h = -h \circ J^{\mathbf{S}},$$

and that the restriction of h to $V_\pm$ vanishes so that $V_- \simeq V_+^*$.

There is an evident involution $\theta^* : [\mathfrak{g}/G] \rightarrow [\mathfrak{g}/G]$ induced by $\theta^*(g) = J^*gJ^{*, -1}$ and $\theta^*(x) = -J^*xJ^{*, -1}$ on the Lie algebra.

Recall that an S -point of $[\mathfrak{g}/G]$ consists of the data of a triple $(\mathcal{E}', h, \phi')$, with $\mathcal{E}'$ a vector bundle on $S \otimes_k k'$, $\phi : \mathcal{E}' \rightarrow \mathcal{E}'$ a linear map, and $h : \mathcal{E}' \rightarrow \sigma^*(\mathcal{E}')^\vee$ a Hermitian form on $\mathcal{E}'$, i.e. $h = \sigma^*(h)^\vee$.

Lemma 4.2. *There is an automorphism $[\mathfrak{g}/G] \rightarrow [\mathfrak{g}/G]$ which brings θ^* to the involution ι^* which sends $(\mathcal{E}', h, \phi') \in [\mathfrak{g}/G](S)$ to*

$$\iota^*(\mathcal{E}', h, \phi') = \begin{cases} (\mathcal{E}', -h, -\phi') & \text{if } \star = \mathbf{S} \\ (\mathcal{E}', h, -\phi') & \text{if } \star = \mathbf{U} \end{cases}$$

Proof. To a triple $(\mathcal{E}', h, \phi') \in [\mathfrak{g}/G](S)$, θ^* induces a triple $(\theta^*(\mathcal{E}'), \theta^*(h), \theta^*(\phi'))$ as follows: Fix an étale covering $\{U_\alpha \rightarrow S \otimes_k k'\}$ which is stable under the involution σ , with local trivializations $\nu_\alpha : \mathcal{E}'|_{U_\alpha} \xrightarrow{\sim} V \otimes \mathcal{O}_\alpha$ and descent data $\{g_{\alpha, \beta}\}$. By definition the diagrams

$$\begin{array}{ccc} V \otimes \mathcal{O}_{\alpha, \beta} & \xrightarrow{h_{\alpha, \beta}} & \sigma^*(V)^\vee \otimes \mathcal{O}_{\alpha, \beta} \\ \downarrow g_{\alpha, \beta} & & \downarrow g_{\alpha, \beta}^\dagger \\ V \otimes \mathcal{O}_{\alpha, \beta} & \xrightarrow{h_{\alpha, \beta}} & \sigma^*(V)^\vee \otimes \mathcal{O}_{\alpha, \beta} \end{array}$$

commute for all α, β . Then,

- (1) the bundle $\theta_*^\star(\mathcal{E}')$ is obtained from the descent data $\{J^\star g_{\alpha,\beta} J^{\star,-1}\}$. Intuitively, the local sections ν_α induce local sections $J^\star v$ of $\theta_*^\star(\mathcal{E}')$.
- (2) the map $\theta_*^\star(\phi') : \theta_*^\star(\mathcal{E}') \rightarrow \theta_*^\star(\mathcal{E}')$ is obtained by locally conjugating by $J^\star$ and gluing together via $\{J^\star g_{\alpha,\beta} J^{\star,-1}\}$
- (3) We first claim that there is a natural isomorphism

$$\theta_*^\star(\sigma^*(\mathcal{E}')^\vee) \simeq \sigma^*(\theta_*^\star(\mathcal{E}'))^\vee.$$

To see this, identify $\{g_{\alpha,\beta}^\dagger\}$ as the descent data for $\sigma^*(\mathcal{E}')^\vee$. Then the isomorphism follows from the matrix identity

$$J^{\star,\dagger} g_{\alpha,\beta}^\dagger (J^{\star,-1})^\dagger = (J^\star g_{\alpha,\beta} J^{\star,-1})^\dagger.$$

With this identity, the map $\theta_*^\star(h) : \theta_*^\star(\mathcal{E}') \rightarrow \sigma^*(\theta_*^\star(\mathcal{E}'))^\vee$ is obtained by gluing the local maps $\theta_*^\star(h)|_{U_\alpha} \equiv h$. We note that is well defined since

$$(J^\star g_{\alpha,\beta} J^{\star,-1})^\dagger \circ h = h \circ (J^\star g_{\alpha,\beta} J^{\star,-1}),$$

which can be checked using the compatibility between $J^\star$ and h .

From the above, we see that the adjoint action by $J^{\star,-1}$ realizes an isomorphism $\theta_*^\star(\mathcal{E}) \simeq \mathcal{E}$ which identifies $\theta_*^\star(\phi)$ with $-\phi$. Finally, the Hermitian form h is carried to $J^{\star,\dagger} h J^{\star,-1} = \pm h$, with negative sign for $\star = \mathbf{S}$ and positive for $\star = \mathbf{U}$ due to the relations between $J^\star$ and h above the lemma. $\square$

As a consequence of Lemma 4.2, we will work with the more explicit involutions $\iota^\star$. The involutions $\iota^\star$ induce involutions on the moduli of unitary Higgs bundles $\mathcal{N}$ by sending $(\mathcal{E}', h, \Phi') \in \mathcal{N}(S)$ to

$$(4.5) \quad \iota^\star(\mathcal{E}', h, \Phi') = \begin{cases} (\mathcal{E}', -h, -\Phi') & \text{if } \star = \mathbf{S} \\ (\mathcal{E}', h, -\Phi') & \text{if } \star = \mathbf{U} \end{cases}$$

Form the homotopy fixed point stack $\mathcal{M}^{\star,b} = \mathcal{N}^{\iota^\star}$ as the fixed points in $\mathcal{N}$ of $\mathbb{Z}/2\mathbb{Z}$, with nontrivial element acting by $\iota^\star$. Explicitly, for a test scheme S ,

$$(4.6) \quad \mathcal{M}^{\star,b}(S) = \left\{ (\mathcal{E}', h, \Phi', \alpha) \left| \begin{array}{l} (\mathcal{E}', h, \Phi') \in \mathcal{N}(S) \\ \alpha : (\mathcal{E}', h, \Phi') \simeq \iota^\star(\mathcal{E}', h, \Phi') \\ \iota^\star \alpha \circ \alpha = Id_{(\mathcal{E}', h, \Phi')} \end{array} \right. \right\}$$

where in the above, α is an isomorphism in the groupoid $\mathcal{N}(S)$. The functor $\mathcal{M}^{\star,b}$ is representable by an algebraic stack by [Rom05, Theorem 3.3] and [LN08, Prop 2.2.1]. Moreover, since $\mathcal{N}$ is locally of finite type, $\mathcal{M}^{\star,b}$ is also locally of finite type by [Rom05, Proposition 3.7].

For $r + s = 2n$, let $\mathcal{M}_{r,s}^{\mathbf{U}}$ denote the moduli space whose S -points are

$$\mathcal{M}_{r,s}^{\mathbf{U}}(S) = \left\{ (\mathcal{E}'_1, \mathcal{E}'_2, h_1, h_2, \phi') \left| \begin{array}{l} \mathcal{E}'_1, \mathcal{E}'_2 \text{ are rank } r, s \text{ vector bundles on } \Sigma' \times S, \text{ respectively,} \\ h_i : \mathcal{E}'_i \rightarrow \sigma^*(\mathcal{E}'_i)^\vee \text{ is a Hermitian form, i.e. } \sigma^* h_i^\vee = h_i, \\ \text{and } \phi' : \mathcal{E}'_1 \rightarrow \mathcal{E}'_2 \otimes D \text{ such that } \sigma^*(\phi')^\vee \circ h_1 + h_2 \circ \phi' = 0 \end{array} \right. \right\}$$

Proposition 4.3. *There are isomorphisms $\mathcal{M}^{\mathbf{S},b} \simeq \mathcal{M}^{\mathbf{S}}$ and*

$$\mathcal{M}^{\mathbf{U},b} \simeq \bigsqcup_{r+s=2n} \mathcal{M}_{r,s}^{\mathbf{U}}.$$

In particular, $\mathcal{M}^{\mathbf{U}} = \mathcal{M}_{n,n}^{\mathbf{U}} \subset \mathcal{M}^{\mathbf{U},b}$ is a distinguished (disjoint) component.

Proof. Consider the data of a point $(\mathcal{E}', h, \Phi', \alpha) \in \mathcal{M}^{\star,b}(S)$ as in the right-hand side of (4.6). We can consider $\mathcal{E}' = \mathcal{E}'_1 \oplus \mathcal{E}'_2$, with $\mathcal{E}'_1$ and $\mathcal{E}'_2$ being the $+1$ and -1 eigenspaces of α acting on $\mathcal{E}'$,

respectively. Then, the condition on α implies that the following diagram commutes

$$\begin{array}{ccc} \mathcal{E}' & \xrightarrow{\alpha} & \mathcal{E}' \\ \Phi' \downarrow & & \downarrow -\Phi' \\ \mathcal{E}' \otimes D & \xrightarrow{\alpha(D)} & \mathcal{E}' \otimes D \end{array}$$

In particular, restricting the top left corner to $\mathcal{E}'_i \subset \mathcal{E}'$, we see that Φ' must be in the form $\Phi' = \Phi'_{12} \oplus \Phi'_{21}$ where $\Phi'_{ij}: \mathcal{E}'_i \rightarrow \mathcal{E}'_j \otimes D$.

Now, we specialize to the case $\star = \mathbf{S}$. In this case, the Hermitian form h satisfies the following condition: let $\langle \cdot, \cdot \rangle: \mathcal{E}' \otimes \sigma^*(\mathcal{E}') \rightarrow \mathcal{O}_{X'}$ denote the pairing associated to h . Then the following diagram commutes

$$\begin{array}{ccc} \mathcal{E}' \otimes \sigma^*(\mathcal{E}') & \xrightarrow{\langle \cdot, \cdot \rangle} & \mathcal{O}_{X'} \\ \alpha \otimes \sigma^* \alpha \downarrow & \nearrow -\langle \cdot, \cdot \rangle & \\ \mathcal{E}' \otimes \sigma^*(\mathcal{E}') & & \end{array}$$

In particular, for all open sets $U \subset \Sigma'$ and all sections $s \in \Gamma(\mathcal{E}'_i, U')$ and $t \in \Gamma(\mathcal{E}'_j, U')$ ($i, j \in \{1, 2\}$), we have

$$(4.7) \quad \langle s, t \rangle = -\langle (-1)^{i-1}s, (-1)^{j-1}t \rangle = (-1)^{i+j-1}\langle s, t \rangle.$$

That is, the condition on h is equivalent to requiring

$$\langle \cdot, \cdot \rangle \in (\mathcal{E}'_1 \otimes \mathcal{E}'_2)^* \oplus (\mathcal{E}'_2 \otimes \mathcal{E}'_1)^*$$

which is to say that h determines an isomorphism $h: \mathcal{E}'_2 \xrightarrow{\sim} \sigma^*(\mathcal{E}'_1)^\vee$. This allows us to identify the tuple $(\mathcal{E}', h, \Phi', \alpha)$ with the equivalent data $(\mathcal{E}'_1, h, \Phi'_{12}, \Phi'_{21}) \in \mathcal{M}^{\mathbf{S}}(S)$.

Consider now the case $\star = \mathbf{U}$. For this case, the condition on h is that the pairing $\langle \cdot, \cdot \rangle$ fit into the commutative diagram

$$\begin{array}{ccc} \mathcal{E}' \otimes \sigma^*(\mathcal{E}') & \xrightarrow{\langle \cdot, \cdot \rangle} & \mathcal{O}_{X'} \\ \alpha \otimes \sigma^* \alpha \downarrow & \nearrow \langle \cdot, \cdot \rangle & \\ \mathcal{E}' \otimes \sigma^*(\mathcal{E}') & & \end{array}$$

This condition implies that, for all open sets $U \subset \Sigma'$ and all sections $s \in \Gamma(\mathcal{E}'_i, U')$ and $t \in \Gamma(\mathcal{E}'_j, U')$ ($i, j \in \{1, 2\}$), we have

$$(4.8) \quad \langle s, t \rangle = -\langle (-1)^{i-1}s, (-1)^{j-1}t \rangle = (-1)^{i+j}\langle s, t \rangle.$$

so that

$$\langle \cdot, \cdot \rangle \in (\mathcal{E}'_1 \otimes \mathcal{E}'_1)^* \oplus (\mathcal{E}'_2 \otimes \mathcal{E}'_2)^*.$$

In other words, h is a direct sum of Hermitian forms h_i on $\mathcal{E}'_i$. From the relation

$$\sigma^*(\Phi')^\vee \circ h + h \circ \Phi' = 0$$

in the definition of unitary Higgs bundles, it follows that Φ' is completely determined by h_1, h_2 , and Φ'_{12} , and so the data of $(\mathcal{E}', h, \Phi', \alpha)$ is equivalent to the data of $(\mathcal{E}'_1, \mathcal{E}'_2, h_1, h_2, \Phi'_{12})$. depending on the ranks of $\mathcal{E}'_1$ and $\mathcal{E}'_2$, this determines a point in $\mathcal{M}_{r,s}^{\mathbf{U}}$.

It is not hard to show the maps on S -points above are functorial, fully faithful, and essentially surjective. $\square$

4.3. The relative Hitchin fibration. We now introduce the relative Hitchin fibration, whose geometry is governed by the invariant theory of §2.1.

Definition 4.4. The relative Hitchin base $\mathcal{A} = \mathcal{A}_D$ is the space of sections

$$\mathcal{A} = \Gamma(\Sigma, \mathfrak{c}_{\mathcal{L}(D)}) = \bigoplus_{i=1}^n H^0(\Sigma, \mathcal{L}(D)^{\otimes 2i}) = \bigoplus_{i=1}^n H^0(\Sigma, D^{\otimes 2i})$$

where $\mathfrak{c}_{\mathcal{L}(D)} = \mathfrak{c} \times^{\mathbb{G}_m} \mathcal{L}(D)$, and we note in the last equality that $\mathcal{L}^{\otimes 2} = \mathcal{O}$.

As $\mathfrak{c}_{\mathcal{L}(D)} = \mathfrak{c}_D$, we will use these two interchangeably. Note that the second equality relies on Lemma 2.5.

The Hitchin base $\mathcal{A}$ is an affine space, and composition with the map $[\chi]: [\mathfrak{h}^\perp/\mathbf{H}] \rightarrow \mathfrak{c}$ induces morphisms

$$f^*: \mathcal{M}^* \rightarrow \mathcal{A},$$

which we call the *relative Hitchin map* for X^* . On S -points, the maps are explicitly given by (using notation as in §4)

$$\begin{aligned} f^{\mathbf{U}}(\mathcal{E}'_1, \mathcal{E}'_2, h_1, h_2, \phi') &= \text{char.poly}(\sigma^*(\phi')^\vee \circ \phi') \\ f^{\mathbf{S}}(\mathcal{E}', h, \phi'_{12}, \phi'_{21}) &= \text{char.poly}(\phi'_{12} \circ \phi'_{21}) \end{aligned}$$

Note that although the constructions differ, Lemma 2.8 ensures that the Hitchin bases for $\mathcal{M}^{\mathbf{U}}$ and $\mathcal{M}^{\mathbf{S}}$ are indeed identified.

Given a choice of D' such that $(D')^2 = D$, for the form $X^{\mathbf{S}}$, Theorem 2.10 along with an argument mirroring [Ngo06, Proposition 2.5] produces a Kostant-Rallis section which globalizes to a section of $h^{\mathbf{S}}$, which we denote by

$$(4.9) \quad [\epsilon] = [\epsilon_{D'}]: \mathcal{A} \rightarrow \mathcal{M}^{\mathbf{S}}.$$

Note that no such section exists in the unitary case $X^{\mathbf{U}}$.

4.4. Spectral curves. Recall that $\mathfrak{s} \rightarrow \mathfrak{c}$ denotes the universal spectral cover restricted to $\mathfrak{c}$, see §2.1. Note that $\mathfrak{s}$ carries a canonical $\mathbb{G}_m$ -action (given by scaling x) and the map $\mathfrak{s} \rightarrow \mathfrak{c}$ is $\mathbb{G}_m$ -equivariant with respect to this action. Hence we may form the twist $\mathfrak{s}_D = \mathfrak{s} \times^{\mathbb{G}_m} D^\times$.

Recall that in §3, we defined the spectral covers $\tilde{\Sigma}'_a$ and $\tilde{\Sigma}_a$ associated to a point $a \in \mathcal{A}(S)$ by taking the base change of the following diagram

$$\begin{array}{ccccc} \tilde{\Sigma}' & \longrightarrow & \tilde{\Sigma} & \longrightarrow & \mathfrak{s}_D \\ \downarrow & & \downarrow & & \downarrow \\ \Sigma' \times \mathcal{A} & \longrightarrow & \Sigma \times \mathcal{A} & \xrightarrow{\text{ev}} & \mathfrak{c}_D \end{array}$$

The involution $\iota(x) = -x$ on $\mathfrak{s}$ induces an involution, which we also denote by ι , on $\tilde{\Sigma}'$ and $\tilde{\Sigma}$ by base change. We denote by $\bar{\Sigma}' = \tilde{\Sigma}'/\iota$ and $\bar{\Sigma} = \tilde{\Sigma}/\iota$. The action of $\sigma \in \text{Aut}(\Sigma'/\Sigma)$ lifts to an involution on $\tilde{\Sigma}'_a$ by letting $\sigma(x) = -x$ while σ acts via the Galois action on coefficients. As previously noted, $\tilde{\Sigma}_a = \tilde{\Sigma}'_a/\sigma$. In particular, for any $a \in \mathcal{A}(S)$, we have a commutative diagram of spectral curves

$$(4.10) \quad \begin{array}{ccc} \tilde{\Sigma}'_a & \xrightarrow{/\sigma} & \tilde{\Sigma}_a \\ \downarrow / \iota & & \downarrow \\ \bar{\Sigma}'_a & \longrightarrow & \bar{\Sigma}_a, \end{array}$$

with each arrow being an étale 2-cover over k . There is clearly a universal such diagram over $\Sigma \times \mathcal{A}$ pulling back to this one for any $a \in \mathcal{A}(S)$

For a general $a \in \mathcal{A}(S)$, the spectral cover $\tilde{\Sigma}_a$ need not be smooth or even reduced. To control the behavior of $\tilde{\Sigma}_a$, we will now introduce the smooth locus $\mathcal{A}^\diamond$ and the heart locus $\mathcal{A}^\heartsuit$ in the Hitchin base. The notation and definitions below are chosen to mirror those of [Ngo10, §4.5, §4.7].

Definition 4.5. Let $\text{ev}: \Sigma \times \mathcal{A} \rightarrow \mathfrak{c}_D$ denote the evaluation map. We denote by $\mathcal{A}^\heartsuit$ the projection of $\text{ev}^{-1}((\mathfrak{c} \setminus \mathfrak{D}^{\text{cent}})_D)$ to $\mathcal{A}$. That is, $\mathcal{A}^\heartsuit \subset \mathcal{A}$ is the open subvariety whose $\bar{k}$ -points are given by

$$\mathcal{A}^\heartsuit(\bar{k}) = \{a \in \mathcal{A}(\bar{k}) \mid a(\Sigma_{\bar{k}}) \not\subset \mathfrak{D}_D^{\text{cent}}\}.$$

Likewise, the *smooth locus* $\mathcal{A}^\diamond$ is the open subvariety of $\mathcal{A}$ consisting of $a \in \mathcal{A}(S)$ whose associated spectral curve $\tilde{\Sigma}'_a$ is smooth and irreducible. It contains all points a of $\mathcal{A}$ whose image $a(\Sigma)$ meets the divisor $\mathfrak{D}_D^{\text{cent}}$ transversely.

We denote $\mathcal{M}^{\star, \heartsuit} = (f^\star)^{-1}(\mathcal{A}^\heartsuit)$ and $\mathcal{M}^{\star, \diamond} = (f^\star)^{-1}(\mathcal{A}^\diamond)$.

Lemma 4.6. *Assume that $\deg(D) > 2g$. The smooth locus $\mathcal{A}^\diamond$ is a nonempty open subset of $\mathcal{A}$, and $\mathcal{A}^\diamond \subset \mathcal{A}^\heartsuit$.*

Proof. The only nontrivial claim is that $\mathcal{A}^\diamond$ is nonempty. Since $a \in \mathcal{A}^\diamond(S)$ holds in particular whenever a intersects the full divisor $\mathfrak{D}_X = \mathfrak{D}^{\text{cent}} + \mathfrak{D}^{\text{br}}$ transversely, an identical argument to [Ngo10, Prop 4.7.1] proves this. $\square$

Similar to [Ngo10, §4.5 and §4.7], we have the following interpretations of $\mathcal{A}^\heartsuit$ in terms of the spectral cover $\tilde{\Sigma}'_a$, with identical proof.

Proposition 4.7. *If $a \in \mathcal{A}^\heartsuit(k)$, the spectral curve $\tilde{\Sigma}'_a$ (and hence also $\tilde{\Sigma}_a$, $\bar{\Sigma}'_a$, $\bar{\Sigma}_a$) is reduced over k .*

4.5. Spectral description of relative Hitchin fibers. In this section we interpret the moduli spaces $\mathcal{M}^\star$ in terms of appropriate spectral covers. Recall our Convention 3.3.

Proposition 3.4 gives the description of $\mathcal{N}^\heartsuit$ in terms of spectral curve

$$(4.11) \quad \mathcal{N}^\heartsuit(S) = \{(a, \mathcal{F}', h) \mid a \in \mathcal{B}^\heartsuit(S), \mathcal{F}' \in \overline{\text{Pic}}(\tilde{\Sigma}'_a), h: \mathcal{F}' \xrightarrow{\sim} \sigma^*(\mathcal{F}')^\vee \text{ a Hermitian form}\}.$$

Recall that a triple $(a, \mathcal{F}', h)$ gives a U_{2n} Higgs bundle by considering the vector bundle $E' = \pi'_{a,*}(\mathcal{F}')$ on $\Sigma' \times S$ (with $\pi'_a: \tilde{\Sigma}'_a \rightarrow \Sigma' \times S$ the covering map) together with the Hermitian form $E \rightarrow \sigma^*(E)^\vee$ obtained by descent from h . The section Φ' is obtained by multiplication by the tautological section.

We have an involution on $\mathcal{B}$ given by sending $a = (a_1, \dots, a_{2n})$ to $\iota_{\mathcal{B}}(a) = ((-1)^i a_i)$. The fixed points $\mathcal{B}^{\iota_{\mathcal{B}}}$ gives the relative Hitchin base $\mathcal{A}$. Recall further that we are denoting by ι the involution on the spectral curve $\tilde{\Sigma}'_a$ defined over Σ' (given by negating the tautological section).

Lemma 4.8. *The involution $\iota^\star$ defined in (4.5) restricts to the involution on $\mathcal{N}^\heartsuit$ sending the data $(a, \mathcal{F}', h) \in \mathcal{N}^\heartsuit(S)$ in the right hand side of (4.11) to*

$$\iota^\star(a, \mathcal{F}', h) = \begin{cases} (\iota_{\mathcal{B}}(a), \iota^* \mathcal{F}', -\iota^* h) & \text{if } \star = \mathbf{S} \\ (\iota_{\mathcal{B}}(a), \iota^* \mathcal{F}', \iota^* h) & \text{if } \star = \mathbf{U}. \end{cases}$$

Proof. For spectral data $(a, \mathcal{F}', h) \in \mathcal{N}^\heartsuit(S)$ in the right hand side of (4.11), let $(\mathcal{E}', h, \Phi')$ be the associated U_{2n} Higgs bundle. We have $\pi'_{a,*}(\mathcal{F}') = \pi'_{a,*}(\iota^* \mathcal{F}')$ since $\pi'_a \circ \iota = \pi'_a$. Moreover, ι has the effect of negating the tautological section, so that multiplication by the tautological section on $\mathcal{E}' = \pi'_{a,*}(\iota^* \mathcal{F}')$ is $-\Phi'$. The Hermitian form $\pm \iota^* h$ descends to $\pm h$ on $\mathcal{E}'$, and so we deduce that $\iota^\star(a, \mathcal{F}', h)$ corresponds exactly to $\iota^\star(\mathcal{E}', h, \Phi')$. $\square$

The following result can be seen as a unitary version of [Sch15, Proposition 2.2].

Corollary 4.9. *The functor $\mathcal{M}^{\star, \heartsuit}$ is the homotopy fixed point stack of $\mathcal{N}^{\heartsuit}$ under the action of $\iota^{\star}$. In particular, $\mathcal{M}^{\star, \heartsuit}$ is isomorphic to the functor*

$$\mathcal{M}^{\star, \heartsuit}(S) = \left\{ (a, \mathcal{F}', h, \alpha) \left| \begin{array}{l} a \in \mathcal{A}^{\heartsuit}(S), \mathcal{F}' \in \overline{\text{Pic}}(\tilde{\Sigma}'_a/S), \\ h: \mathcal{F}' \xrightarrow{\sim} \sigma^*(\mathcal{F}')^{\vee} \text{ is Hermitian,} \\ \alpha: (a, \mathcal{F}', h) \xrightarrow{\sim} \iota^{\star}(a, \mathcal{F}', h) \text{ such that } \iota^{\star}\alpha \circ \alpha = \text{Id} \end{array} \right. \right\}$$

where in the above α is the data of an isomorphism $\alpha: \mathcal{F}' \rightarrow \iota^{\star}\mathcal{F}'$ which transports h to $-\iota^{\star}h$ in the Siegel case and $\iota^{\star}h$ in the unitary case. That is, α must fit into a commutative diagram

$$\begin{array}{ccc} \mathcal{F} \otimes \mathcal{F} & \xrightarrow{h} & \omega_{\tilde{\Sigma}'_a/\Sigma' \times S} \\ \alpha \otimes \alpha \downarrow & \nearrow \pm \iota^{\star}h & \\ \iota^{\star}\mathcal{F} \otimes \iota^{\star}\mathcal{F} & & \end{array}$$

with the horizontal maps being the pairings induced by h and $\pm \iota^{\star}h$ and the sign on the latter determined by the case $\star \in \{\mathbf{S}, \mathbf{U}\}$ (+1 in the case $\star = \mathbf{U}$ and -1 in the case $\star = \mathbf{S}$).

Proof. The statement mostly follows from Proposition 4.3 and Lemma 4.8. It suffices to note that, in the case $\star = \mathbf{U}$, the image of $\mathcal{M}_{r,s}^{\mathbf{U}}$ under the relative Hitchin map $h^{\mathbf{U}}$ is disjoint from $\mathcal{A}^{\heartsuit}$, so that $\mathcal{M}^{\mathbf{U}, \heartsuit} = h^{\mathbf{U}, -1}(\mathcal{A}^{\heartsuit})$. $\square$

5. SYMMETRIES OF RELATIVE HITCHIN FIBERS

It was observed in [Ngo06] that globalization of endoscopic groups is intimately tied to the groups of symmetries acting on the Hitchin fibration. For example, in type A , this manifests in an action of Picard stacks of spectral covers on the moduli of Higgs bundles. Here, we define a smooth, commutative group stack $\mathcal{P}$ which acts on the relative Hitchin moduli $\mathcal{M}^{\star}$ over $\mathcal{A}$, and give a spectral description of $\mathcal{P}$ and its action. We will study its relation to endoscopy further in §5.3.

Recall that the Hitchin base $\mathcal{A}$ is identified with the space of sections $\Gamma(\Sigma, \mathfrak{c}_D)$. For any $a \in \mathcal{A}(S)$, we consider the group scheme $J_a = a^*J$ defined as the pullback of the group scheme J from Theorem 2.9 to $\Sigma \times S$. In particular, J_a is a smooth, commutative group scheme over $\Sigma \times S$.

Definition 5.1. Let $\mathcal{P}$ denote the functor sending a k -scheme S to

$$\mathcal{P}(S) = \{(a, \mathcal{T}) \mid a \in \mathcal{A}(S), \mathcal{T} \text{ is a } J_a\text{-torsor on } \Sigma \times S\}.$$

We again denote $\mathcal{P}^{\heartsuit} = \mathcal{P}|_{\mathcal{A}^{\heartsuit}}$.

Let $\tilde{\Sigma}'$ be the spectral cover of $\Sigma' \times \mathcal{A}$ defined in §4.4. Recall from (4.10) the quotients $\bar{\Sigma}' = \tilde{\Sigma}'/\iota$ and $\bar{\Sigma} = \tilde{\Sigma}'/\sigma$. The spectral description of J (see Theorem 2.9) gives the following spectral description for $\mathcal{P}$.

Proposition 5.2. *Over $\mathcal{A}^{\heartsuit}$, we have an identification $\mathcal{P}^{\heartsuit} \simeq \text{Prym}_{/\mathcal{A}^{\heartsuit}}(\bar{\Sigma}', \bar{\Sigma})$, where $\text{Prym}_{/\mathcal{A}^{\heartsuit}}(\bar{\Sigma}', \bar{\Sigma})$ denotes the group of pairs $(\mathcal{G}', \xi)$ where $\mathcal{G}'$ is in the kernel of the norm map*

$$\text{Pic}(\bar{\Sigma}'/\mathcal{A}^{\heartsuit}) \rightarrow \text{Pic}(\bar{\Sigma}/\mathcal{A}^{\heartsuit}), \quad \mathcal{G}' \mapsto \mathcal{G}' \otimes \sigma^*\mathcal{G}'$$

and $\xi: \mathcal{G}' \simeq \sigma^*\mathcal{G}'^{\otimes -1}$ is an isomorphism realizing this.

Moreover, the action of $\mathcal{P}^{\heartsuit}$ on $\mathcal{M}^{\star, \heartsuit}$ is identified with the map

$$\mathcal{P}^{\heartsuit} \times \mathcal{M}^{\star, \heartsuit} \rightarrow \mathcal{M}^{\star, \heartsuit}$$

$$(\mathcal{G}', \mathcal{F}') \mapsto q'^*\mathcal{G}' \otimes \mathcal{F}'$$

where in the above, $q': \tilde{\Sigma}' \rightarrow \bar{\Sigma}'$ is the quotient map by ι . In this statement, we view $\mathcal{M}^{*,\heartsuit}$ in terms of the spectral description (cf. Corollary 4.9). The Hermitian form on $q'^*\mathcal{G}' \otimes \mathcal{F}'$ is induced by the corresponding forms on $\mathcal{G}'$ and $\mathcal{F}'$ by

$$q'^*\mathcal{G}' \otimes \mathcal{F}' \xrightarrow{q'^*\xi \otimes h} q'^*\sigma^*\mathcal{G}'^{\otimes -1} \otimes (\sigma^*\mathcal{F}')^\vee = \sigma^*(q'^*\mathcal{G}' \otimes \mathcal{F}')^\vee,$$

and α is induced by

$$q'^*\mathcal{G} \otimes \mathcal{F} \xrightarrow{1 \otimes \alpha} q'^*\mathcal{G} \otimes \iota^*\mathcal{F} = \iota^*(q'^*\mathcal{G} \otimes \mathcal{F}).$$

Remark 5.3. In the above result, Prym refers to the full kernel of the norm map, not just the identity component. For example, since ϑ is unramified, the Prym above has two connected components when $\bar{\Sigma}'_a$ is irreducible. In fact, it is easy to compute the component group in general; see Lemma 5.18.

Proof. By Theorem 2.9, the regular centralizer group J can be described as the kernel of the morphism

$$\mathrm{Nm}: \mathrm{Res}_{\mathbb{C}}^{(\mathfrak{s}/\iota)_{k'}}(\mathbb{G}_m) \rightarrow \mathrm{Res}_{\mathbb{C}}^{\mathfrak{s}/\iota}(\mathbb{G}_m),$$

where Nm is norm map. Since torsors over J_a for $a \in \mathcal{A}^\heartsuit(S)$ are identified with $H^1(\Sigma \times S, J_a)$, the above description of J_a describes $\mathcal{P}_a$ as the kernel of the norm map

$$\mathrm{Pic}(\bar{\Sigma}'_a) \rightarrow \mathrm{Pic}(\bar{\Sigma}_a), \quad \mathcal{G} \mapsto \mathcal{G} \otimes \sigma^*\mathcal{G}.$$

The group $\mathcal{P}_a$ is identified with the sections of $\pi_{a,*}(\mathcal{O}_{\tilde{\Sigma}_a}^\times)$ that preserve the standard Hermitian form $h: \mathcal{O}_{\tilde{\Sigma}_a} \simeq \sigma^*\mathcal{O}_{\tilde{\Sigma}_a}^\vee$ via the map $\mathcal{G} \mapsto q'^*\mathcal{G}$, and the action is given by the natural action of this unitary group on $\mathrm{Pic}(\tilde{\Sigma}'_a)$. $\square$

5.1. Open subvarieties of the relative Hitchin base. In this section, we introduce some more (open) subvarieties of the Hitchin base that will be relevant later, namely the anisotropic locus $\mathcal{A}^{\mathrm{ani}}$ (Definition 5.4) and the dagger locus $\mathcal{A}^\dagger$ (Definition 5.11), both of which will be crucial in geometric stabilization.

Definition 5.4. The *anisotropic locus* in $\mathcal{A}$, denoted by $\mathcal{A}^{\mathrm{ani}}$, is the set of all $a \in \mathcal{A}^\heartsuit(S)$ such that the double cover $\tilde{\Sigma}'_a \rightarrow \tilde{\Sigma}_a = \tilde{\Sigma}'_a/\sigma$ induces a bijection on irreducible components, i.e. the involution σ acts trivially on the set of irreducible components of the spectral cover $\tilde{\Sigma}'_a$.

We denote $\mathcal{M}^{*,\mathrm{ani}} = \mathcal{M}^* \times_{\mathcal{A}} \mathcal{A}^{\mathrm{ani}}$ and $f^{*,\mathrm{ani}}: \mathcal{M}^{*,\mathrm{ani}} \rightarrow \mathcal{A}^{\mathrm{ani}}$ the restriction of f^* to $\mathcal{M}^{*,\mathrm{ani}}$, and likewise for $\mathcal{P}^{\mathrm{ani}}$ and $g^{\mathrm{ani}}: \mathcal{P}^{\mathrm{ani}} \rightarrow \mathcal{A}^{\mathrm{ani}}$.

Remark 5.5. Note that the above definition uses the spectral curve $\tilde{\Sigma}'_a$, not $\bar{\Sigma}'_a$. The above condition implies the corresponding condition for $\bar{\Sigma}'_a$. As such, the above can be thought of as an intersection between 2 elliptic loci, one for the relative Hitchin system itself and one for unitary Hitchin system $\mathcal{N}_{2n}$ from [LN08]. While analysis of the full elliptic locus for the relative Hitchin system is important for the relative trace formula [LXZ25b], this smaller locus suffices for our local aims.

Lemma 5.6. *For $a \in \mathcal{A}(S)$, we have $a \in \mathcal{A}^{\mathrm{ani}}(S)$ if and only if $\pi_0(\mathrm{Prym}(\tilde{\Sigma}'_a, \tilde{\Sigma}_a))$ is finite. In this case, $\pi_0(\mathcal{P}_a)$ is also finite.*

Proof. It is easy to see that the Prym variety $\mathrm{Prym}(\tilde{\Sigma}'_a, \tilde{\Sigma}_a)$ has finite component group if and only if for each component of $\tilde{\Sigma}_a$, its preimage under the cover $\tilde{\Sigma}'_a \rightarrow \tilde{\Sigma}_a$ is a geometrically connected cover, which is exactly the condition of $\mathcal{A}^{\mathrm{ani}}$.

Now, suppose that $\tilde{\Sigma}'_a \rightarrow \tilde{\Sigma}_a$ induces a bijection on irreducible components. This map is ι equivariant, so after passing to the quotient by ι in diagram (4.10), we deduce that $\bar{\Sigma}'_a \rightarrow \bar{\Sigma}_a$ also induces a bijection on irreducible components. Therefore, finiteness of $\pi_0(\mathcal{P}_a)$ follows from the spectral description Proposition 5.2 using an identical argument to the first paragraph. $\square$

Lemma 5.7. *The locus $\mathcal{A}^{\text{ani}}$ is open in $\mathcal{A}$, and we have inclusions $\mathcal{A}^\diamond \subset \mathcal{A}^{\text{ani}} \subset \mathcal{A}^\heartsuit$. In particular, $\mathcal{A}^{\text{ani}}$ is nonempty.*

Proof. That this locus is open follows from [LN08, Lem 2.8.1]. For $a \in \mathcal{A}^\diamond(S)$, the spectral cover $\tilde{\Sigma}'_a \rightarrow \Sigma' \times S$ restricts to a smooth and connected, hence irreducible, curve over any geometric point s of S . Hence, $\tilde{\Sigma}'_a \rightarrow \tilde{\Sigma}_a$ induces an injection (and so also a bijection) on irreducible components, and the inclusion $\mathcal{A}^\diamond \subset \mathcal{A}^{\text{ani}}$ follows. $\square$

Proposition 5.8. *For $\star \in \{\mathbf{U}, \mathbf{S}\}$, the relative Hitchin morphism $f^{\star, \text{ani}}: \mathcal{M}^{\star, \text{ani}} \rightarrow \mathcal{A}^{\text{ani}}$ is proper and $\mathcal{M}^{\star, \text{ani}}$ is a smooth, finite type Deligne–Mumford stack.*

Proof. Since the anisotropic condition is the intersection of the anisotropic condition for $\mathbf{U}_{2n}$ Higgs bundles, $\mathcal{M}^{\star, \text{ani}}$ is the fixed point stack of $\iota^\star$ acting on $\mathcal{N}^{\text{ani}}$ by Proposition 4.3 and Corollary 4.9. Hence, smoothness follows from [Rom22, Theorem 1.2.1]. By [Rom05, Theorem 3.3], the morphism $\mathcal{M}^{\star, \text{ani}} \rightarrow \mathcal{N}^{\text{ani}}$ is representable, and by [LN08, Prop 2.8.4], $\mathcal{N}^{\text{ani}}$ is Deligne–Mumford. Hence, $\mathcal{M}^{\star, \text{ani}}$ is Deligne–Mumford.

We now note that since $\mathcal{B}$ is affine, it is separated. Therefore, as $f^{\text{ani}}: \mathcal{N}^{\text{ani}} \rightarrow \mathcal{B}^{\text{ani}}$ is proper (and so, in particular, separated), $\mathcal{N}^{\text{ani}}$ is also separated. Properness of $f^{\star, \text{ani}}$ now follows by [Rom05, Proposition 3.7] since f^{ani} is proper (see §3 and [LN08, Prop 2.8.4]). Moreover, [Rom05, Proposition 3.7] implies the map $\mathcal{M}^{\star, \text{ani}} \rightarrow \mathcal{N}^{\text{ani}}$ is proper and so in particular is finite type. Hence, $\mathcal{M}^{\star, \text{ani}}$ is locally of finite type since $\mathcal{N}^{\text{ani}}$ is. (See [Ngo10, Prop 6.1.3].) $\square$

In fact, the open subset $\mathcal{A}^{\text{ani}} \subset \mathcal{A}^\heartsuit$ can be arbitrarily large relative to the degree of D , as the following lemma makes plain.

Lemma 5.9. *The codimension of the complement $\mathcal{A}^\heartsuit - \mathcal{A}^{\text{ani}}$ in $\mathcal{A}^\heartsuit$ is at least $\deg(D)$.*

Proof. Identical to [Ngo10, Cor 6.3.6]. $\square$

Remark 5.10. Note that, while we use the above lemma in §6.8, we will need a slightly stronger version (see Lemma 9.4) for the proof of the main result.

We finally introduce the following subset of the anisotropic base.

Definition 5.11. Let $\mathcal{A}^\dagger \subset \mathcal{A}^{\text{ani}}$ denote the subvariety of $\mathcal{A}^{\text{ani}}$ whose S -points $a: \Sigma \times S \rightarrow \mathfrak{c}_D$ satisfy that image intersects transversely with the branching divisor $\mathfrak{D}_D^{\text{br}}$.

We likewise denote $\mathcal{M}^\dagger = \mathcal{M} \times_{\mathcal{A}} \mathcal{A}^\dagger$ and $h^\dagger = h|_{\mathcal{A}^\dagger}$ for the restrictions of $\mathcal{M}$ and h , respectively, to $\mathcal{A}^\dagger$.

The subset $\mathcal{A}^\dagger$ will be used in 8.2 in tandem with the support theorem. We note here the following density statement, which follows from an argument mirroring that of [Ngo10, Prop 4.7.1].

Lemma 5.12. *The set $\mathcal{A}^\dagger$ is open and dense in $\mathcal{A}$ as long as $\deg(D) > 2g$.*

5.2. Geometry over the regular locus. We consider here the geometry of the regular locus in the moduli spaces $\mathcal{M}^\star$, which is the global analogue of the regular locus of the representation $V^\star$ of $H^\star$. It turns out that this geometry is considerably richer than that of usual Higgs bundles, which will be an important technical tool in our proof of geometric stabilization.

Definition 5.13. The *regular locus* in $\mathcal{M}^\star$ is the mapping space

$$\mathcal{M}^{\star, \text{reg}} := \text{Maps}(\Sigma, [V_D^{\star, \text{reg}}/H^\star])$$

where $V^{\star, \text{reg}} = [(\mathfrak{h}^\perp)^{\text{reg}}]^{\tau_\star \sigma}$ is the open locus in $V_D^\star$ with the notion of regularity defined in §2.1.

We recall from Theorem 2.19 that

$$[\chi^{\text{reg}}]: [(\mathfrak{h}^\perp)^{\text{reg}}/\mathbf{H}] \rightarrow \mathfrak{c}$$

factors through the nonseparated scheme $(\mathfrak{h}^\perp)^{\text{reg}} // \mathbf{H}$ and the map

$$[(\mathfrak{h}^\perp)^{\text{reg}} / \mathbf{H}] \rightarrow (\mathfrak{h}^\perp)^{\text{reg}} // \mathbf{H}$$

is a gerbe banded by the regular centralizer group scheme J . We introduce the following global analogue of this intermediate space, which is studied in [HM25, §5].

Definition 5.14. The *regular base* is the mapping space $\mathcal{A}^{\star, \text{reg}} = \text{Maps}(\Sigma, V_D^{\star, \text{reg}} // \mathbf{H}^\star)$.

Note that the action of $\text{Aut}(\Sigma'/\Sigma)$ on the regular nilpotent orbits in $V_D^{\star, \text{reg}}$ is different in the unitary and Siegel cases; in particular, $\mathcal{A}^{\mathbf{U}, \text{reg}}$ and $\mathcal{A}^{\mathbf{S}, \text{reg}}$ are not isomorphic. We have the following important properties of $\mathcal{A}^{\star, \text{reg}}$.

Proposition 5.15. *The relative Hitchin morphism*

$$f^\star|_{\mathcal{M}^{\star, \text{reg}}} : \mathcal{M}^{\star, \text{reg}} \rightarrow \mathcal{A}$$

factors canonically through morphisms

$$\mathcal{M}^{\star, \text{reg}} \xrightarrow{f^{\star, \text{reg}}} \mathcal{A}^{\star, \text{reg}} \xrightarrow{\varphi^\star} \mathcal{A}$$

Moreover, $\varphi^\star$ is étale and the smooth, commutative group stack $(\varphi^\star)^*\mathcal{P}$ acts on $\mathcal{M}^{\star, \text{reg}}$ over $\mathcal{A}^{\text{reg}}$, realizing $\mathcal{M}^{\star, \text{reg}}$ as a $(\varphi^\star)^*\mathcal{P}$ -torsor.

Proof. This is a restatement of [HM25, Theorem 5.7 and Proposition 5.6]. In *loc. cit.* is assumed that k is algebraically closed, but all properties in the proposition statement can be checked after base changing to $\bar{k}$. $\square$

In fact, it is relatively straightforward to describe $\mathcal{A}^{\star, \text{reg}}$ over the open set $\mathcal{A}^\dagger \subset \mathcal{A}$ (see Definition 5.11). To prepare for this, we first prove the following lemma.

Lemma 5.16. *Consider the map*

$$(5.1) \quad V_D^{\star, \text{reg}} // H^\star \rightarrow \mathfrak{c}_D.$$

In the Siegel case ($\star = \mathbf{S}$), the restriction to the zero fiber $\Sigma = 0_D \subset \mathfrak{c}_D$ is the trivial 2-cover

$$\Sigma \sqcup \Sigma \rightarrow \Sigma.$$

In the unitary case ($\star = \mathbf{U}$), the restriction to the zero fiber is our fixed étale double cover

$$\vartheta : \Sigma' \rightarrow \Sigma.$$

Proof. Choose $\infty \in \Sigma(k)$. The action of $\pi_1(\Sigma, \infty)$ on the fiber of (5.1) over $\infty \in \Sigma = 0_D$ acts through the quotient map

$$\pi_1(\Sigma, \infty) \rightarrow \text{Aut}(\Sigma'/\Sigma),$$

where the latter acts through via the Frobenius on the set of regular nilpotent $H^\star$ -orbits in $V^\star$. This action is nontrivial on the set of regular nilpotent orbits in the unitary case and trivial in the Siegel case. Since the cover $\vartheta^{-1}(0_D) \rightarrow \Sigma$ is étale, it is determined by this action. $\square$

Consider the evaluation map $\mathcal{A}^\dagger \rightarrow \text{Sym}^{2n \deg(D)}(\Sigma)$ sending $a : \Sigma \times S \rightarrow \mathfrak{c}_D$ to $a^{-1}(\mathfrak{D}_D^{\text{br}})$. By Lemma 5.16, this evaluation map lifts to morphisms

$$\mathcal{A}^{\mathbf{S}, \text{reg}} \times_{\mathcal{A}} \mathcal{A}^\dagger \rightarrow \text{Sym}^{2n \deg(D)}(\Sigma \sqcup \Sigma) = \bigsqcup_{i=0}^{2n \deg(D)} \text{Sym}^i(\Sigma) \times \text{Sym}^{2n \deg(D)-i}(\Sigma)$$

$$\mathcal{A}^{\mathbf{U}, \text{reg}} \times_{\mathcal{A}} \mathcal{A}^\dagger \rightarrow \text{Sym}^{2n \deg(D)}(\Sigma')$$

These evaluation maps capture the entire behavior of the morphism $\varphi^\star$ over $\mathcal{A}^\dagger$.

Proposition 5.17. *The following diagrams, with horizontal arrows given by the evaluation morphisms described above and with the right vertical arrows induced by the double covers $\Sigma \sqcup \Sigma \rightarrow \Sigma$ and $\Sigma' \rightarrow \Sigma$ respectively, are Cartesian.*

$$\begin{array}{ccc} \mathcal{A}^{\mathbf{S}, \text{reg}} \times_{\mathcal{A}} \mathcal{A}^{\dagger} & \longrightarrow & \text{Sym}^{2n \deg(D)}(\Sigma \sqcup \Sigma) \\ \downarrow & & \downarrow \\ \mathcal{A}^{\dagger} & \longrightarrow & \text{Sym}^{2n \deg(D)}(\Sigma) \end{array} \quad \begin{array}{ccc} \mathcal{A}^{\mathbf{U}, \text{reg}} \times_{\mathcal{A}} \mathcal{A}^{\dagger} & \longrightarrow & \text{Sym}^{2n \deg(D)}(\Sigma') \\ \downarrow & & \downarrow \\ \mathcal{A}^{\dagger} & \longrightarrow & \text{Sym}^{2n \deg(D)}(\Sigma) \end{array}$$

Proof. Since the diagrams above are commutative, and since the left vertical arrows are étale of the same degree as the right vertical arrows, the proposition follows. $\square$

5.3. The component groups $\pi_0(\mathcal{P}_a)$. We will now consider the component groups $\pi_0(\mathcal{P}_a)$ for $a \in \mathcal{A}^{\text{ani}}(S)$ and relate them to the character lattice of the dual torus $\widehat{\mathbf{C}}$ (see Lemma 5.18 and Proposition 5.24). In addition to utility in calculations, this will be necessary for decomposing $\text{R}f_{*}^{\star, \text{ani}} \overline{\mathbb{Q}}_{\ell}$ into κ isotypic components.

The first description is a simple calculation using the spectral description in Proposition 5.2.

Lemma 5.18. *For $a \in \mathcal{A}^{\text{ani}}(S)$, let $\text{Irr}(\overline{\Sigma}_a)$ denote the set of irreducible components of $\overline{\Sigma}_a$ (which is in bijection with $\text{Irr}(\overline{\Sigma}'_a)$). Then, we have*

$$\pi_0(\mathcal{P}_a) = (\mathbb{Z}/2\mathbb{Z})^{\#\text{Irr}(\overline{\Sigma}_a)}.$$

Proof. The proof is identical to [LN08, Lem 2.8.2]. Namely, the unramified morphism $\overline{\Sigma}'_a \rightarrow \overline{\Sigma}_a$ induces an exact sequence

$$1 \rightarrow \mathbb{G}_{m, \overline{\Sigma}_a} \rightarrow \text{Res}_{\overline{\Sigma}'_a / \overline{\Sigma}_a}(\mathbb{G}_m) \xrightarrow{\beta} \text{Res}_{\overline{\Sigma}'_a / \overline{\Sigma}_a}(\mathbb{G}_m)^{\sigma^* = (\cdot)^{-1}} \rightarrow 1$$

where β is the anti-norm map $\beta(\psi) = \psi \cdot \sigma(\psi)^{-1}$. This sequence therefore induces an exact sequence on cohomology

$$\text{Pic}(\overline{\Sigma}_a) \rightarrow \text{Pic}(\overline{\Sigma}'_a) \rightarrow \mathcal{P}_a \rightarrow 1.$$

The above sequence remains exact on passing to π_0 since taking component groups is right exact, and by the condition that $a \in \mathcal{A}^{\text{ani}}(S)$, we get $\text{Pic}(\overline{\Sigma}_a) \simeq \text{Pic}(\overline{\Sigma}'_a) \simeq \mathbb{Z}^{\#\text{Irr}(\overline{\Sigma}_a)}$. The natural map

$$\pi_0(\text{Pic}(\overline{\Sigma}_a)) \rightarrow \pi_0(\text{Pic}(\overline{\Sigma}'_a))$$

is identified with the multiplication-by-2 map $\mathbb{Z}^{\#\text{Irr}(\overline{\Sigma}_a)} \xrightarrow{2} \mathbb{Z}^{\#\text{Irr}(\overline{\Sigma}_a)}$, and so the lemma follows. $\square$

We now connect the component groups of fibers with the character group of the dual torus $\widehat{\mathbf{C}}$. To do so, we must first briefly introduce a cameral-type cover. For wider applications of the geometry of Hitchin systems to relative endoscopic fundamental lemmas, we expect a greater use of similar cameral covers will be necessary.

Definition 5.19. The cameral cover $\underline{\Sigma}$ over Σ and $\underline{\Sigma}'$ over Σ' are determined by the Cartesian diagrams

$$\begin{array}{ccccc} \underline{\Sigma}' & \longrightarrow & \underline{\Sigma} & \longrightarrow & (\mathfrak{a} // \{\pm 1\}^n)_D \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{A}^{\text{ani}} \times \Sigma' & \longrightarrow & \mathcal{A}^{\text{ani}} \times \Sigma & \longrightarrow & \mathfrak{c}_D. \end{array}$$

Since $\mathfrak{s} = \mathfrak{a} // (\mathfrak{S}_{n-1} \ltimes \{\pm 1\}^n)$, any of the n natural embeddings $\mathfrak{S}_{n-1} \rightarrow \mathfrak{S}_n$ induce factorizations

$$(5.2) \quad \begin{array}{ccccc} \underline{\Sigma}' & \longrightarrow & \overline{\Sigma}' & \longrightarrow & \mathcal{A}^{\text{ani}} \times \Sigma' \\ \downarrow & & \downarrow & & \downarrow \\ \underline{\Sigma} & \longrightarrow & \overline{\Sigma} & \longrightarrow & \mathcal{A}^{\text{ani}} \times \Sigma. \end{array}$$

Now, fix $\infty \in \Sigma(k)$ which splits in Σ' and let $\infty'_1, \infty'_2 \in \Sigma'(k)$ denotes the two points lying over ∞ . We define the following modifications of the anisotropic Hitchin base.

Definition 5.20. Denote by $\mathcal{A}_{\infty}^{\text{ani}} \subset \mathcal{A}^{\text{ani}}$ the subset of points a of $\mathcal{A}^{\text{ani}}$ such that the finite, flat map

$$\underline{\Sigma}'_a \rightarrow \Sigma' \rightarrow \Sigma$$

is unramified at ∞ . Note that since ϑ is unramified, the diagram (5.2) induces a commutative diagram

$$\begin{array}{ccccc} \underline{\Sigma}'_a & \longrightarrow & \overline{\Sigma}'_a & \longrightarrow & \Sigma' \\ \downarrow & & \downarrow & & \downarrow \vartheta \\ \underline{\Sigma}_a & \longrightarrow & \overline{\Sigma}_a & \longrightarrow & \Sigma \end{array}$$

with étale maps.

Denote $\underline{\mathcal{A}}^{\text{ani}} = \underline{\mathcal{A}}_{\infty}^{\text{ani}} \subset \mathcal{A}_{\infty}^{\text{ani}} \times \Sigma'$ for the space of pairs $(a, \underline{\infty}')$ such that $a \in \mathcal{A}_{\infty}^{\text{ani}}$ and $\underline{\infty}' \in \underline{\Sigma}'_a$ has image ∞ under $\underline{\Sigma}'_a \rightarrow \Sigma$. We will drop the subscript ∞ in what follows for notational simplicity. By definition of $\mathcal{A}_{\infty}^{\text{ani}}$, the map $\underline{\mathcal{A}}^{\text{ani}} \rightarrow \mathcal{A}_{\infty}^{\text{ani}}$ is étale of degree $2(n!)$. Note that $\sigma \in \text{Aut}(\Sigma'/\Sigma)$ lifts to an action on $\underline{\mathcal{A}}^{\text{ani}}$ over $\mathcal{A}_{\infty}^{\text{ani}}$ via its action on $\underline{\Sigma}'$.

Now for $i = 1, 2$, we let $Z_i \subset \mathcal{A}_{\infty}^{\text{ani}} \times \overline{\Sigma}'$ denote the space of pairs $(a, \overline{\infty}')$ such that $\overline{\infty}' \in \overline{\Sigma}'_a$ has image $\infty'_i \in \Sigma'(k)$. Note that $Z_i \rightarrow \mathcal{A}_{\infty}^{\text{ani}}$ is étale of degree n and σ induces an isomorphism $Z_1 \rightarrow Z_2$.

We have a concrete realization of $\underline{\mathcal{A}}^{\text{ani}}$ as follows.

Lemma 5.21. *There is an isomorphism of spaces over $\mathcal{A}_{\infty}^{\text{ani}}$*

$$\underline{\mathcal{A}}^{\text{ani}} \simeq (Z_1^n \setminus \Delta_1) \sqcup (Z_2^n \setminus \Delta_2),$$

where Δ_i denotes the union of all pairwise diagonals in the n -fold product Z_i^n . This isomorphism intertwines the action of σ on $\underline{\mathcal{A}}^{\text{ani}}$ with the induced isomorphism $(Z_1^n \setminus \Delta_1) \rightarrow (Z_2^n \setminus \Delta_2)$.

Proof. The n distinct factorizations of (5.2) induce a map

$$(5.3) \quad \mathfrak{a} // \{\pm 1\}^n \rightarrow \prod_{i=1}^n \mathfrak{a} // (\mathfrak{S}_{n-1} \ltimes \{\pm 1\}^n)$$

sending

$$(x_1, \dots, x_n) \mapsto \prod_i (x_i, [x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n]),$$

where square brackets denote unordered tuples. The resulting morphism $\underline{\mathcal{A}}^{\text{ani}} \rightarrow Z_1^n \sqcup Z_2^n$ obtained by composing the evaluation map is defined over $\mathcal{A}_{\infty}^{\text{ani}}$ has image in the complement of Δ since we are assuming the underlying point a is in $\mathcal{A}_{\infty}^{\text{ani}}$. Moreover, both sides are finite and étale over $\mathcal{A}_{\infty}^{\text{ani}}$ of degree $2(n!)$. Since (5.3) is an injection, the map $\underline{\mathcal{A}}^{\text{ani}} \rightarrow Z_1^n \sqcup Z_2^n$ is injective and hence an isomorphism. $\square$

Denote $Z = Z_1 \sqcup Z_2$ for the space of pairs $(a, \overline{\infty}') \in \mathcal{A}_{\infty}^{\text{ani}} \times_{\mathcal{A}} \overline{\Sigma}'$ such that $\overline{\infty}' \mapsto \infty$. There is a natural action of σ on Z from the isomorphism $\sigma: Z_1 \rightarrow Z_2$.

Corollary 5.22. *The base change*

$$Z \times_{\mathcal{A}_{\infty}^{\text{ani}}} \underline{\mathcal{A}}^{\text{ani}} \rightarrow \underline{\mathcal{A}}^{\text{ani}}$$

is isomorphic to the trivial cover

$$\prod_{i=1}^{2n} \underline{\mathcal{A}}^{\text{ani}} \rightarrow \underline{\mathcal{A}}^{\text{ani}}.$$

with σ acting freely on the components. In particular, there are n distinct sections

$$y_1, \dots, y_n: \underline{\mathcal{A}}^{\text{ani}} \rightarrow \overline{\Sigma}' \times_{\mathcal{A}_{\infty}^{\text{ani}}} \underline{\mathcal{A}}^{\text{ani}}$$

so that $y_1, \dots, y_n, \sigma(y_1), \dots, \sigma(y_n)$ index all components of $Z \times_{\mathcal{A}_{\infty}^{\text{ani}}} \underline{\mathcal{A}}^{\text{ani}}$.

Proof. It suffices to show that for $i = 1, 2$, the cover

$$Z_i \times_{\mathcal{A}_{\infty}^{\text{ani}}} \underline{\mathcal{A}}^{\text{ani}} \rightarrow \underline{\mathcal{A}}^{\text{ani}}$$

is trivial. Moreover, since $Z_1 \simeq Z_2$, by Lemma 5.21 it suffices to show

$$Z_i \times_{\mathcal{A}_{\infty}^{\text{ani}}} (Z_i^n \setminus \Delta) \rightarrow (Z_i^n \setminus \Delta)$$

is trivial. A point in the left hand side is a tuple $(a, \overline{\omega}', (\overline{\omega}'_1, \dots, \overline{\omega}'_n))$ where $\overline{\omega}', \overline{\omega}'_i \in \overline{\Sigma}'_a$ all have image ω'_i and $\overline{\omega}'_j \neq \overline{\omega}'_\ell$ for all $j \neq \ell$. In particular, since the map $\overline{\Sigma}'_a \rightarrow \Sigma'$ is of degree n , we must have $\overline{\omega}' = \overline{\omega}'_j$ for some j . Therefore, we have an explicit isomorphism

$$\bigsqcup_{j=1}^n (Z_i^n \setminus \Delta) \rightarrow Z_i \times_{\mathcal{A}_{\infty}^{\text{ani}}} (Z_i^n \setminus \Delta)$$

$$(\overline{\omega}'_1, \dots, \overline{\omega}'_n)_j \mapsto (a, \overline{\omega}'_j, (\overline{\omega}'_1, \dots, \overline{\omega}'_n))$$

where the subscript j denotes the j -th copy of $(Z_i^n \setminus \Delta)$. □

Recall that the canonical torus $\mathbf{T}$ of $G = \text{U}_{2n}$ fits into an exact sequence

$$1 \rightarrow \mathbf{C} \rightarrow \mathbf{T} \rightarrow \mathbf{A} \rightarrow 1,$$

where $\mathbf{C}$ is the fixed points of a canonical involution on $\mathbf{T}$. Let Ω be the set of weights of the defining representation of GL_{2n} . That is, $\Omega = \{\varepsilon_1, \dots, \varepsilon_{2n}\}$ where ε_i is the i -th coordinate function on the lattice $\mathbb{X}^*(\mathbf{T})$. The Frobenius σ acts on Ω and we can choose an ordering of the ε_i so that $\varepsilon_{i+n} = \sigma(\varepsilon_i)$ for all $i = 1, \dots, n$.

For $\omega \in \Omega$, we have an induced function $\omega: \underline{\mathcal{A}}^{\text{ani}} \rightarrow Z$ which, under the identification $\underline{\mathcal{A}}^{\text{ani}} \simeq (Z_1^n \setminus \Delta_1) \sqcup (Z_2^n \setminus \Delta_2)$ of Lemma 5.21, is expressed by

$$\varepsilon_i(\overline{\omega}'_1, \dots, \overline{\omega}'_n) = \overline{\omega}'_i$$

$$\varepsilon_{i+n}(\overline{\omega}'_1, \dots, \overline{\omega}'_n) = \sigma(\overline{\omega}'_i)$$

for $i = 1, \dots, n$.

Denote $Z = Z_1 \sqcup Z_2$ for the space of pairs $(a, \overline{\omega}') \in \mathcal{A}_{\infty}^{\text{ani}} \times_{\mathcal{A}} \overline{\Sigma}'$ such that $\overline{\omega}' \mapsto \infty$. There is a natural action of σ on Z from the isomorphism $\sigma: Z_1 \rightarrow Z_2$.

Corollary 5.23. *The base change*

$$Z \times_{\mathcal{A}_{\infty}^{\text{ani}}} \underline{\mathcal{A}}^{\text{ani}} \rightarrow \underline{\mathcal{A}}^{\text{ani}}$$

is canonically isomorphic to the trivial cover

$$\prod_{\omega \in \Omega} \underline{\mathcal{A}}^{\text{ani}} \rightarrow \underline{\mathcal{A}}^{\text{ani}},$$

where Ω is the set of weights of the defining representation of U_{2n} , with σ acting freely on the components via the conjugation action on the weight lattice of U_{2n} . In particular, for every weight $\omega \in \Omega$, we have a section

$$y_\omega: \underline{\mathcal{A}}^{\text{ani}} \rightarrow \overline{\Sigma}' \times_{\mathcal{A}_\infty^{\text{ani}}} \underline{\mathcal{A}}^{\text{ani}}.$$

Proof. By Lemma 5.21, a point in the left hand side is a tuple $\underline{a} = (a, \overline{\omega}', \underline{\omega}')$ where $\overline{\omega}'$ has image ∞ ; $\underline{\omega}' = (\overline{\omega}'_1, \dots, \overline{\omega}'_n)$ for $\overline{\omega}'_i \in \overline{\Sigma}'_a$; there exists a choice of $i \in \{1, 2\}$ so that each $\overline{\omega}'_i$ has image $\overline{\omega}'_i$; and $\overline{\omega}'_j \neq \overline{\omega}'_\ell$ for all $j \neq \ell$. In particular, since the map $\mathfrak{a} // \{\pm 1\}^n \rightarrow \mathfrak{a} // (\mathfrak{S}_{n-1} \ltimes \{\pm 1\}^n)$ is of degree $(n-1)!$, there must exist some index j such that $\overline{\omega}'$ and $\overline{\omega}'_j$ have the same image in $\mathfrak{s}_D$. In other words, such that either $\overline{\omega}' = \overline{\omega}'_j$ or $\overline{\omega}' = \sigma(\overline{\omega}'_j)$; and so there exists a unique $\omega_{\underline{a}} \in \Omega$ such that $\omega_{\underline{a}}(\underline{\omega}') = \overline{\omega}'$. Therefore, we have an explicit isomorphism

$$Z \times_{\mathcal{A}_\infty^{\text{ani}}} \underline{\mathcal{A}}^{\text{ani}} \rightarrow \bigsqcup_{\omega \in \Omega} \underline{\mathcal{A}}^{\text{ani}}$$

$$\underline{a} = (a, \overline{\omega}', \underline{\omega}') \mapsto (\underline{\omega}')_{\omega_{\underline{a}}}$$

where the subscript $\omega_{\underline{a}}$ denotes the $\omega_{\underline{a}}$ -th copy of $\underline{\mathcal{A}}^{\text{ani}}$. The inverse of the above map is given by $(\underline{\omega}')_\omega \mapsto (a, \omega(\underline{\omega}'), \underline{\omega}')$ $\square$

We now apply this to study the components of $\mathcal{P}$. Denote by $\underline{p}: \underline{\mathcal{A}}^{\text{ani}} \rightarrow \mathcal{A}_\infty^{\text{ani}}$, and $\underline{\mathcal{P}} = \underline{p}^*(\mathcal{P}|_{\mathcal{A}_\infty^{\text{ani}}})$. By [Ngo06, Proposition 6.2], there is a constructible sheaf $\pi_0(\underline{\mathcal{P}}/\underline{\mathcal{A}}^{\text{ani}})$ on $\underline{\mathcal{A}}^{\text{ani}}$ whose fiber over any a in $\underline{\mathcal{A}}^{\text{ani}}$ is the component group of the fiber $\mathcal{P}_a$.

Corollary 5.23 gives a canonical map

$$\mathbb{X}_*(\mathbf{T}) \times \underline{\mathcal{A}}^{\text{ani}} \rightarrow \text{Pic}(\overline{\Sigma}' \times_{\mathcal{A}_\infty^{\text{ani}}} \underline{\mathcal{A}}^{\text{ani}} / \underline{\mathcal{A}}^{\text{ani}})$$

$$(\lambda, a) \mapsto \mathcal{O}_{\overline{\Sigma}'_a} \left(\sum_{\omega \in \Omega} \langle \omega, \lambda \rangle y_\omega \right)$$

When $\lambda \in \mathbb{X}_*(\mathbf{C})$, we have $\langle \omega, \lambda \rangle = -\langle \sigma(\omega), \lambda \rangle$ for $\omega \in \Omega$, where λ is viewed in $\mathbb{X}_*(\mathbf{T})$. The line bundle above then becomes

$$\mathcal{O}_{\overline{\Sigma}'_a} \left(\sum_{\omega \in \Omega} \langle \omega, \lambda \rangle y_\omega \right) = \mathcal{O}_{\overline{\Sigma}'_a} \left(\sum_{i=1}^n \langle \varepsilon_i, \lambda \rangle (y_i - \sigma(y_i)) \right).$$

In particular, this carries a unique Hermitian structure, and so the image lies in the Prym variety $\underline{\mathcal{P}}$, and we deduce a morphism

$$\mathbb{X}_*(\mathbf{C}) \times \underline{\mathcal{A}}^{\text{ani}} \rightarrow \underline{\mathcal{P}}$$

over $\underline{\mathcal{A}}^{\text{ani}}$. This induces a surjection on π_0 by Lemma 5.18, and hence we obtain the following.

Proposition 5.24. *There is a canonical surjective morphism of sheaves on $\underline{\mathcal{A}}^{\text{ani}}$*

$$(5.4) \quad \mathbb{X}^*(\widehat{\mathbf{C}}) \times \underline{\mathcal{A}}^{\text{ani}} \rightarrow \pi_0(\underline{\mathcal{P}}/\underline{\mathcal{A}}^{\text{ani}})$$

which is given explicitly by identifying $\mathbb{X}^*(\widehat{\mathbf{C}}) \simeq \mathbb{Z}^n$ and sending $((d_1, \dots, d_n), a)$ to the component of the bundle

$$\mathcal{O} \left(\sum_{i=1}^n d_i (y_i - \sigma(y_i)) \right) \in \underline{\mathcal{P}}_a$$

where $y_1, \dots, y_n$ are the sections in Corollary 5.23 for the elements $\varepsilon_1, \dots, \varepsilon_n \in \Omega$.

Proof. Immediate from Corollary 5.23 and the discussion above. $\square$

We denote $\underline{g}^{\text{ani}}: \underline{\mathcal{P}}^{\text{ani}} \rightarrow \underline{\mathcal{A}}^{\text{ani}}$ and $\underline{f}^{\star, \text{ani}}: \underline{\mathcal{M}}^{\star, \text{ani}} \rightarrow \underline{\mathcal{A}}^{\text{ani}}$ for the base change of the respective structure maps g and f to $\underline{\mathcal{A}}^{\text{ani}}$. Note that $\underline{\mathcal{P}}$ acts on $\text{R}\underline{g}_!^{\text{ani}}\overline{\mathbb{Q}}_\ell$ and $\text{R}\underline{f}_*^{\text{ani}}\overline{\mathbb{Q}}_\ell$ through $\pi_0(\underline{\mathcal{P}}/\underline{\mathcal{A}}^{\text{ani}})$. Hence, the map (5.4) defines κ -decompositions

$$(5.5) \quad \text{R}\underline{g}_!^{\text{ani}}\overline{\mathbb{Q}}_\ell = \bigoplus_{\kappa} (\text{R}\underline{g}_!^{\text{ani}}\overline{\mathbb{Q}}_\ell)_{\kappa}, \quad \text{R}\underline{f}_*^{\text{ani}}\overline{\mathbb{Q}}_\ell = \bigoplus_{\kappa} (\text{R}\underline{f}_*^{\star, \text{ani}}\overline{\mathbb{Q}}_\ell)_{\kappa},$$

where the each direct sum is finite-indexed over the same set of $\kappa \in \widehat{\mathbf{C}}$, and the subscript κ indicates the subsheaf on which the character group $\mathbb{X}^*(\widehat{\mathbf{C}})$ acts through the evaluation map at κ .

Corollary 5.25. *For $\kappa \in \widehat{\mathbf{C}}$, we have $(\text{R}\underline{f}_*^{\star, \text{ani}}\overline{\mathbb{Q}}_\ell)_{\kappa} \neq 0$ if and only if κ is 2-torsion in $\widehat{\mathbf{C}}$. In particular, $(\text{R}\underline{f}_*^{\star, \text{ani}}\overline{\mathbb{Q}}_\ell)_{\kappa} \neq 0$ if and only if κ extends to an elliptic endoscopic datum $\mathfrak{e}$.*

Proof. By Proposition 5.24, for each $a \in \mathcal{A}^{\text{ani}}(S)$ the map $\mathbb{X}^*(\widehat{\mathbf{C}}) \rightarrow \pi_0(\mathcal{P}_a)$ factors through the quotient $\mathbb{X}^*(\widehat{\mathbf{C}})/2\mathbb{X}^*(\widehat{\mathbf{C}})$. In particular, if $(\text{R}\underline{f}_*^{\star, \text{ani}}\overline{\mathbb{Q}}_\ell)_{\kappa}$ has nonzero stalk at a , κ must be order 2 in $\widehat{\mathbf{C}}$.

Conversely, we note that for any distinct nonzero sections $s_i \in H^0(\Sigma, L)$, we can consider the point $a \in \mathcal{A}^{\text{ani}}(k)$ associated to the characteristic polynomial

$$\prod_{i=1}^n (\lambda^2 - s_i).$$

That is, a_{2i} is the coefficient of $\lambda^{2(n-i)}$ in the above polynomial, which is a section in $H^0(\Sigma, L^{2i})$. The corresponding spectral cover $\overline{\Sigma}_a$ has n components, each of which is isomorphic to Σ , and since $\overline{\Sigma}'_a = \overline{\Sigma}_a \times_{\Sigma} \Sigma'$, the same is true for $\overline{\Sigma}'_a$. We conclude that $a \in \mathcal{A}^{\text{ani}}$, and that $\pi_0(\mathcal{P}_a) = \mathbb{Z}^n$ is maximal. Therefore, the stalk of $(\text{R}\underline{f}_*^{\star, \text{ani}}\overline{\mathbb{Q}}_\ell)_{\kappa}$ is nontrivial at a for every 2 torsion element κ and the converse holds. $\square$

5.4. The Endoscopic Case. Let $\mathfrak{e} = (\kappa, \rho)$ be a relative endoscopic datum for $X^{\mathbf{S}}$ over Σ . From §2.3, we have the associated endoscopic space $X^{\mathfrak{e}}$, defined over Σ , which are products of Siegel and unitary cases of complementary ranks. We may define all the same structures using the bundle of $H^{\mathfrak{e}}$ -representations $V_D^{\mathfrak{e}}$. In particular, we have a moduli of $X^{\mathfrak{e}}$ -Higgs bundles, $\mathcal{M}^{\mathfrak{e}}$, which comes equipped with a Hitchin fibration

$$f^{\mathfrak{e}}: \mathcal{M}^{\mathfrak{e}} \rightarrow \mathcal{A}_{\mathfrak{e}},$$

which itself is a product of Siegel and unitary cases. The transfer morphism $\nu_{\mathfrak{e}}: \mathfrak{c}_{\mathfrak{e}} \rightarrow \mathfrak{c}$ induces a morphism of Hitchin bases, which we also denote

$$\nu_{\mathfrak{e}}: \mathcal{A}_{\mathfrak{e}} \rightarrow \mathcal{A}.$$

Recall from Proposition 2.14 that $\mathfrak{s}_{\mathfrak{e}}/\iota_{\mathfrak{e}} \rightarrow \mathfrak{s}/\iota \times_{\mathfrak{c}} \mathfrak{c}_{\mathfrak{e}}$ is a partial normalization which separates two components of $\mathfrak{s}/\iota \times_{\mathfrak{c}} \mathfrak{c}_{\mathfrak{e}}$ canonically. In particular, for $\nu_{\mathfrak{e}}(a_{\mathfrak{e}}) = a$ there is a map $i_{\mathfrak{e}}: \overline{\Sigma}'_{a_{\mathfrak{e}}} \rightarrow \overline{\Sigma}'_a$ which realizes $\overline{\Sigma}'_{a_{\mathfrak{e}}}$ as a union of components of $\overline{\Sigma}'_a$. We describe this explicitly as follows: Suppose that $\mathfrak{e} = (\kappa, \rho)$ is an endoscopic datum associated to a 2-torsion element $\kappa \in \widehat{\mathbf{C}}$. Let $S_{\kappa} \subset \{1, \dots, n\}$ be the subset of indices for which $\kappa_i = 1$ and set $\#S_{\kappa} = r$. For $a \in \mathcal{A}^{\heartsuit}(k)$, we have an associated characteristic polynomial $Q_a(\lambda^2)$, where $Q(\lambda)$ is a polynomial of degree n with i -th coefficient in $H^0(\Sigma, L^{2i})$. A lift $a_{\mathfrak{e}} \in \mathcal{A}_{\mathfrak{e}}^{\heartsuit}(k)$ is the data of a factorization $Q_a = Q_1 Q_2$, where $\deg(Q_1) = r$ and $\deg(Q_2) = n - r$. Furthermore, if $\overline{\Sigma}_{a_{\mathfrak{e}}, i}$ denotes the vanishing locus of $Q_i(\lambda)$ in the total space of L , then the image of the sections $\{y_j \mid j \in S_{\kappa}\}$ of Corollary 5.23 are in $\overline{\Sigma}_{a_{\mathfrak{e}}, 1}$ and $\{y_j \mid j \notin S_{\kappa}\}$ are in $\overline{\Sigma}_{a_{\mathfrak{e}}, 2}$.

Therefore, for any such $a_{\mathfrak{e}} \in \mathcal{A}_{\mathfrak{e}}^{\heartsuit}$, the spectral curve $\overline{\Sigma}_{a_{\mathfrak{e}}}$ is a disjoint union $\overline{\Sigma}_{a_{\mathfrak{e}}} = \overline{\Sigma}_{a_{\mathfrak{e}}, 1} \sqcup \overline{\Sigma}_{a_{\mathfrak{e}}, 2}$ while $\overline{\Sigma}_a = \overline{\Sigma}_{a_{\mathfrak{e}}, 1} \cup \overline{\Sigma}_{a_{\mathfrak{e}}, 2}$ is a (not necessarily disjoint) union. Each $\overline{\Sigma}_{a_{\mathfrak{e}}, i}$ is a union of components of $\overline{\Sigma}_a$ and the map $\overline{\Sigma}_{a_{\mathfrak{e}}} \rightarrow \overline{\Sigma}_a$ is the inclusion map of $\overline{\Sigma}_{a_{\mathfrak{e}}, i}$ into $\overline{\Sigma}_a$. By the anisotropic condition, this can all be lifted to the double covers $i_{\mathfrak{e}}: \overline{\Sigma}'_{a_{\mathfrak{e}}} = \overline{\Sigma}'_{a_{\mathfrak{e}}, 1} \sqcup \overline{\Sigma}'_{a_{\mathfrak{e}}, 2} \rightarrow \overline{\Sigma}'_a$.

By Lemma 5.21, the map $\nu_{\mathfrak{e}}$ lifts to the cover

$$(5.6) \quad \underline{\nu}_{\mathfrak{e}}: \underline{\mathcal{A}}_{\mathfrak{e}} \rightarrow \underline{\mathcal{A}}.$$

More precisely, if $\kappa = (1_r, -1_{n-r})$ the corresponding map is

$$\underline{\nu}_{\mathfrak{e}}[(a_{\mathfrak{e}}, (\overline{\omega}'_1, \dots, \overline{\omega}'_r), (\overline{\omega}'_{r+1}, \dots, \overline{\omega}'_n))] = (\nu_{\mathfrak{e}}(a_{\mathfrak{e}}), (i_{\mathfrak{e}}(\overline{\omega}'_1), \dots, i_{\mathfrak{e}}(\overline{\omega}'_n))).$$

(When κ is a permutation of $(1_r, -1_s)$, we interlace the orders as specified by the order of the signs on κ .)

Lemma 5.26. *The map (5.6) is a closed embedding over the heart locus $\underline{\mathcal{A}}^{\heartsuit}$.*

Proof. Since the map $\mathfrak{c}_{\mathfrak{e}} \rightarrow \mathfrak{c}$ is induced by the inclusion of Weyl groups $W_{X^{\mathfrak{e}}} \subset W_X$, it follows from [Ngo06, Proposition 7.2] that (5.6) is finite and unramified. Therefore, it is enough to check the map is injective on geometric points. Suppose that $(a, (\overline{\omega}'_1, \dots, \overline{\omega}'_n)) \in \underline{\mathcal{A}}^{\heartsuit}(\overline{k})$ is in the image of $\nu_{\mathfrak{e}}$. Then, the data of κ and the ordering $(\overline{\omega}'_1, \dots, \overline{\omega}'_n)$ determine a decomposition $\overline{\Sigma}'_a = \overline{\Sigma}'_{a,1} \cup \overline{\Sigma}'_{a,2}$, which forces a unique preimage for $(a, (\overline{\omega}'_1, \dots, \overline{\omega}'_n))$ in $\underline{\mathcal{A}}_{\mathfrak{e}}(\overline{k})$. $\square$

For any $\kappa \in \widehat{\mathbf{C}}$, we let $\underline{\mathcal{A}}_{\kappa}$ denote the support of $(R\underline{g}_!^{\text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}$, which is a closed subvariety of $\underline{\mathcal{A}}^{\text{ani}}$.

Proposition 5.27. *For any $\kappa \in \widehat{\mathbf{C}}$, we have a decomposition of top degree cohomology*

$$R^{2d} \underline{g}_!^{\text{ani}} \overline{\mathbb{Q}}_{\ell} = \bigoplus_{\mathfrak{e}} \underline{\nu}_{\mathfrak{e},*}(\overline{\mathbb{Q}}_{\ell})$$

where the sum is over all endoscopic data $\mathfrak{e} = (\kappa, \rho)$. Moreover, the sheaf $\pi_0(\underline{\mathcal{P}}/\underline{\mathcal{A}}^{\text{ani}})$ is locally constant on an open subset of $\underline{\mathcal{A}}_{\kappa}$, and $\underline{\mathcal{A}}_{\kappa} = \underline{\nu}_{\mathfrak{e}}(\underline{\mathcal{A}}_{\mathfrak{e}})$, where $\mathfrak{e} = (\kappa, \rho)$ is the (unique) endoscopic datum containing κ .

Proof. The support of $(R\underline{g}_!^{\text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}$ can be computed by Lemma 5.18 and Proposition 5.24. Namely, the support is the locus in $\underline{\mathcal{A}}^{\text{ani}}$ for which the kernel of the evaluation map at κ ,

$$(5.7) \quad \mathbb{X}_*(\mathbf{C}) \rightarrow \overline{\mathbb{Q}}_{\ell}^{\times}$$

contains the kernel of (5.4). Corollary 5.25, the only κ with nontrivial $(\underline{g}_!^{\text{ani}} \overline{\mathbb{Q}}_{\ell, \mathcal{P}})_{\kappa}$ are those for which $\kappa = (\kappa_1, \dots, \kappa_n) \in \widehat{\mathbf{C}}$ is 2-torsion; thus, $\kappa_i \in \{\pm 1\}$ for all i . Let $\kappa \in \widehat{\mathbf{C}}[2]$ and let $S_{\kappa} \subset \{1, \dots, n\}$ denote the set of all indices i for which $\kappa_i = -1$. The kernel of the map (5.7) is given explicitly by

$$\left\{ (x_1, \dots, x_n) \in \mathbb{X}_*(\widehat{\mathbf{C}}) \mid \sum_{i \in S_{\kappa}} x_i \equiv 0 \pmod{2} \right\}$$

The map (5.4) is also quite explicit: Fix $\varepsilon_1, \dots, \varepsilon_n \in \Omega$ so that $\Omega = \{\varepsilon_1, \dots, \varepsilon_n, \sigma(\varepsilon_1), \dots, \sigma(\varepsilon_n)\}$. At $a \in \underline{\mathcal{A}}^{\text{ani}}(k)$, the spectral curve $\overline{\Sigma}'_a$ decomposes into irreducible components C_i . We put $S_i \subset \{1, \dots, n\}$ for the set of indices such that the image $y_j(a)$ of the section y_j of Corollary 5.23 lies in C_i . We have a partition

$$\{1, \dots, n\} = S_1 \sqcup \dots \sqcup S_{\ell}.$$

Then the map sends

$$(x_1, \dots, x_n) \mapsto \left(\sum_{j \in C_i} y_j(a) \right)_i \in (\mathbb{Z}/2\mathbb{Z})^{\#\text{Irr}(\overline{\Sigma}_a)}$$

over a . In particular, we see that the containment of kernels happens exactly when S_{κ} is one of the S_i ; that is, the sections $y_j(a)$ for $j \in S_{\kappa}$ determine a component of $\overline{\Sigma}_a$. It follows immediately that $\underline{\mathcal{A}}_{\kappa}$ is the preimage of $\underline{\nu}_{\mathfrak{e}}(\underline{\mathcal{A}}_{\mathfrak{e}})$. Moreover, the above gives a bijection between the set of irreducible components of $\overline{\Sigma}_a$ over any point $a \in \underline{\mathcal{A}}^{\text{ani}}$ and the set of κ with $a \in \underline{\mathcal{A}}_{\kappa}$, so that the natural map

$$\bigoplus_{\mathfrak{e}} \underline{\nu}_{\mathfrak{e},*}(\overline{\mathbb{Q}}_{\ell}) \rightarrow R^{2d} \underline{g}_!^{\text{ani}} \overline{\mathbb{Q}}_{\ell, \mathcal{P}}$$

is an isomorphism. $\square$

5.5. The δ -stratification on $\mathcal{A}^{\text{ani}}$. The group stack $\mathcal{P}$ is a gerbe over a commutative group scheme P . By Chevalley’s structure theorem, for any $a \in \mathcal{A}^{\text{ani}}$, we have an exact sequence

$$0 \rightarrow R_a \rightarrow P_a \rightarrow A_a \rightarrow 0,$$

where R_a is the unique maximal connected affine subgroup of P_a and A_a is an abelian variety.

Definition 5.28. The (global) δ -invariant at $a \in \mathcal{A}^{\text{ani}}$ is the dimension of the affine part of P_a , i.e. $\delta(a) = \dim R_a$.

We denote by $\mathcal{A}_\delta$ the locally closed subset of $\mathcal{A}^{\text{ani}}$ consisting of a in $\mathcal{A}^{\text{ani}}$ with $\delta(a) = \delta$.

Lemma 5.29. *Let $a \in \mathcal{A}^{\text{ani}}(\bar{k})$, and let $\delta_{n,a}$ denote the dimension of the affine part of the Jacobian $\text{Jac}(\bar{\Sigma}_a)$. Then, we have $\delta(a) = \delta_{n,a}$.*

Proof. Let $\delta'_{n,a}$ denote the dimension of the affine part of $\text{Jac}(\bar{\Sigma}'_a)$. Since $\bar{\Sigma}'_a \rightarrow \bar{\Sigma}_a$ is an unramified double cover, we have $\delta'_{n,a} = 2\delta_{n,a}$. Hence, the result follows from the short exact sequence

$$1 \rightarrow \text{Prym}(\bar{\Sigma}'_a, \bar{\Sigma}_a) \rightarrow \text{Jac}(\bar{\Sigma}'_a) \rightarrow \text{Jac}(\bar{\Sigma}_a) \rightarrow 1$$

by Proposition 5.2. $\square$

We may form the relative Jacobian

$$\text{Jac}(\bar{\Sigma}/\mathcal{A}) \rightarrow \mathcal{A},$$

which over the locus $\mathcal{A}^\heartsuit$ is a relative Jacobian of a family of curves with reduced fibers. As the dimension of the affine part of $\text{Jac}(\bar{\Sigma}_a)$, $\delta_{n,a}$ is an upper semi-continuous function, which agrees with $\delta_a = \dim \mathcal{P}_a$ by Lemma 5.29. By [Ngo10, §4.4.1] and [Ngo10, Prop 5.7.2], we obtain the following as a corollary with the necessary constraint on characteristic of k .

Corollary 5.30. *Assume that $\text{char}(k) > n$. For every integer $\delta_0 \geq 0$, there exists some integer N_0 such that if $\deg(D) \geq N_0$, then for every $\delta \leq \delta_0$, we have*

$$\text{codim}_{\mathcal{A}^{\text{ani}}}(\mathcal{A}_\delta) \geq \delta.$$

6. RELATIVE AFFINE SPRINGER FIBERS

In this section, we prove basic results about certain relative affine Springer fibers (RASFs) which arise in the local study of endoscopic orbital integrals for Friedberg–Jacquet varieties. In particular, we give moduli and lattice descriptions of RASFs; we study the component groups of RASFs for split elements (Proposition 6.20); and we relate the RASFs to relative Hitchin fibers via a product formula (Proposition 6.24). We use the product formula together with a global argument to obtain a dimension formula for RASFs (Theorem 6.28).

6.1. Lattice definition of RASFs. Similar to the GL_n case of usual affine Springer fibers (cf. [Ngo10, §3.6] for example), relative affine Springer fibers admit a definition in terms of moduli spaces of lattices. We choose to define these local spaces first in terms of lattices because this definition is at once elementary and makes clear questions of representability. We then give a definition that is likely to generalize to other spherical varieties in §6.2, relying on an important lifting property (Proposition 6.6) which we prove at the end of the present section.

We begin with a definition of lattices.

Definition 6.1. Let F be a non-archimedean local field and let $\mathcal{O} \subset F$ be the corresponding ring of integers. Let R denote an $\mathcal{O}$ -algebra which is free as an $\mathcal{O}$ -module. An R -lattice in $R \otimes_{\mathcal{O}} F$ is an R -submodule $\Lambda \subset R \otimes_{\mathcal{O}} F$ which is free as an $\mathcal{O}$ -module and for which $\Lambda \otimes_{\mathcal{O}} F = R \otimes_{\mathcal{O}} F$.

Now we specialize to our settings. Throughout this section, we let $F = k((\varpi_v))$ and we let $F' = F'_v$ be a fixed unramified quadratic extension of F with $\sigma \in \text{Gal}(F'/F)$ the nontrivial element in the Galois group. In particular, $F' = k'((\varpi_v))$, where k' is the unique quadratic extension k' of k . (For simplicity, we will drop the subscripts v throughout this section.) Let $\mathfrak{c}^\heartsuit(\mathcal{O}) := \mathfrak{c}(F)^{\text{rs}} \cap \mathfrak{c}(\mathcal{O}_F)$ denote all elements $a \in \mathfrak{c}(\mathcal{O}_F)$ which are regular, semisimple over F , i.e. for which the centralizer J_a is a torus. For $a = (a_{2i})_{i=1}^n \in \mathfrak{c}^\heartsuit(\mathcal{O}_F)$, let $P_a(t) = x^{2n} + a_2x^{2n-2} + \cdots + a_{2n} \in \mathcal{O}_F[t]$ denote the corresponding characteristic polynomial. Let

$$(6.1) \quad R_a(F') = F'[x]/(P_a(x)) \text{ and } R_a(\mathcal{O}_{F'}) = \mathcal{O}_{F'}[x]/(P_a(x)).$$

There are two natural involutions on these algebras: We let $\iota_R(f(x)) = f(-x)$ and

$$\sigma_R \left(\sum_{i=0}^{2n-1} \alpha_i x^i \right) = \sum_{i=0}^{2n-1} \sigma(\alpha_i) \cdot (-x)^i,$$

We set $R_a(F) = R_a(F')^{\sigma_R}$ and $R_a(\mathcal{O}_F) = R_a(\mathcal{O}_{F'})^{\sigma_R}$. The involution ι_R descends to an involution on R_a which we also denote by ι_R .

A non-unitary version of a relative affine Springer fiber could be defined as the functor classifying $R_a(\mathcal{O}_F)$ lattices Λ in $R_a(F)$ satisfying $\iota_R(\Lambda) = \Lambda$. To describe our unitary versions $\mathcal{M}^\star(a)$, we need to consider the additional structure of a Hermitian form on $R_a(F')$ whose compatibility with ι_R is dependent on whether we are in the Siegel or unitary case.

We begin with $\mathcal{M}^\mathbf{S}(a)$. Consider the natural pairing $h^\mathbf{S}: R_a(F') \times R_a(F') \rightarrow F'$ by

$$h^\mathbf{S}(f, g) = \text{tr}_{R_a(F')/F'} \left(\frac{f \cdot \sigma_R(g)}{P'_a(x)} \right)$$

The pairing $h^\mathbf{S}$ is clearly Hermitian and is nondegenerate by Euler's formula. We call an $\mathcal{O}_{F'}$ -lattice Λ in $R_a(F')$ $h^\mathbf{S}$ -self-dual if $\Lambda = \Lambda^{\mathbf{S}-\vee}$, where $\Lambda^{\mathbf{S}-\vee}$ is the dual lattice with respect to $h^\mathbf{S}$, i.e.

$$(6.2) \quad \Lambda^{\mathbf{S}-\vee} = \{f \in R_a(F') \mid h^\mathbf{S}(f, g) \in \mathcal{O}_{F'} \text{ for all } g \in \Lambda\}.$$

Definition 6.2. The Siegel relative affine Springer fiber $\mathcal{M}^\mathbf{S}(a)$ is the functor classifying all $R_a(\mathcal{O}_{F'})$ -lattices Λ in $R_a(F')$ such that $\iota_R(\Lambda) = \Lambda$ and Λ is $h^\mathbf{S}$ -self-dual, i.e. for any finitely generated k -algebra R ,

$$\mathcal{M}_v^\mathbf{S}(a)(R) = \{\Lambda \subset R_a(F') \hat{\otimes}_k R \mid \Lambda \text{ is an } R_a(\mathcal{O}_{F'})\text{-lattice satisfying } \iota_R(\Lambda) = \Lambda \text{ and } \Lambda^{\mathbf{S}-\vee} = \Lambda\},$$

where $\hat{\otimes}_k$ is the completed tensor product.

Note that the Hermitian form $h^\mathbf{S}$ restricts to zero on the vector subspace $R_a(F')^{\iota_R}$. In particular, it is ill-suited for $\mathcal{M}_v^\mathbf{U}(a)$, which is meant to model the local case where the Hermitian form restricts to a non-degenerate Hermitian form on $R_a(F')^{\iota_R}$. We instead define the form $h^\mathbf{U}: R_a(F') \times R_a(F') \rightarrow F'$ as follows: Denote by $R_a(F')^{\pm \iota_R}$ the subset of invariant/anti-invariant functions $f \in R_a(F')$ under the involution ι_R . For any $f, g \in R_a(F')^{\iota_R}$, respectively $f, g \in R_a(F')^{-\iota_R}$, we have $f \cdot \sigma_R(g) \in R_a(F')^{\iota_R}$ and so we may define

$$h^\mathbf{U}(f, g) = \text{tr}_{R_a(F')^\vee/F'} \left(\frac{f \cdot \sigma_R(g)}{Q'_a(x^2)} \right)$$

where $Q_a(x) = x^n + a_2x^{n-1} + \cdots + a_{2n}$ is the polynomial in t such that $Q_a(x^2) = P_a(x)$. When $f \in R_a(F')^{\iota_R}$ and $g \in R_a(F')^{-\iota_R}$, we put $h^\mathbf{U}(f, g) = 0$. One may now easily verify that $h^\mathbf{U}$ induces a non-degenerate Hermitian form on $R_a(F')^{\iota_R}$.

For any lattice Λ , we may define the dual lattice $\Lambda^{\mathbf{U}-\vee}$ in the same way as (6.2), and we again say that λ is $h^\mathbf{U}$ -self-dual if $\Lambda = \Lambda^{\mathbf{U}-\vee}$.

Definition 6.3. The unitary relative affine Springer fiber $\mathcal{M}^\mathbf{U}(a)$ is the functor classifying all $R_a(\mathcal{O}_{F'})$ lattices Λ in $R_a(F')$ such that $\iota_R(\Lambda) = \Lambda$ and Λ is $h^\mathbf{U}$ -self-dual, i.e. For any k algebra R , $\mathcal{M}^\mathbf{U}(a)(R) = \{\Lambda \subset R_a(F') \hat{\otimes}_k R \mid \Lambda \text{ is an } R_a(\mathcal{O}_{F'}) \hat{\otimes}_k R\text{-lattice satisfying } \iota_R(\Lambda) = \Lambda \text{ and } \Lambda^{\mathbf{U}-\vee} = \Lambda\}.$

Example 6.4. We compute the k -points of the spaces $\mathcal{M}_v^*(a)$ above in the case $n = 1$. Let $a = \varpi_v^m$. We have

$$R_a(F') = F'[x]/(x^2 + \varpi_v^m).$$

It is easy to see that a lattice $\Lambda \subset R$ which is fixed by the involution ι_R is generated by $\varpi^i, \varpi^j x$, where $j \leq i \leq j + m$.

The Hermitian form $h^{\mathbf{S}}$ can be computed by means of Euler's formula: in terms of the F' -basis $\{1, t\}$ for $R_a(F')$, it has matrix

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

In particular, we can compute $\Lambda^{\mathbf{S}-\vee} = \langle \varpi_v^{-j}, \varpi_v^{-i} x \rangle$, and so Λ is $h^{\mathbf{S}}$ -self-dual if and only if $\Lambda = \langle \varpi_v^i, \varpi_v^{-i} x \rangle$ for $0 \leq 2i \leq m$.

The Hermitian form $h^{\mathbf{U}}$ can be computed in terms of the same F' -basis as

$$\begin{pmatrix} 1 & 0 \\ 0 & \varpi_v^m \end{pmatrix}.$$

We have $\Lambda^{\mathbf{U}-\vee} = \langle \varpi_v^{-i}, \varpi_v^{m-j} x \rangle$, and so Λ is $h^{\mathbf{U}}$ -self-dual if and only if m is even and $\Lambda = \langle 1, \varpi_v^{m/2} x \rangle$.

We note that we have $\mathcal{M}_v^{\mathbf{S}}(a)(k') = \mathcal{M}_v^{\mathbf{U}}(a)(k')$ where k' is the quadratic extension k' of k for which $F' \otimes_k k'$ splits, which in all cases is an infinite disjoint union of points.

Corollary 6.5. *The functors $\mathcal{M}^*(a)$ (for $\star = \mathbf{S}$ or $\mathbf{U}$) are representable by projective ind-schemes which are locally of finite type.*

Proof. The space of $R_a(F')$ -lattices is representable by a projective ind-scheme which is locally of finite type by [KL88, §3, Proposition 1]. As the above moduli spaces of lattices are closed subvarieties of the moduli space of all $R_a(F')$ -lattices, the result is now immediate. $\square$

Before concluding this section, we remark on another application of the local rings $R_a(F')$ to lifting elements $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$. The following result will be used to give a more conceptual definition of RASFs in the following section.

Proposition 6.6. *For $\star \in \{\mathbf{S}, \mathbf{U}\}$ and for any $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_F)$, there is a canonical lift $[\gamma] \in [V^*/H^*](F)$ of a .*

Proof. Let $[x]$ denote the multiplication-by- x endomorphism on the ring $R_a(F')$. We claim that the tuple $(R_a(F'), h^*, [x])$ gives an F -point of the stack $[V^*/H^*]$, with the characteristic polynomial of $[x]$ being given by $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_F)$. For this, we note that an F -point of the stack $[V^*/H^*]$ is the following data:

- Two rank n vector bundles $\mathcal{E}'_1, \mathcal{E}'_2$ on $\text{Spec}(F')$ with Hermitian structure h on $\mathcal{E}'_1 \oplus \mathcal{E}'_2$ which either ($\star = \mathbf{U}$) is a direct sum of Hermitian forms on $\mathcal{E}'_1$ and $\mathcal{E}'_2$ or ($\star = \mathbf{S}$) is a direct sum of isomorphisms $\mathcal{E}'_1 \simeq \sigma^* \mathcal{E}'_2^\vee$ and $\mathcal{E}'_2 \simeq \sigma^* \mathcal{E}'_1^\vee$;
- Endomorphisms $\phi'_{ij}: \mathcal{E}'_i \rightarrow \mathcal{E}'_j$ for $i \neq j$ in $\{1, 2\}$ such that $\phi' = \phi'_{12} \oplus \phi'_{21}$ satisfies $\sigma^*(\phi')^\vee \circ h + h \circ \phi' = 0$.

Then, we identify $\mathcal{E}'_1 = R_a(F')^\iota$ with elements represented by polynomials concentrated in even degrees and $\mathcal{E}'_2 = x \cdot R_a(F')^\iota$ with those in odd degree. We take $\phi = [x]$, which interchanges the two. We can check that the condition on the Hermitian form is satisfied for the forms h^* on $R_a(F')$. $\square$

6.2. General RASFs. For the rest of this paper, we denote $\mathcal{O}_v = k[[\varpi_v]]$ and $F_v = k((\varpi_v))$, together with their unramified extensions $\mathcal{O}'_v$ and F'_v , which we assume to be nonsplit. We also denote $\Sigma_v = \text{Spec}(\mathcal{O}_v)$ and $\Sigma_v^\bullet = \text{Spec}(F_v)$. For the purposes of this section, H^* denotes the quasisplit forms of the relevant symmetric subgroups defined over $\mathcal{O}_v$ constructed in Section 2.2. In particular, H^* splits over the $\mathcal{O}'_v$. Let $\text{Gr}_{H^*} = H^*(F_v)/H^*(\mathcal{O}_v)$ denote the affine Grassmannian of H^* .

To an element $\gamma \in V^{\star, \text{rs}}(F_v) \cap V^{\star}(\mathcal{O}_v)$, we may define $\mathfrak{X}_{\gamma}^{\star, \text{pre}}$ to be the functor sending a k -algebra R to

$$(6.3) \quad \mathfrak{X}_{\gamma}^{\star, \text{pre}}(R) = \{h \in \text{Gr}_{H^{\star}}(R) \mid h^{-1} \cdot \gamma \in V^{\star}(R \hat{\otimes}_k \mathcal{O}_v)\}.$$

We define $\mathfrak{X}_{\gamma}^{\star}$ to be the reduced subfunctor of $\mathfrak{X}_{\gamma}^{\star, \text{pre}}$.

We immediately note that $\mathfrak{X}_{\gamma}^{\star}$ only depends on the image of γ in the stack $[V^{\star}/H^{\star}](F_v)$. Indeed, the image $[\gamma] \in [V^{\star}/H^{\star}](F_v)$ is the data of a tuple:

- when $\star = \mathbf{S}$, $[\gamma] = (\mathcal{E}', h, \phi'_{12}, \phi'_{21})$, where $\mathcal{E}'$ is a rank- n vector bundle over $\Sigma_v^{\bullet} \hat{\otimes}_k k'$; $h: \mathcal{E}' \rightarrow \sigma^*(\mathcal{E}')^{\vee}$ such that $h = \sigma h^{\vee}$; $\phi'_{12}: \mathcal{E}' \rightarrow \sigma^*(\mathcal{E}')^{\vee}$; and $\phi'_{21}: \sigma^*(\mathcal{E}')^{\vee} \rightarrow \mathcal{E}'$; satisfying $\sigma^* \phi'_{12}^{\vee} = \phi'_{21}$.
- when $\star = \mathbf{U}$, $[\gamma] = (\mathcal{E}'_1, \mathcal{E}'_2, h_1, h_2, \phi')$, where $\mathcal{E}'_i$ are rank n vector bundles on $\Sigma_v^{\bullet} \hat{\otimes}_k k'$; $h_i: \mathcal{E}'_i \rightarrow \sigma^*(\mathcal{E}'_i)^{\vee}$ such that $h = \sigma^* h_i^{\vee}$; and $\phi': \mathcal{E}'_1 \rightarrow \mathcal{E}'_2$ satisfying $\sigma^* \phi'^{\vee} \circ h_1 + h_2 \circ \phi' = 0$.

By base change, $[\gamma]$ induces similar data $[\gamma]_R$ for any k scheme R , replacing $\Sigma_v^{\bullet}$ with $\Sigma_v^{\bullet} \hat{\otimes}_k R$. We may reformulate (6.3) as

$$\mathfrak{X}_{[\gamma]}^{\star, \text{pre}}(R) = \left\{ (\mathcal{E}, \xi) \left| \begin{array}{l} \mathcal{E} \in [V^{\star}/H^{\star}](R \hat{\otimes}_k \mathcal{O}_v) \text{ and} \\ \xi: \mathcal{E}|_{\Sigma_v^{\bullet} \hat{\otimes}_k R} \xrightarrow{\sim} [\gamma]_R \text{ is a choice of trivialization} \end{array} \right. \right\}.$$

Proposition 6.7. *Let $a \in \mathfrak{c}^{\heartsuit}(\mathcal{O}_v)$. For $\star \in \{\mathbf{S}, \mathbf{U}\}$, the RASF $\mathcal{M}_v^{\star}(a)$ (see Definitions 6.2 and 6.3) is isomorphic to $\mathfrak{X}_{[\gamma]}^{\star}$ where $[\gamma] \in [V^{\star}/H^{\star}](F_v)$ is the canonical lift of a of Proposition 6.6.*

Proof. Suppose that $(\mathcal{E}, \xi) \in \mathfrak{X}_{[\gamma]}^{\star}(R)$. In particular, $\mathcal{E}$ determines a point $(\Lambda, h_{\Lambda}) \in [\mathfrak{u}_{2n}/\mathfrak{U}_{2n}](\mathcal{O}_v \hat{\otimes} R)$, which is the data of a $R_a(\mathcal{O}_{F'_v})$ -module Λ equipped with a Hermitian form h_{Λ} so that $h_{\Lambda} \circ [x] + [x] \circ h_{\Lambda} = 0$, where $[x]$ denotes the multiplication-by- x operation on Λ . The isomorphism ξ determines, in particular, a choice of isomorphism $\Lambda \otimes_{\mathcal{O}_{F'_v}} F'_v \simeq R_a(F'_v)$, which allows us to view Λ as a submodule of $R_a(F'_v)$.

The Hermitian form h_{Λ} extends to a Hermitian form h_{Λ} on $R_a(F'_v)$ under which Λ is h_{Λ} self dual. The remaining data in the pair $(\mathcal{E}, \xi)$ involve the compatibility of this form with the involution ι . In particular, we see that the induced Hermitian form h_{Λ} must restrict to zero on $R_a(F'_v)^{\iota_R}$ in the Siegel case and must be a direct sum of Hermitian forms on $R_a(F'_v)^{\pm \iota_R}$ in the unitary case. Up to automorphism of $R_a(F_v)$ commuting with ι we can therefore normalize h_{Λ} to $h^{\star}$, which gives a bijection between $\mathfrak{X}_{[\gamma]}^{\star}(R)$ and $\mathcal{M}_v^{\star}(a)(R)$. $\square$

Remark 6.8. While this lattice definition of $\mathcal{M}_v^{\star}(a)$ is special to the unitary Friedberg–Jacquet varieties and other symmetric varieties for classical groups, we expect an identical construction to $\mathfrak{X}_{[\gamma]}^{\star}$ to work for any spherical variety X . The obstruction is to find lifts $[\gamma] \in [T^{\star}X/G](F_v)$ for any $a \in (T^{\star}X//G)^{\heartsuit}(\mathcal{O}_v)$. This is of course nontrivial, and we leave this point to future work.

For the Siegel case, there is a Kostant–Rallis section, which permits us to use the more naive definition in terms of $\mathfrak{X}_{\gamma}^{\mathbf{S}}$. The following ensures this is well defined.

Lemma 6.9. *For $a \in \mathfrak{c}^{\heartsuit}(\mathcal{O}_v)$ and a fixed Kostant–Rallis section $[\epsilon]$, the canonical lift $[\gamma] \in [V^{\mathbf{S}}/H^{\mathbf{S}}](F_v)$ of Proposition 6.6 is isomorphic to the image of $[\epsilon](a) \in V^{\mathbf{S}}(F)$. In particular, $\mathcal{M}_v^{\mathbf{S}}(a) \simeq \mathfrak{X}_{[\gamma]}^{\mathbf{S}} \simeq \mathfrak{X}_{[\epsilon](a)}^{\mathbf{S}}$.*

Proof. The image of $[\epsilon](a) \in V^{\mathbf{S}}(F_v)$ in $[V^{\mathbf{S}}/H^{\mathbf{S}}](F_v)$ consists of the trivial $H^{\mathbf{S}}$ -bundle together equipped with an endomorphism—which can be used to identify it with $R_a(F'_v)$ —and a compatible Hermitian form h^{ϵ} . The Hermitian form h^{ϵ} is characterized up to scaling by the following properties:

- h^{ϵ} is compatible with the endomorphism $[x]$ in the sense that $h^{\epsilon}(x \cdot f, g) + h^{\epsilon}(f, x \cdot g) = 0$ for all $f, g \in R_a(F'_v)$;
- h^{ϵ} restricts to zero on $R_a(F'_v)^{\iota}$.

We may write $h^\epsilon(f, g) = h^{\mathbf{S}}(\eta f, g)$ for some automorphism η , which by the first condition is in $R_a(F'_v)^\times$ and by the second is in $(R_a(F'_v)^\times)^\iota$. In particular, η determines an isomorphism between the image of $[\epsilon](a)$ and $[\gamma]$. $\square$

Remark 6.10. If k'/k is quadratic extension over which our unitary groups split, then $\mathcal{M}_v^{\mathbf{S}}(a) \otimes_k k' \simeq \mathcal{M}_v^{\mathbf{U}}(a) \otimes_k k'$. Hence, Lemma 6.9 together with the conjugation invariance of $\mathfrak{X}_\gamma^*$ imply that the spaces $\mathcal{M}_v^{\mathbf{S}}(a) \otimes_k k'$ and $\mathcal{M}_v^{\mathbf{U}}(a) \otimes_k k'$ can be computed using any lift $\gamma' \in V^*(k')$. We will use this fact for algebraically closed fields in §6.5.

6.3. The Obstruction Class. In this section, we study a class $\text{Ob}_v^*(a)$, which serves as the obstruction to the existence of k -points in the RASFs $\mathcal{M}_v^*(a)$. This obstruction only exists in the unitary case ($\star = \mathbf{U}$) and that it is controlled by the parity of $\text{val}(a_{2n})$.

Given $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$, recall that we may form the ring $R_a(F'_v)$, which comes equipped with a Hermitian form h^* . The tuple $(R_a(F'_v), h^*, [x])$ gives an F_v point of the stack $[\mathbf{u}_{2n}/\mathbf{U}_{2n}](F_v)$, which is the image of the canonical lift $[\gamma]$ of a (see Proposition 6.6) under the map $[V^*/H^*] \rightarrow [\mathbf{u}_{2n}/\mathbf{U}_{2n}]$. In particular, $(R_a(F'_v), h^*)$ gives a $\mathbf{U}_{2n}$ torsor on $\Sigma_v^\bullet = \text{Spec}(F_v)$.

Definition 6.11. The obstruction class $\text{Ob}_v^*(a)$ is the class

$$\text{Ob}_v^*(a) = [(R_a(F'_v), h^*)] \in H^1(F_v, \mathbf{U}_{2n}) \simeq \mathbb{Z}/2\mathbb{Z}$$

If no superscript is written, we mean the unitary case, i.e. $\text{Ob}_v(a) = \text{Ob}_v^{\mathbf{U}}(a)$.

Our convention to drop the superscripts in the unitary case is justified by the following lemma.

Lemma 6.12. *If $\star = \mathbf{S}$, then $\text{Ob}_v^{\mathbf{S}}(a)$ is trivial for all $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$.*

Proof. We can calculate $H^1(F_v, \mathbf{U}_{2n}) = \mathbb{Z}/2\mathbb{Z}$ with a $\mathbf{U}_{2n}$ -bundle (E, h) being taken to $\text{val}_v(\det(h)) \bmod 2$. But by our definition of $h^{\mathbf{S}}$ and Euler's formula for trace, the matrix of $h^{\mathbf{S}}$ with respect to the basis $\{1, x, \dots, x^{2n-1}\}$ of $R_a(F'_v)$ is lower triangular, with 1's on the anti-diagonal and nonzero entries only possible in the lower right triangular entries. In particular, $\det(h^{\mathbf{S}}) = 1$, so the class of $(R_a(F'_v), h^{\mathbf{S}})$ in $H^1(F_v, \mathbf{U}_{2n})$ is trivial. $\square$

The unitary case is more interesting. The following computes $\text{Ob}_v(a) = \text{Ob}_v^{\mathbf{U}}(a)$.

Lemma 6.13. *If $\star = \mathbf{U}$, then*

$$\text{Ob}_v(a) = \text{val}(a^* \mathfrak{D}^{\text{br}}) = \text{val}(a_{2n}) \in \mathbb{Z}/2\mathbb{Z}$$

is the parity of the valuation of the determinant component of $a = (a_2, \dots, a_{2n}) \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$.

Proof. Similar to Lemma 6.12, it suffices to calculate the valuation of $\det(h^{\mathbf{U}})$. By Euler's formulas (using an identical argument to Lemma 6.12 using the basis $1, x^2, \dots, x^{2n-2}$), we know the restriction of $h^{\mathbf{U}}$ to $R_a(F'_v)^{\iota_R}$ has determinant 1. On the other hand, the restriction of $h^{\mathbf{U}}$ to $R_a(F'_v)^{-\iota_R}$ can be computed with respect to the basis $x, \dots, x^{2n-1}$ as follows. We can identify $R_a(F'_v)^{\iota_R} \simeq R_a(F'_v)^{-\iota_R}$ as F'_v -vector spaces by multiplying by x and under this identification, the form $h^{\mathbf{U}}|_{R_a(F'_v)^{-\iota_R}}$ is identified with the composition of $h^{\mathbf{U}}|_{R_a(F'_v)^{\iota_R}}$ and the multiplication-by- x^2 endomorphism. As the former has determinant 1 and the latter has determinant a_{2n} , we conclude that $h^{\mathbf{U}}$ restricted to $R_a(F'_v)^{-\iota_R}$ has determinant a_{2n} . Since $h^{\mathbf{U}}$ is defined so that $R_a(F'_v)^{\iota_R}$ and $R_a(F'_v)^{-\iota_R}$ are orthogonal with respect to $h^{\mathbf{U}}$, it follows that $\det(h^{\mathbf{U}}) = a_{2n}$, and the lemma follows. $\square$

Finally, we prove that the obstruction class indeed captures the existence of rational points. This will end up being a non-vanishing statement for stable orbital integrals.

Proposition 6.14. *The obstruction class $\text{Ob}_v^*(a)$ is trivial if and only if the RASF $\mathcal{M}_v^*(a)$ has a k point. In particular, $\mathcal{M}_v^{\mathbf{U}}(a)$ has a k point if and only if $\text{val}(a^* \mathfrak{D}^{\text{br}}) = \text{val}(a_{2n})$ is even.*

Proof. We always have $\text{Ob}_v^S(a) = 0$ by Lemma 6.12. Moreover, by the existence of the Kostant-Rallis section, we have $\mathcal{M}_v^S(a)(k) \neq \emptyset$ for all $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$. Therefore, the only nontrivial statement is for $\star = \mathbf{U}$.

Suppose first that there exists a $h^\mathbf{U}$ -self-dual lattice Λ such that $\iota_R(\Lambda) = \Lambda$. Then, the Hermitian form $h^\mathbf{U}$ restricts to a perfect pairing on Λ , and so must have determinant a norm. Hence, the $\mathbf{U}_{2n}$ bundle $(R_a(F'_v), h^\mathbf{U})$ is trivial, and $\text{Ob}_v(a) = 0$. We note that this argument proves the stronger claim that, if $\text{Ob}_v(a) \neq 0$, then there are no $h^\mathbf{U}$ -self-dual lattices (regardless of stability under ι_R).

Now, suppose that $\text{Ob}_v(a) = 0$. Then, $(R_a(F'_v), h^\mathbf{U})$ is the trivial $\mathbf{U}_{2n}$ bundle, and moreover, by our formula for $h^\mathbf{U}$ and the proof of Lemma 6.13, the restriction of $h^\mathbf{U}$ to $R_a(F'_v)^{\iota_R}$ and to $R_a(F'_v)^{-\iota_R}$ are each isometric to the trivial Hermitian form. That is, there exist bases $\mathcal{B}_\pm$ for $R_a(F'_v)^{\pm\iota_R}$ with respect to which $h^\mathbf{U}$ has matrix the identity I_n when restricted to $R_a(F'_v)^{\pm\iota_R}$. In particular, we can consider the lattices $\Lambda_\pm \subset R_a(F'_v)^{\pm\iota_R}$, defined by

$$\Lambda_\pm = \bigoplus_{e \in \mathcal{B}_\pm} \mathcal{O}_{F_v} \cdot e$$

Then $\Lambda = \Lambda_+ \oplus \Lambda_-$ is self dual and satisfies $\iota_R(\Lambda) = \Lambda$. $\square$

6.4. Symmetries of RASFs. Let $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v) := \mathfrak{c}^{\text{rs}}(F_v) \cap \mathfrak{c}(\mathcal{O}_v)$. For any group J'_a over $\mathcal{O}_v$ with generic fiber $J'_a(F_v) \simeq J_a(F_v)$ agreeing with the group scheme of regular centralizers J_a , we consider the affine Grassmannian

$$\mathcal{P}_v(J'_a) := \text{Gr}_{J'_a} = J_a(F_v)/J'_a(\mathcal{O}_v)$$

For each $\gamma \in V^\star(F_v)$ lifting a , we obtain an identification of the fiber $J_a(F_v)$ with the centralizer $I_\gamma \subset H^\star(F_v)$. The latter group acts on the scheme $\mathfrak{X}_\gamma^\star$ by left multiplication. This action factors through $\mathcal{P}_v(J'_a)$, and we deduce the following.

Proposition 6.15. *For any $\gamma \in V^\star(F_v)$ lifting a , the group scheme $\mathcal{P}_v(J'_a)$ acts on $\mathfrak{X}_\gamma^\star$. In particular, there is an action of $\mathcal{P}_v(J'_a)$ on $\mathcal{M}_v^\star(a)$.*

Analogously to the classical setting, this action consists of a free action by a lattice in $\mathcal{P}_v(J'_a)$ together with an action of a group scheme of finite type. More precisely, for $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$, we let Γ_a denote the lattice $\Gamma_a = \mathbb{X}_*(J_a)_{\check{F}_v}$ of $\check{F}_v$ -rational cocharacters. Then Γ_a inherits an action of Frobenius, and hence descends to an ind-scheme over k . If $\gamma \in V^\star(F_v)$ is a lift of a , and $I_\gamma \subset H^\star(F_v)$ its centralizer group, then we have a canonical identification $\Gamma_a \simeq \mathbb{X}_*(I_\gamma)_{\check{F}_v}$. More explicitly, suppose that we have a factorization

$$R_a(F_v)^{\iota_R} \simeq \prod_{j=1}^r K_j$$

where K_j is a finite extension of F_v for all j . Let ϖ_j be a uniformizer of K_j . Then, $(m_1, \dots, m_r) \in \Gamma_a \otimes_k k' \simeq \mathbb{Z}^r$ acts on $\mathcal{M}_v^\star(a) \otimes_k k'$ by multiplication by $(\varpi_1^{m_1}, \dots, \varpi_r^{m_r}) \in R_a(F_v)^{\iota_R}$.

Proposition 6.16. *Fix $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$.*

- (1) *The action of Γ_a on $\mathcal{M}_v^\star(a)$ is free;*
- (2) *The quotient $\mathcal{M}_v^\star(a)/\Gamma_a$ is a finite-type projective scheme, and $\mathcal{P}_v(J_a)/\Gamma_a$ is a commutative group scheme of finite type;*
- (3) *The stabilizers of the action of $\mathcal{P}_v(J_a)$ on $\mathcal{M}_v^\star(a)$ are finite type and affine.*

Proof. As all claims are geometric, we will assume throughout that $k = \bar{k}$. Put $c = \text{val} \left(\text{Disc}(P_a) \right)$.

The proof will be via explicit computation. For simplicity throughout this proof, we denote $R = R_v(\mathcal{O}'_v) \otimes_k \bar{k}$ and $\tilde{R}$ the normalization of R . If we decompose

$$R \otimes_{\mathcal{O}'_v} F'_v = R_v(F'_v) \simeq \prod_j E_j$$

as a product of finite extensions E_i of F'_v , then $\tilde{R} \simeq \prod_j \mathcal{O}_{E_j}$. We denote by

$$v: R_a(F'_v)^\times / \tilde{R}^\times \xrightarrow{\sim} \prod_j \mathbb{Z}$$

the valuation map on each $E_j^\times$. As $\tilde{R}$ is stable under the involution ι , we can transfer ι to the right-hand side, and so can define $\Gamma = \left(\prod_j \mathbb{Z} \right)^\iota$. The valuation map v determines an embedding $v: \Gamma_a \rightarrow \Gamma$, which because Γ_a and Γ are free abelian groups of the same rank, has finite cokernel.

Since $\tilde{R}$ is integral, we must have $\tilde{R} \subset R^\vee$, where $R^\vee = (P'_a)^{-1} \cdot R$ is the dual $\mathcal{O}'_v$ -lattice to R in $R_a(F'_v)$, where P_a is as in (6.1) and $(P'_a)^{-1} \cdot R$ denotes the fractional ideal generated by $(P'_a)^{-1}$. As $\text{Disc}(P_a) \in P'_a \cdot R$, it follows that $\varpi_v^c \tilde{R} \subset R$. On the other hand, since every $\tilde{R}$ -lattice in $R_v(F'_v)$ is principal for every R -lattice Λ , there exists some $f_\Lambda \in R_v(F'_v)$ such that $\tilde{R} \cdot \Lambda = f_\Lambda \cdot \tilde{R}$. In particular, we can define $v(\Lambda) := v(f_\Lambda)$. Note that $v(\Lambda) \in \Gamma$ when Λ is an ι -fixed lattice. From the inclusion $\varpi_v^c \tilde{R} \subset R$ of the previous paragraph, we get

$$(6.4) \quad \varpi_v^c f_\Lambda \cdot \tilde{R} \subset \Lambda \subset f_\Lambda \cdot \tilde{R}.$$

Now, let us prove (1). If $\check{\lambda} \cdot \Lambda = \Lambda$ for $\check{\lambda} \in \Gamma_a$ and $\Lambda \in \mathcal{M}_v^*(a)(\bar{k})$, then taking valuations gives

$$v(\Lambda) = v(\check{\lambda} \cdot \Lambda) = v(\check{\lambda}(\varpi_v)) + v(\Lambda),$$

so that $v(\check{\lambda}(\varpi_v)) = 0$. Since v is injective on Γ_a , this implies that $\check{\lambda} = 0$, and so Γ_a acts freely on $\mathcal{M}_v^*(a)$.

We prove (2) by constructing a finite type, projective subvariety of $\mathcal{M}_v^*(a)$ which contains a fundamental domain for the Γ_a -action. For each of the finitely many cosets $s \in \Gamma/\Gamma_a$, we take a representative $f_s \in R_v(F'_v)^\times$. Define Y_s to be the moduli of all ι -fixed R -lattices Λ such that

$$\varpi_v^c f_s \cdot \tilde{R} \subset \Lambda \subset f_s \cdot \tilde{R}$$

We can view Y_s as a closed subvariety of the Grassmannian of the finite dimensional vector space $f_s \cdot \tilde{R} / \varpi_v^c f_s \cdot \tilde{R}$, so Y_s is a projective algebraic variety of finite type. Consider the space

$$Y_{\mathcal{M}} := \mathcal{M}_v^*(a) \cap \left(\bigcup_{s \in \Gamma/\Gamma_a} Y_s \right).$$

Then, for any $\Lambda \in \mathcal{M}_v^*(a)$, pick $\check{\lambda} \in \Gamma_a$ and $s \in \Gamma/\Gamma_a$ such that $v(\Lambda) - v(\check{\lambda}(\varpi_v)) = v(f_s)$. Then, $\tilde{R} \cdot (\check{\lambda}^{-1} \Lambda) = f_s \cdot \tilde{R}$, and so by (6.4), we see that $\check{\lambda}^{-1} \cdot \Lambda \in Y_s \subset Y_{\mathcal{M}}$. We conclude that $Y_{\mathcal{M}}$ contains a fundamental domain for the Γ_a -action, and part (2) follows.

Consider now the symmetry group $\mathcal{P}_v(J_a)$. We compute $\mathcal{P}_v(J_a) \simeq (R^\iota \otimes_{\mathcal{O}'_v} F'_v)^\times / (R^\iota)^\times$. Therefore, $\mathcal{P}_v(J_a)$ fits into a short exact sequence

$$0 \rightarrow (\tilde{R}^\iota)^\times / (R^\iota)^\times \rightarrow \mathcal{P}_v(J_a) \rightarrow (R^\iota \otimes_{\mathcal{O}'_v} F'_v)^\times / (\tilde{R}^\iota)^\times \rightarrow 0$$

where $\tilde{R}^\iota$ is the normalization of R^ι . In particular, $(R^\iota \otimes_{\mathcal{O}'_v} F'_v)^\times / (\tilde{R}^\iota)^\times \simeq \prod_i \mathbb{Z}$ is a product of copies of $\mathbb{Z}$ since $(\tilde{R}^\iota)^\times$ splits as a product of DVR's, and the composition $\Gamma_a \rightarrow (R^\iota \otimes_{\mathcal{O}'_v} F'_v)^\times / (\tilde{R}^\iota)^\times$ is injective between abelian groups of the same rank and hence is of finite index. Therefore, in the resulting exact sequence

$$0 \rightarrow (\tilde{R}^\iota)^\times / (R^\iota)^\times \rightarrow \mathcal{P}_v(J_a) / \Gamma_a \rightarrow \left[(R^\iota \otimes_{\mathcal{O}'_v} F'_v)^\times / (\tilde{R}^\iota)^\times \right] / \Gamma_a \rightarrow 0,$$

the cokernel is finite and the kernel is affine and of finite type. Hence, $\mathcal{P}_v(J_a) / \Gamma_a$ is finite type. By (1), the stabilizer subgroup of any $m \in \mathcal{M}_v^*(a)(\bar{k})$ under the action of $\mathcal{P}_v(J_a)$ can be viewed as a subgroup of $\mathcal{P}_v(J_a) / \Gamma_a$, which necessarily must be affine and of finite type since it is an extension of a finite type, affine group by a finite group. $\square$

Remark 6.17. In contrast to the case of usual affine Springer fibers, the lattice Γ_a does not act transitively on the set of irreducible components of $\mathcal{M}_v^*(a)$. However, for each connected component of $\mathcal{M}_v^*(a)$, we will see that the lattice action is free and transitive on the set of irreducible components in that connected component of $\mathcal{M}_v^*(a)$.

6.5. Dimension in the split and transverse cases. The relative affine springer fibers $\mathcal{M}_v^*(a)$ are particularly simple in the case of *split* $a \in \mathfrak{c}^\vee(\mathcal{O}_v)$. We will give an explicit description of the dimension and component groups in this case, which we will use in Section 6.8 to describe a general dimension formula.

Definition 6.18. We say $a \in \mathfrak{c}^\vee(\mathcal{O}_v)$ is *split* if the centralizer J_a is a split torus over F_v .

Recall that we use a superscript breve to denote base change to $\bar{k}$; for example $\check{F}_v = F_v \hat{\otimes}_k \bar{k}$, $\check{\mathcal{O}}_v = \mathcal{O}_v \hat{\otimes}_k \bar{k}$, etc.

Lemma 6.19. *Let $a \in \mathfrak{c}^\vee(\mathcal{O}_v)$ be split. Then, there is a lift $\gamma \in \mathfrak{h}^\perp(\check{F}_v)$ of a of the form*

$$\gamma = \begin{pmatrix} 0 & A \\ I_n & 0 \end{pmatrix} \quad \text{where } A = \begin{pmatrix} u_1 \varpi_v^{\delta_1} & & & \\ & u_2 \varpi_v^{\delta_2} & & \\ & & \ddots & \\ & & & u_n \varpi_v^{\delta_n} \end{pmatrix}$$

where $0 \leq \delta_1 \leq \dots \leq \delta_n$ and $u_i \in \mathcal{O}_v^\times$ such that $u_i \neq u_j$ whenever $\delta_i = \delta_j$. This lift is unique up to reordering of pairs u_i, u_j such that $\delta_i = \delta_j$

Proof. Choose any lift $\gamma \in \mathfrak{h}^\perp(\check{F}_v)$ of a . We have a canonical identification $I_\gamma \simeq J_a(\check{F}_v)$, which is a split torus over $\check{F}_v$ by assumption. Hence, the image of I_γ under the each of the two projections $H(\check{F}_v) = \mathrm{GL}_n(\check{F}_v) \times \mathrm{GL}_n(\check{F}_v) \rightarrow \mathrm{GL}_n(\check{F}_v)$ is a split maximal torus of $\mathrm{GL}_n(\check{F}_v)$, and up to $H(\check{F}_v)$ conjugacy, we may assume that the image of I_γ under each projection is the standard maximal torus consisting of diagonal matrices. Now, let

$$\gamma = \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$$

and denote $T_{\mathrm{GL}_n} \subset \mathrm{GL}_n$ and $T \subset H$ the standard diagonal maximal tori. Then, it is easy to see that $(g, h) \in I_\gamma \subset T(\check{F}_v)$ implies that $g \in C_{\mathrm{GL}_n(\check{F}_v)}(BA)$ and $h \in C_{\mathrm{GL}_n(\check{F}_v)}(AB)$. In particular, we see that $T_{\mathrm{GL}_n}(\check{F}_v) \subset C_{\mathrm{GL}_n(\check{F}_v)}(BA)$ and $T_{\mathrm{GL}_n}(\check{F}_v) \subset C_{\mathrm{GL}_n(\check{F}_v)}(AB)$. Since we have an equality of characteristic polynomials (with λ a variable)

$$\mathrm{char.poly}(\gamma, \lambda) = \mathrm{char.poly}(AB, \lambda^2) = \mathrm{char.poly}(BA, \lambda^2),$$

it follows that AB and BA are regular, semisimple over $\check{F}_v$ with no zero eigenvalues. Hence, we conclude that AB and BA are diagonal, and since A and B have no zero eigenvalues, A and B are invertible. Therefore, we have

$$\begin{pmatrix} I_n & 0 \\ 0 & B^{-1} \end{pmatrix} \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \begin{pmatrix} I_n & 0 \\ 0 & B \end{pmatrix} = \begin{pmatrix} 0 & AB \\ I_n & 0 \end{pmatrix}$$

where AB is diagonal, regular semisimple. This representation is unique up to reordering of the diagonal entries of AB , and so the lemma follows. $\square$

We now prove a dimension formula for affine Springer fibers for split a . Note that, by Remark 6.10, we can compute $\mathcal{M}_v^*(a) \otimes_k \bar{k} \simeq \mathfrak{X}_\gamma^*$ using the lift $\gamma \in V^*(\check{F}_v) = \mathfrak{h}^\perp(\check{F}_v)$ of Lemma 6.19.

We may identify $\mathfrak{c} \simeq \mathfrak{c}_{\mathrm{GL}_n}$, with the corresponding $\mathbb{G}_m$ actions doubled in $\mathfrak{c}$. More explicitly, for $a = (a_1, \dots, a_n) \in \mathfrak{c}_{\mathrm{GL}_n}$ thought of as a characteristic polynomial

$$P_a(\lambda) = \lambda^n + a_1 \lambda^{n-1} + \dots + a_n,$$

we send a to the same n -tuple in $\mathfrak{c}$, thought of as the polynomial $P_a(\lambda^2)$. If $a \in \mathfrak{c}$, we call $a_{\text{GL}} \in \mathfrak{c}_{\text{GL}_n}$ as above the *associated element to a in $\mathfrak{c}_{\text{GL}_n}$* .

The following (without component groups) appears in [Pol15]. We give a simpler argument here.

Proposition 6.20. *Let $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$ be split with a_{GL} the associated element in $\mathfrak{c}_{\text{GL}_n}(\check{\mathcal{O}}_v)$. Then, the RASF $\mathcal{M}_v^*(a)$ is equidimensional with*

$$(6.5) \quad \dim \mathcal{M}_v^*(a) = \dim \mathcal{P}_v(J_a) = \dim \mathcal{M}_v(a_{\text{GL}})$$

where $\mathcal{M}_v(a_{\text{GL}})$ denotes the affine Springer fiber for the group $G = \text{GL}_n$ over a_{GL} . Moreover, if

$$\gamma = \begin{pmatrix} 0 & A \\ I_n & 0 \end{pmatrix} \quad \text{where } A = \begin{pmatrix} u_1 \varpi_v^{\delta_1} & & & \\ & u_2 \varpi_v^{\delta_2} & & \\ & & \ddots & \\ & & & u_n \varpi_v^{\delta_n} \end{pmatrix}$$

is a lift of a as in Lemma 6.19, then

$$\text{Irr}(\mathcal{M}_v^*(a)) \simeq \{(m^{(1)}, m^{(2)}) \in \mathbb{Z}^n \times \mathbb{Z}^n : 0 \leq m_i^{(1)} - m_i^{(2)} \leq \delta_i\}$$

with the action of the Frobenius σ for k coming from

$$\sigma(m^{(1)}, m^{(2)}) = \begin{cases} (-m^{(1)}, -m^{(2)}) & \text{in the unitary case, } \star = \mathbf{U} \\ (-m^{(2)}, -m^{(1)}) & \text{in the Siegel case, } \star = \mathbf{S} \end{cases}$$

Proof. To compute dimension and components, it is harmless to assume that $k = \bar{k}$ is algebraically closed and identify H with $\text{GL}_n \times \text{GL}_n$. We begin by taking γ to be slightly more general than the above form: Assume that

$$(6.6) \quad \gamma = \begin{pmatrix} 0 & A' \\ A'' & 0 \end{pmatrix}$$

where A', A'' are diagonal and $A'A'' = A$ is of the form of Prop 6.19. Consider the unipotent subgroup of H , which we denote by $N = N_n \times N_n$, with N_n the standard upper triangular unipotent subgroup of GL_n . Denote by $\mathfrak{X}_\gamma^N$ the unipotent RASF whose $\bar{k}$ points are given by

$$\mathfrak{X}_\gamma^N(\bar{k}) = \{n \in \text{Gr}_N \mid \text{Ad}(n)^{-1} \cdot \gamma \in \mathfrak{h}^\perp(\mathcal{O}_v)\}.$$

We may similarly define $\mathfrak{X}_{\gamma'}^{N_n}$ for $\gamma' \in \mathfrak{gl}_n(\mathcal{O}_v)$ a regular semisimple nil-element. Let $\gamma' = A = A'A''$ denote the diagonal matrix in the upper right block of γ . We have a map

$$(6.7) \quad \mathfrak{X}_\gamma^N \rightarrow \mathfrak{X}_{\gamma'}^{N_n} \times \mathfrak{X}_{\gamma'}^{N_n}$$

taking n to each of its components (n_1, n_2) . A simple calculation shows that $n = (n_1, n_2) \in N(F_v)$ represents an element of $\mathfrak{X}_\gamma^N$ if and only if $n_2^{-1}n_1, n_1^{-1}An_2 \in \mathfrak{n}_n(\mathcal{O}_v)$. Since n_2 and n_1 are invertible in a unipotent subgroup, the former condition holds if and only if $n_1 = n_2 \cdot N_n(\mathcal{O}_v)$, and so the above two conditions are equivalent to the conditions $n_1, n_2 \in \mathfrak{X}_{\gamma'}^{N_n}$ and $n_1 = n_2 \cdot N_n(\mathcal{O}_v)$. That is, the map (6.7) identifies $\mathfrak{X}_\gamma^N$ with the diagonal in the product $\mathfrak{X}_{\gamma'}^{N_n} \times \mathfrak{X}_{\gamma'}^{N_n}$.

Now consider the Iwasawa decomposition $h = tnk$ for $t \in T(F_v)$ and $n \in N(F_v)$ uniquely determined up to right multiplication by $T(\mathcal{O}_v)$ and $N(\mathcal{O}_v)$, respectively. Note that when $t^{-1} \cdot \gamma$ is in $\mathfrak{h}^\perp(\mathcal{O}_v)$, it is of the form (6.6), and hence, $\mathfrak{X}_{t^{-1} \cdot \gamma}^N \simeq \mathfrak{X}_\gamma^N$. Let $\delta = (\delta_1, \dots, \delta_n)$ denote the exponents of the diagonal entries of A . Then $t^{-1} \cdot \gamma$ is integral valued when $t^{-1} \cdot \gamma = \gamma_{\eta, \eta'}$ where η, η' are n -tuples of nonnegative integers satisfying $\eta + \eta' = \delta$. We have a morphism

$$\mathfrak{X}_\gamma \rightarrow \bigsqcup_{\substack{\eta + \eta' = \delta \\ \eta, \eta' \geq 0}} \mathfrak{X}_{\gamma_{\eta, \eta'}}^N$$

whose fibers are discrete ind-schemes of dimension zero, which are torsors for the action of the lattice Γ_a . In particular, we at once deduce the claimed dimension formula and the description of component groups.

It is a simple exercise to see the σ -action on $\mathrm{Gr}_H \simeq \mathrm{Gr}_{\mathrm{GL}_n} \times \mathrm{Gr}_{\mathrm{GL}_n}$ restricts to the actions described in the proposition statement. $\square$

Remark 6.21. An explicit formula for the dimension (6.5) will be given in §6.8 using basic invariants associated to a .

We conclude this section with the *transverse* case. Namely, we prove:

Proposition 6.22. *Suppose that $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$ intersects with the full discriminant divisor $\mathfrak{D}_X$ transversely. Then, the ring $R_a(\mathcal{O}'_v)$ is regular. Moreover, the affine Springer fiber $\mathcal{M}_v^*(a)$ has dimension zero.*

Proof. It suffices to show that every $\bar{k}$ -point of $\mathrm{Spec} R_a(\mathcal{O}'_v)$ is regular. An $\bar{k}$ -point of $R_a(\mathcal{O}'_v)$ is an element $x_0 \in \bar{k}$ such that $P_a(x_0) \equiv 0 \pmod{\varpi_v}$. Such an x_0 is non-regular if and only if $P_a(x) \in \mathfrak{m}_{x_0}^2$ where $\mathfrak{m}_{x_0} = (\varpi_v, x - x_0) \subset R_a(\mathcal{O}'_v)$ is the maximal ideal corresponding to x_0 . More explicitly, let us write

$$P_a(x) = \sum_{i \geq 0} P_i(x) \varpi_v^i, \quad P_i(x) \in k[x].$$

Then, x_0 is non-regular if $(x - x_0)^2$ divides $P_0(x)$ and $P_1(x_0) = 0$.

Suppose that x_0 is a non-regular $\bar{k}$ -point of $\mathrm{Spec} R_a(\mathcal{O}'_v)$. If $x_0 = 0$, then $P_a(x_0) \equiv 0 \pmod{\varpi_v}$ implies $a_{2n} \equiv 0 \pmod{\varpi_v}$, and $P_1(x_0) = 0$ then implies that $a_{2n} \equiv 0 \pmod{\varpi_v^2}$. Therefore, $\mathrm{val}(a_{2n}) \geq 2$, contradicting transversality with $\mathfrak{D}^{\mathrm{br}} = (a_{2n}) \subset \mathfrak{D}_X$.

If instead $x_0 \neq 0$, then we claim that x_0^2 is a nonzero, non-regular point of $\mathrm{Spec} R_a(\mathcal{O}'_v)^{\iota_R}$. Indeed,

$$R_a(\mathcal{O}'_v)^{\iota_R} = \mathcal{O}'_v[y]/(Q_a(x)), \quad Q_a(x) = x^n + a_2 x^{n-1} + a_{2n-2} x^{n-2} + \cdots + a_{2n}$$

with $Q_a(x^2) = P_a(x)$. In particular, x_0^2 is a closed point of $\mathrm{Spec} R_a(\mathcal{O}'_v)^{\iota_R}$. If we express

$$Q_a(x) = \sum_{i \geq 0} Q_i(x) \varpi_v^i, \quad Q_i(x) \in k[x],$$

then one can further check that $Q_1(x_0^2) = P_1(x_0)$ so that x_0^2 is non-regular as a point of $\mathrm{Spec} R_a(\mathcal{O}'_v)^{\iota_R}$ if and only if x_0 is a non-regular point of $\mathrm{Spec} R_a(\mathcal{O}'_v)$. Since $\mathfrak{D}^{\mathrm{cent}} = (\mathrm{Disc} Q_a(x^2))$, transversality implies $\mathrm{val} \mathrm{Disc} Q_a(x_0^2) \leq 1$. Hence, the regularity comes from the following lemma.

Lemma 6.23. *Let $Q \in \mathcal{O}'_v[x]$ be a monic polynomial with $\mathrm{val} \mathrm{Disc}(Q) \leq 1$. Then $R_Q = \mathcal{O}'_v[x]/(Q)$ is a regular local ring.*

Proof. Since $\mathrm{Disc}(Q) \neq 0$, R_Q is reduced. Let $\widetilde{R_Q}$ be its normalization. Since the trace of an integral element is integral, we have $R_Q \subset \widetilde{R_Q} \subset (R_Q)^\vee$, where $(R_Q)^\vee$ is the dual lattice to R_Q in $R_Q \otimes_{\mathcal{O}'_v} F'_v$. Hence $\widetilde{R_Q}$ is also a $\mathcal{O}'_v$ lattice in $R_Q \otimes_{\mathcal{O}'_v} F'_v$. Let M be the matrix expressing a basis of R_Q in terms of a basis of $\widetilde{R_Q}$. Then, we can compute the discriminant via the discriminant of lattices

$$\mathrm{Disc}(Q) = \mathrm{Disc}(R_Q) = \det(M)^2 \mathrm{Disc}(\widetilde{R_Q}), \quad \mathrm{Disc}(\widetilde{R_Q}) \in \mathcal{O}'_v.$$

Since $\mathrm{val} \mathrm{Disc}(Q) \leq 1$, we are forced to have $\mathrm{val} \det M = 0$, so that $\widetilde{R_Q} = R_Q$ is normal. Therefore, R_Q is regular since it is a reduced, normal, one-dimensional ring. $\square$

We now complete the proof of Proposition 6.22. Since $R_a(\mathcal{O}'_v)$ is regular, it is a product of DVR's $\prod_{i=1}^{2n} R_i$, each of which is finite over $\mathcal{O}'_v$. Let ϖ_i denote a uniformizer in R_i for each i . As a consequence, any $R_a(\mathcal{O}'_v)$ lattice Λ in $R_a(F'_v)$ must be of the form $\Lambda = \langle \ell_1 \varpi_1, \dots, \ell_n \varpi_n \rangle$. Hence, $\mathcal{M}_v^*(a) \otimes_k \bar{k}$ embeds in a discrete space and $\dim \mathcal{M}_v^*(a) = 0$. $\square$

6.6. The product formula. We will now build a bridge between the local models (RASFs) and global moduli spaces (relative Hitchin fibration) by means of a product formula (Proposition 6.24). The proofs of relevant statements are similar to the case of [Ngo10, §4.15].

Let $a \in \mathcal{A}^\heartsuit(k)$, and for any place $v \in |\Sigma|$, denote by a_v the restriction of the map $a: \Sigma \rightarrow \mathfrak{c}_D$ to the infinitesimal disk $\Sigma_v \subset \Sigma$. It is easy to see that $a_v \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$.

Let $U_a = a^{-1}(\mathfrak{c}_D^{\text{rs}})$ denote the preimage of the regular semisimple locus in Σ . We denote by $\check{\Sigma} = \Sigma \otimes_k \bar{k}$ the base change to the algebraic closure $\bar{k}$. Recall that we denote by $\mathcal{M}_a^* = f^{\star,-1}(a)$ the fiber of the relative Hitchin map at a . We will first describe a gluing map

$$(6.8) \quad \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{M}_v^*(a_v) \rightarrow \mathcal{M}_a^*.$$

Let $\tilde{U}'_a \subset \tilde{\Sigma}'_a$ be the preimage of U_a . First, note that the cover $\tilde{U}'_a \rightarrow U_a$ is smooth and étale, so that $\omega_{\tilde{U}'_a/U_a} = \mathcal{O}_{\tilde{U}'_a}$ and the canonical global section $x \in \Gamma(\tilde{U}'_a, \mathcal{O}_{\tilde{U}'_a})$ induced by the element $x \in k[\mathfrak{s}]$ is invertible. Note also that $\iota^*x = -x$. Using the description from Corollary (4.9), we obtain a canonical X^* -Higgs bundle on U_a with characteristic data $a|_{U_a}$ by considering the structure sheaf $\mathcal{O}_{\tilde{U}'_a}$ together with the identity map

$$\alpha = \text{id}: \mathcal{O}_{\tilde{U}'_a} \rightarrow \iota^* \mathcal{O}_{\tilde{U}'_a} \simeq \mathcal{O}_{\tilde{U}'_a}$$

and the Hermitian form

$$h^*: \mathcal{O}_{\tilde{U}'_a} \rightarrow \sigma^* \mathcal{O}_{\tilde{U}'_a} \simeq \mathcal{O}_{\tilde{U}'_a}$$

where $h^{\mathbf{U}} = \text{id}$ and $h^{\mathbf{S}} = x$ is multiplication by the section x . We call the triple $(\mathcal{O}_{\tilde{U}'_a}, h^*, \alpha)$ the trivial X^* -bundle.

The data of an $R_a(\mathcal{O}'_v)$ -lattice in $R_a(F'_v)$ gives gluing data along the punctured disk for a trivial rank $2n$ vector bundle on Σ'_v . The assumptions on self duality and ι fixed imply that we can glue the Hermitian forms h^* on Σ'_v with h^* on U'_a compatibly with the isomorphisms α . By the formal gluing of Beauville and Laszlo [BL95], we may glue such lattices with the trivial X^* -bundle and therefore obtain the map (6.8).

Likewise, we have a map $\prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J_a) \rightarrow \mathcal{P}_a$, and the map (6.8) is compatible with the actions of these symmetry groups. We deduce a map

$$(6.9) \quad \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{M}_v^*(a_v) \wedge \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J_a) \rightarrow \mathcal{M}_a^*$$

We prove the following close analogue of [Ngo10, Prop 4.15.1], [Ngo06, Thm 4.6].

Proposition 6.24. *Let $a \in \mathcal{A}^{\text{ani}}(\bar{k})$. Then the quotient of $\prod_{v \in |\check{\Sigma} - U_a|} \mathcal{M}_v^*(a) \times \mathcal{P}_a$ by $\prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J_a)$ is a proper Deligne–Mumford stack. Moreover, the map (6.9) is a universal homeomorphism; namely, it induces a bijection of groupoids of $\bar{k}$ -points.*

If $a \in \mathcal{A}^{\text{ani}}(K)$ for an extension K/k , the bijection is compatible with the respective Galois actions by $\text{Gal}(k/K)$. In particular, we have an equivalence of groupoids

$$\prod_{v \in |\Sigma_K - U_a|} [\mathcal{M}_v^*(a)/\mathcal{P}_v(J_a)](K_v) \xrightarrow{\sim} [\mathcal{M}_a^*/\mathcal{P}_a](K),$$

for every K/k , with $\Sigma_K := \Sigma \otimes_k K$ and K_v the residue field of Σ_K at the place v .

Proof. To show that the space $\prod_{v \in |\check{\Sigma} - U_a|} \mathcal{M}_v^*(a) \wedge \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J_a) \rightarrow \mathcal{P}_a$ is a proper Deligne–Mumford stack, we can argue identically to [Ngo10, Lemme 4.15.1] using the fact that the lattice Γ_a acts freely on $\mathcal{M}_v(a)$ with quotient $\mathcal{M}_v(a)/\Gamma_a$ a projective, finite type scheme; see Proposition 6.16. Alternatively, one can deduce this from the morphism $\mathcal{M}_v^*(a) \rightarrow \mathcal{N}_v(a)$ using [Rom05, Theorem 3.3].

By considering the quotient by $\mathcal{P}_a$ on each side of (6.9), it suffices to show the following map is an equivalence of groupoids

$$(6.10) \quad \beta: \prod_{v \in |\check{\Sigma} - U_a|} [\mathcal{M}_v^*(a)/\mathcal{P}_v(J_a)](\bar{k}) \rightarrow [\mathcal{M}_a^*/\mathcal{P}_a](\bar{k}),$$

compatible with the Galois action. First, we prove β is fully faithful. If $\Lambda = \{\Lambda_v\}_v$ and $\Lambda' = \{\Lambda'_v\}_v$ are two $\bar{k}$ -points in the left hand side such that $\beta(\Lambda)$ and $\beta(\Lambda')$ are isomorphic, then there exists $p \in \mathcal{P}_a(\bar{k})$ so that $p \cdot \beta(\Lambda) = \beta(\Lambda')$. Since $\beta(\Lambda)$ and $\beta(\Lambda')$ are glued from the trivial X^* -Higgs bundle over U_a , the element p restricts to the identity over U_a , and therefore, can be obtained by gluing of $p_v \in \mathcal{P}_v(J_a)$ for $v \in |\check{\Sigma} - U_a|$. Then, $p_v \cdot \Lambda_v = \Lambda'_v$ for all places v , and we conclude that β is fully faithful.

We now prove that β is essentially surjective. Let m_0 denote the trivial X^* -Higgs bundle over U_a , and let $m \in \mathcal{M}_a^*(\bar{k})$. Since $m|_{U_a}$ is regular semisimple, there exists a J_a -torsor p' on U_a such that $m|_{U_0} = p' \cdot m_0$. Since $H^0(\bar{k}, H^*) = 0$, p' extends to a J_a -torsor on Σ , i.e. $p \in \mathcal{P}_a(\bar{k})$. Then, $p \cdot m$ carries a trivialization identifying $p \cdot m|_{U_a} \simeq m_0$. For $v \in |\check{\Sigma} - U_a|$, the restriction of $p \cdot m$ to the infinitesimal disk Σ_v comes equipped with gluing data over the punctured disk $\Sigma_v^\bullet$, which by the moduli description of $\mathcal{M}_v^*(a)$ is a point $m_v \in \mathcal{M}_v^*(a)(\bar{k})$. The element p gives an explicit isomorphism between $\beta((m_v))$ and m , proving surjectivity.

Finally, we note that the gluing map (6.8) intertwines the Galois actions. Indeed, the Frobenius σ acts on $\mathcal{M}_a^*$ through $\text{Aut}(\Sigma'/\Sigma)$, which restricts to actions of the Galois group $\text{Gal}(F'_v/F_v)$ for every $v \in |\check{\Sigma}|$. $\square$

Remark 6.25. Let $a \in \mathcal{A}^{\text{ani}}(\bar{k})$ be as above. Suppose that J'_a is a smooth group scheme equipped with a map $J'_a \rightarrow J_a$, which is an isomorphism on $U_a \subset \check{\Sigma}$. Then $\mathcal{P}_v(J'_a)$ acts on $\mathcal{M}_v^*(a)$ through $\mathcal{P}_v(J_a)$, and moreover, we may form a Picard stack $\mathcal{P}'_a$ of J'_a torsors on Σ which acts on $\mathcal{M}_a^*$. We claim that Proposition 6.24 holds replacing $\mathcal{P}_a$ with $\mathcal{P}'_a$ and $\mathcal{P}_v(J_a)$ with $\mathcal{P}_v(J'_a)$. Indeed, by a Beauville-Laszlo gluing argument, the morphism

$$\mathcal{P}'_a \rightarrow \mathcal{P}_a$$

is a surjective map with kernel a torsor under the group $\prod_{v \in |\check{\Sigma} - U_a|} J_{a,v}(\mathcal{O}_v)/J'_{a,v}(\mathcal{O}_v)$. In particular,

$$\mathcal{P}_a \simeq \mathcal{P}'_a \wedge \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J'_a) \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J_a).$$

Therefore, the map $J'_a \rightarrow J_a$ induces an isomorphism

$$\begin{aligned} \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{M}_v^*(a) \wedge \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J'_a) \mathcal{P}'_a &\simeq \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{M}_v^*(a) \wedge \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J_a) \left(\prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J_a) \wedge \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J'_a) \mathcal{P}'_a \right) \\ &\simeq \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{M}_v^*(a) \wedge \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J_a) \mathcal{P}_a. \end{aligned}$$

Therefore, Proposition 6.24 gives a morphism

$$\prod_{v \in |\check{\Sigma} - U_a|} \mathcal{M}_v^*(a_v) \wedge \prod_{v \in |\check{\Sigma} - U_a|} \mathcal{P}_v(J'_a) \mathcal{P}'_a \xrightarrow{\sim} \mathcal{M}_a^*,$$

which is a universal homeomorphism over k .

6.7. Local Constancy of RASFs. In this section, we will prove the following proposition, mirroring [Ngo10, Proposition 3.5.1] in our relative context. The proof here is simpler due to our use of spectral covers.

Proposition 6.26. *Suppose $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$ satisfies $\mathrm{Ob}_v^*(a) = 0$, and let $\mathcal{M}_v^*(a)$ (for $\star = \mathbf{S}$ or $\mathbf{U}$) denote the associated affine Springer fibers and $\mathcal{P}_v(J_a)$ the group of symmetries. There exists an integer $N > 0$ such that for all finite extensions K of k and for all $a' \in \mathfrak{c}^\heartsuit(\mathcal{O}_v \otimes_k K)$ satisfying*

$$a \equiv a' \pmod{\varpi_v^N},$$

the affine Springer fiber $\mathcal{M}_v^(a')$ with the action of $\mathcal{P}_v(J_{a'})$ is isomorphic to $\mathcal{M}_v^*(a) \otimes_k K$ with the action of $\mathcal{P}_v(J_a) \otimes_k K$.*

We introduce some preliminaries before the proof below. Let $\tilde{\Sigma}'_{a,v} = \mathrm{Spec} R_a(\mathcal{O}_{F'_v})$, where $R_a(\mathcal{O}_{F'_v})$ is as in §6.1. This cover $\tilde{\Sigma}'_{a,v}$ fits into a Cartesian diagram

$$\begin{array}{ccc} \tilde{\Sigma}'_{a,v} & \longrightarrow & \mathfrak{s} \\ \downarrow & & \downarrow \\ \Sigma'_v & \longrightarrow & \mathfrak{c} \end{array}$$

Recall from §6.1 that $\tilde{\Sigma}'_{a,v}$ carries two involutions: σ acting by the nontrivial involution in $\mathrm{Aut}(\Sigma'/\Sigma)$ and τ the involution of the spectral cover induced by $x \mapsto -x$ (see §4.4 for a discussion).

Lemma 6.27. *For any $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$ and integer N_1 , there exists an integer $N > N_1$ such that for all finite extensions K of k and all $a' \in \mathfrak{c}(\mathcal{O}_v \otimes_k K)$ satisfying*

$$a \equiv a' \pmod{\varpi_v^N},$$

there is a σ - and τ -equivariant isomorphism $\tilde{\Sigma}'_{a',v} \simeq \tilde{\Sigma}'_{a,v} \otimes_k K$ over $\Sigma'_v \otimes_k K$ which lifts the obvious one mod $\varpi_v^{N_1}$.

Proof. This is a consequence of [Art69, Lemma 3.12] and is very close to the statement recorded in [Ngo10, Lemma 3.5.2]. $\square$

Proof of Proposition 6.26. Fix $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$. Let N be as in Lemma 6.27 for $N_1 = 1$. Without loss of generality, we assume that $k = K$, and suppose that $a' \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$ is such that $a' \equiv a \pmod{\varpi_v^N}$. Then, by Lemma 6.27, $R_a(\mathcal{O}_{F'_v}) \simeq R_{a'}(\mathcal{O}_{F'_v})$ compatibly with the involutions σ_R and ι_R . Compatibility with σ_R implies in particular that the isomorphism respects the Hermitian forms h^* on each side, and so from the lattice definition (see §6.1), we obtain an isomorphism $\mathcal{M}_v(a) \simeq \mathcal{M}_v(a')$ $\square$

6.8. Dimension formula. In [Bez96], it was shown that regular locus in (ordinary) affine Springer fibers is dense, and hence one can deduce a formula for the dimension of an affine Springer fiber from the dimension of its group of symmetries. This method fails for RASF's as regularity is no longer a dense condition. It also fails in the multiplicative setting of [Chi22, Wan26] for a similar reason: multiplicative affine Springer fibers have rich component groups, related to Mirković–Vilonen cycles, and in particular, have no dense regular locus. In work of Jingren Chi [Chi22], a dimension formula and equidimensionality property in the multiplicative case is proved by passing to the global setting. We follow an analogous argument, with some simplifications. In particular, for any $a_v \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$, we construct a curve in the Hitchin base which interpolates between a_v and a split element, and use upper semi-continuity of fiber dimensions for proper maps to deduce the dimension bound. We also collect as a corollary that the relative Hitchin morphism is flat over the anisotropic locus, from which equidimensionality follows.

Let Σ be a smooth projective curve of genus $g \geq 2$ and $v \in |\Sigma|$. Recall from Section 6.1 $\mathfrak{c}^\heartsuit(\mathcal{O}_v) = \mathfrak{c}(\mathcal{O}_v) \cap \mathfrak{c}^{\mathrm{rs}}(F_v)$. As we will be exchanging between the local and global settings in this

section, we will denote by a_v a fixed element in the local base $\mathfrak{c}^\heartsuit(\mathcal{O}_v)$ and by a as element in the global Hitchin base $\mathcal{A} = \mathcal{A}_D$.

Associated to any $a_v \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$, we have two local invariants mimicking the invariants that control the dimension of affine Springer fibers in [Bez96]. First, we put

$$(6.11) \quad d_v^{\text{cent}}(a_v) := \deg(a_v^* \mathfrak{D}^{\text{cent}})$$

to be the multiplicity of intersection of a_v at the centralizer divisor. We also have the invariant $c_v(a_v)$ controlling how split a_v is (see Definition 6.18): define

$$c_v(a_v) = n - \text{rank } \Gamma_a,$$

where $\Gamma_a = \mathbb{X}_*(J_{a_v, \bar{F}_v})$ is the lattice of $\check{F}_v$ -rational cocharacters of the torus $J_{a_v, \check{F}_v}$ over $\check{F}_v$ acting on the RASF as defined in §6.4. Note in particular that $c_v(a_v) = 0$ if and only if a_v is split.

We begin by proving the following dimension formula.

Theorem 6.28. *Let $a_v \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$. The RASF $\mathcal{M}_v^*(a_v)$ has dimension equal to that of $\mathcal{P}_v(J_a)$. Explicitly, we have*

$$\dim \mathcal{M}_v(a_v) = \dim \mathcal{P}_v(J_{a_v}) = \frac{d_v^{\text{cent}}(a) - c_v(a)}{2}$$

Proof. Our proof follows [Chi22, §5.2], with some simplifications in our case. As dimension is a geometric property, we will base change to an algebraic closure $\bar{k}$ throughout the rest of this section.

First, we note that, there exists a point $m \in \mathcal{M}_v^{\text{reg}}(a_v)(\bar{k})$ given by the image of a Kostant-Rallis section. The stabilizer of m in $\mathcal{P}_v(J_{a_v})$ is trivial since m is regular, so we have $\dim \mathcal{M}_v(a_v) \geq \dim \mathcal{P}_v(J_{a_v})$. The rest of this proof will concern the opposite inequality.

Recall the algebra of functions $R = R_a(\mathcal{O}_v)$ on $\Sigma_{a_v, v} = \Sigma_v \times_{\mathfrak{c}} \mathfrak{s}$ considered in §6.1, where we recall that $\mathfrak{s}$ is the spectral cover appearing in §2.1. Then $R_a(F_v) = R \otimes_{\mathcal{O}_v} F_v$ splits as a product of extensions $\prod_i E_i$ of F_v of degrees e_i . Put $e = \text{LCM}(e_i)$ to be the least common multiple of the degrees e_i ; in particular, we have $e = 1$ if and only if a_v is split. Consider the family a_t given by viewing a as n -tuple of power series in ϖ_v and substituting $\varpi_v \mapsto t\varpi_v + (1-t)\varpi_v^e$ in a_v , i.e.

$$a_t = a_v(t\varpi_v + (1-t)\varpi_v^e) \in \mathfrak{c}^\heartsuit(\mathcal{O}_v).$$

We note that $\mathcal{M}_v^*(a_t) \simeq \mathcal{M}_v^*(a_1) = \mathcal{M}_v^*(a_v)$ for $t \neq 0$ and that $a_0 = a_v(\varpi_v^e)$ is split since $\varpi_v \mapsto \varpi_v^e$ splits $R_a(F_v)$. Let $N > 0$ be as in Proposition 6.26 for both a_v and a_0 , so that $\mathcal{M}_v^*(a'_v) \simeq \mathcal{M}_v^*(a_v)$ whenever $a'_v \equiv a_v \pmod{\varpi_v^N}$ and likewise for a_0 . Fix a smooth, projective curve Σ of genus $g \geq 2$ and an étale double cover $\vartheta: \Sigma' \rightarrow \Sigma$ so that at a distinguished place v , the extension of Σ'_v by Σ_v is the extension $\mathcal{O}_v \subset \mathcal{O}'_v$. Choose a line bundle D on Σ such that

$$\deg(D) > \max\{N \cdot n, 2g\}.$$

where $n = \dim \mathfrak{a}$ is the rank of the symmetric variety X^* . Let $\mathcal{M}^{*, \text{ani}} = \mathcal{M}_{\Sigma, D}^{*, \text{ani}}$ be the relative Hitchin moduli space, which is smooth by Proposition 5.8. For every $u \neq v$ in $|\Sigma|$, we have an affine linear map

$$(6.12) \quad \mathcal{A} = \mathcal{A}_D \rightarrow \mathfrak{c}(\mathcal{O}_v/\varpi_v^N) \times \mathfrak{c}(\mathcal{O}_u/\varpi_u^2),$$

which is surjective by Riemann-Roch since $\deg(D) \geq \max\{N \cdot n, 2g\} - 1 \geq \frac{1}{2}(2g + N + 1)$. Let C denote the curve in $H^0(\Sigma, \mathfrak{c}(\mathcal{O}_v/\varpi_v^N))$ given by the family a_t . Define the closed subset $L_C \subset \mathcal{A}$ to be the preimage of C , and let Z denote the set of points a such that $a|_{\Sigma - \{v\}}$ meets the discriminant divisor $\mathfrak{D}_X$ transversely. Put $Z_C = Z \cap L_C$. The set Z is open in $\mathcal{A}^\heartsuit$ and intersects each fiber of the map $L_C \rightarrow C$ non-trivially by surjectivity of (6.12). Moreover, since by Lemma 5.9, the codimension of $\mathcal{A} - \mathcal{A}^{\text{ani}}$ in $\mathcal{A}$ is greater than $\deg(D)$ and Z_C is open in L_C , which is of codimension $N \cdot n$ in $\mathcal{A}$, it follows that $Z_C^{\text{ani}} = Z_C \cap \mathcal{A}^{\text{ani}}$ is also nonempty and open in L_C .

Composing (6.12) with projection onto the first factor gives

$$(6.13) \quad \mathcal{A} = \mathcal{A}_D \rightarrow \mathfrak{c}(\mathcal{O}_v/\varpi_v^N),$$

which is also an affine linear map between affine spaces. Choose a section s of the map (6.13) such that $s(a_0) \in Z_C^{\text{ani}}$. As $Z_C^{\text{ani}} \subset L_C$ is open, this implies that $s^{-1}(Z_C^{\text{ani}})$ is open in C . Therefore, we may choose a fixed $t \neq 0$ such that $s(a_t) \in Z_C^{\text{ani}}$.

As $f^{\star, \text{ani}}$ is proper (Proposition 5.8), by upper semi-continuity of fiber dimensions we have

$$\dim \mathcal{M}_{s(a_t)}^{\star} \leq \dim \mathcal{M}_{s(a_0)}^{\star}$$

By Proposition 6.20, we know that $\dim \mathcal{M}_{s(a_0)}^{\star} = \dim \mathcal{P}_{s(a_0)}$, which by smoothness of the group scheme $\mathcal{P}$ is equal to $\dim \mathcal{P}_{s(a_t)}$. Using the product formula 6.24, we deduce that

$$\begin{aligned} \dim \mathcal{M}_{s(a_t)}^{\star} - \dim \mathcal{P}_{s(a_t)} &= \sum_{v' \in |\check{\Sigma} - (U_a \cup \{v\})|} (\dim \mathcal{M}_{v'}^{\star}(a) - \dim \mathcal{P}_{v'}(J_a)) + (\dim \mathcal{M}_v^{\star}(a) - \dim \mathcal{P}_v(J_a)) \\ &= \dim \mathcal{M}_v^{\star}(a) - \dim \mathcal{P}_v(J_a) \end{aligned}$$

where the second equality follows from Proposition 6.22. Hence, we conclude that

$$\dim \mathcal{M}_v^{\star}(a) - \dim \mathcal{P}_v(J_a) = \dim \mathcal{M}_{s(a_t)}^{\star} - \dim \mathcal{P}_{s(a_t)} \leq \dim \mathcal{M}_{s(a_0)}^{\star} - \dim \mathcal{P}_{s(a_0)} = 0$$

so that $\dim \mathcal{M}_v^{\star}(a) = \dim \mathcal{P}_v(J_a)$, as desired. The final formula now follows from Bezrukavnikov's formula computing the dimension of $\mathcal{P}_v(J_a)$ [Bez96]. $\square$

Corollary 6.29. *The relative Hitchin fibration restricted to anisotropic locus $f^{\star, \text{ani}}: \mathcal{M}^{\star, \text{ani}} \rightarrow \mathcal{A}^{\text{ani}}$ is flat.*

Proof. Since $\mathcal{M}^{\star, \text{ani}}$ is smooth by Proposition 5.8, by Miracle Flatness it suffices to show the fibers of $f^{\star, \text{ani}}$ are equidimensional of the same dimension. Since by Proposition 5.8 $f^{\star, \text{ani}}$ is proper, the fiber dimension is upper semicontinuous on the source. Hence, it suffices to show the fiber dimensions are equal over any $a \in \mathcal{A}^{\text{ani}}$. For any $a \in \mathcal{A}^{\text{ani}}$, we have by product formula (Proposition 6.24) and the dimension formula for RASFs (Theorem 6.28) that

$$\dim[\mathcal{M}_a^{\star}/\mathcal{P}_a] = \sum_{v \in |\check{\Sigma} - U_a|} \dim[\mathcal{M}_v^{\star}(a)/\mathcal{P}_v(J_a)] = 0.$$

Hence, $\dim \mathcal{M}_a^{\star} = \dim \mathcal{P}_a$ for all $a \in \mathcal{A}^{\text{ani}}$. Moreover, by uppersemicontinuity of local dimension of fibers, this constrains all components in $\mathcal{M}_a^{\star}$ to also be of dimension $\dim \mathcal{P}_a$. Finally, since $\mathcal{P}$ is smooth over $\mathcal{A}^{\text{ani}}$, we conclude that $\dim \mathcal{M}_a^{\star}$ is constant on $\mathcal{A}^{\text{ani}}$. $\square$

Proposition 6.30. *Let $a_v \in \mathfrak{c}^{\heartsuit}(\mathcal{O}_v)$. The RASF $\mathcal{M}_v^{\star}(a_v)$ is equidimensional.*

Proof. Let $N > 0$ be as in Proposition 6.26. Choose a line bundle D on Σ such that

$$\deg(D) > \max\{N \cdot n, 2g\}.$$

Let $\mathcal{M}^{\star, \text{ani}} = \mathcal{M}_{\Sigma, D}^{\star, \text{ani}}$ be the relative Hitchin moduli space, which is smooth by Proposition 5.8.

Let $L \subset \mathcal{A}$ be the subset of points a such that

$$a \equiv a_v \pmod{\varpi_v^N}.$$

In particular, L is a closed subset of $\mathcal{A}^{\heartsuit}$ since $a_v \in \mathfrak{c}^{\heartsuit}(\mathcal{O}_v) \subset \mathfrak{c}^{\text{rs}}(F_v)$. Alternatively, we note that L is the preimage of a_v under the local evaluation map

$$\mathcal{A} \rightarrow \mathfrak{c}(\mathcal{O}_v/\varpi_v^N),$$

which is surjective using Riemann–Roch by the assumption on $\deg(D)$. Therefore, the codimension of L in $\mathcal{A}$ is given by

$$\text{codim}_{\mathcal{A}}(L) = N \cdot n.$$

The complement of the anisotropic locus has codimension at least $\deg(D) > N \cdot n$ by Lemma 5.9. Hence, L intersects non-trivially with the open set $\mathcal{A}^{\text{ani}}$, and the set $L^{\text{ani}} = L \cap \mathcal{A}^{\text{ani}}$ is locally closed

in $\mathcal{A}$. We claim that L^{ani} contains a point a such that for any $v' \in |\Sigma| \setminus \{v\}$, the image of a under the local evaluation map

$$(6.14) \quad \mathcal{A} = \mathcal{A}_{\Sigma, D} \rightarrow \mathfrak{c}(\mathcal{O}_{v'}/\varpi_{v'}^2)$$

is nonzero. That is, $a|_{\Sigma \setminus \{v\}}$ meets the discriminant divisor transversely away from v and we have

$$a \equiv a_v \pmod{\varpi_v^N}.$$

In this case, by the product formula (Proposition 6.24), we have

$$[\mathcal{M}_a^*/\mathcal{P}_a] \underset{\text{Homeo}}{\simeq} [\mathcal{M}_v^*(a)/\mathcal{P}_v(J_a)] \times \prod_{v' \in |\check{\Sigma} - (U_a \cup \{v\})|} [\mathcal{M}_{v'}^*(a)/\mathcal{P}_{v'}(J_a)],$$

where, for any $v' \in |\check{\Sigma} - (U_a \cup \{v\})|$, we have $[\mathcal{M}_{v'}^*(a)/\mathcal{P}_{v'}(J_a)]$ is equidimensional of dimension zero by Proposition 6.22. Since $f^{\star, \text{ani}}$ is flat (Corollary 6.29), $[\mathcal{M}_a^*/\mathcal{P}_a]$ is equidimensional (of dimension zero), and so $[\mathcal{M}_v^*(a)/\mathcal{P}_v(J_a)]$ must also be equidimensional. The result then follows.

To prove the existence of such a point a , we argue similarly to [Ngo10, Proposition 4.7.1] and [Chi22, Lemma 5.2.6]. By our assumptions on the degree of D , the evaluation map

$$(6.15) \quad L \rightarrow \mathfrak{c}(\mathcal{O}_{v'}/\varpi_{v'}^2)$$

is surjective. Let $\mathfrak{D}^{\text{cent}} = \mathfrak{D}^{\text{sm}} \cup \mathfrak{D}^{\text{sing}}$ where $\mathfrak{D}^{\text{sm}}$ is the smooth locus of $\mathfrak{D}^{\text{cent}}$ and $\mathfrak{D}^{\text{sing}}$ its complement (the intersection of root hyperplanes). Then, define

$$Z_1 = \{(c, a) \in \mathfrak{D}_D^{\text{sm}} \times \mathcal{A} \mid a \text{ meets } \mathfrak{D}_D \text{ at } c \text{ with multiplicity } \geq 2\}$$

$$Z_2 = \{(c, a) \in \mathfrak{D}_D^{\text{sing}} \times L \mid a \text{ meets } \mathfrak{D}_D \text{ at } c\}$$

For each of the Z_j , the surjectivity of (6.15) implies that

$$\dim(Z_j) \leq \dim(L) - 1$$

Hence, the complement of the image of the projection $Z_1 \cup Z_2 \rightarrow L$ is a nonempty open set in L . $\square$

Finally, we collect as a corollary a relationship with dimensions of endoscopic RASF's. Recall from Lemma 2.21 that there exists a divisor $\mathfrak{R}_{\mathfrak{c}} \subset \mathfrak{c}_{\mathfrak{c}}$ so that $\nu_{\mathfrak{c}}^* \mathfrak{D}^{\text{cent}} = \mathfrak{D}_{\mathfrak{c}}^{\text{cent}} + 2\mathfrak{R}_{\mathfrak{c}}$. Define $r_v(a_{\mathfrak{c}}) = \text{val}(a_{\mathfrak{c}}^* \mathfrak{R}_{\mathfrak{c}})$ for $a_{\mathfrak{c}} \in \mathfrak{c}_{\mathfrak{c}}^{\heartsuit}(\mathcal{O}_v)$. We will denote by $\delta(a)$ the dimension of the RASF $\mathcal{M}_v^*(a)$, which is independent of $\star \in \{\mathbf{S}, \mathbf{U}\}$.

Lemma 6.31. *For $a_{\mathfrak{c}} \in \mathfrak{c}_{\mathfrak{c}}^{\heartsuit}(\mathcal{O}_v)$ and $a \in \mathfrak{c}^{\heartsuit}(\mathcal{O}_v)$ such that $\nu_{\mathfrak{c}}(a_{\mathfrak{c}}) = a$, we have*

$$\delta_v(a) = \delta_{\mathfrak{c}, v}(a_{\mathfrak{c}}) + r_v(a_{\mathfrak{c}})$$

Proof. Note that $c_v(a)$ and $c_{\mathfrak{c}, v}(a_{\mathfrak{c}})$ depend only on the generic fiber of J_a , respectively $J_{a_{\mathfrak{c}}}$. By Proposition 2.14, J_a and $J_{a_{\mathfrak{c}}}$ have isomorphic generic fibers, and so $c_v(a) = c_{\mathfrak{c}, v}(a_{\mathfrak{c}})$. On the other hand, by Lemma 2.21, $d_v^{\text{cent}}(a) = d_{\mathfrak{c}, v}^{\text{cent}}(a_{\mathfrak{c}}) + 2r_v(a_{\mathfrak{c}})$. Hence, by the dimension formula, Proposition 6.28, we have

$$\delta_v(a) = \frac{d_v^{\text{cent}}(a) - c_v(a)}{2} = \frac{d_{\mathfrak{c}, v}^{\text{cent}}(a_{\mathfrak{c}}) - c_{\mathfrak{c}, v}(a_{\mathfrak{c}})}{2} + r_v(a_{\mathfrak{c}}) = \delta_{\mathfrak{c}, v}(a_{\mathfrak{c}}) + r_v(a_{\mathfrak{c}}) \quad \square$$

Lemma 6.31 implies a global result as well. We fix global data $\vartheta: \Sigma' \rightarrow \Sigma$ and D on Σ as in §4. The divisor $\mathfrak{R}_{\mathfrak{c}}$ is homogeneous for the $\mathbf{G}_m$ action on $\mathfrak{c}_{\mathfrak{c}}$, and so we can form the divisor $(\mathfrak{R}_{\mathfrak{c}})_D$ in $\mathfrak{c}_{\mathfrak{c}, D}$. Put $r_v(D) = \deg a_{\mathfrak{c}}^*(\mathfrak{R}_{\mathfrak{c}})_D$. When $X^{\mathfrak{c}} = X_r^{\mathbf{S}} \times X_s^{\mathbf{U}}$ for $r + s = n$, we get $r_v(D) = 2rs \deg(D)$.

Corollary 6.32. *Fix a divisor D on Σ , and let $\mathfrak{c} = (\kappa, \rho)$ be an endoscopic datum. For any $a_{\mathfrak{c}} \in \mathcal{A}_{\mathfrak{c}}^{\text{ani}}(\bar{k})$ and $a \in \mathcal{A}^{\text{ani}}(\bar{k})$ such that $\nu_{\mathfrak{c}}(a_{\mathfrak{c}}) = a$, we have*

$$\delta(a) = \delta_{\mathfrak{c}}(a_{\mathfrak{c}}) + r_v(D)$$

Proof. By the product formula Proposition 6.24, it suffices to note that

$$r_v(D) = \sum_{v \in |\check{\Sigma} - U_{a_\epsilon}|} \deg(v) r_v(a_\epsilon). \quad \square$$

6.9. Density of $\mathcal{A}^\dagger$ in δ -strata. We now extract some global corollaries from these local results. Recall the δ -stratification of $\mathcal{A}$ in Section 5.5, as well as the $\dagger$ -locus $\mathcal{A}^\dagger$ in Definition 5.11. We prove the following density result.

Proposition 6.33. *Fix a positive integer δ , and suppose that $\deg(D) \geq g + \frac{3}{2}\delta$. Then the locus $\mathcal{A}_\delta^\dagger := \mathcal{A}^\dagger \cap \mathcal{A}_\delta$ is open and dense in every component of $\mathcal{A}_\delta$.*

The proof of this proposition involves some results in deformation theory. We begin with the following claim.

Lemma 6.34. *Fix distinct points $x_0, x_1, \dots, x_r, y \in \check{\Sigma}(\bar{k})$. If $\deg(D) > g + \frac{3}{2}r$, then the product of local evaluation maps*

$$(6.16) \quad \mathcal{A} \rightarrow H^0(\Sigma_y, \mathfrak{c}(\mathcal{O}_y/\varpi_y)) \oplus H^0(\Sigma_{x_0}, \mathfrak{c}(\mathcal{O}_{x_0}/\varpi_{x_0})) \oplus \prod_{i=1}^r H^0(\Sigma_{x_i}, \mathfrak{c}(\mathcal{O}_{x_i}/\varpi_{x_i}^3))$$

is surjective.

Proof. This morphism has cokernel given by

$$H^1\left(\Sigma, \mathfrak{c}_D\left(-y - x_0 - 3 \sum_{i=1}^r x_i\right)\right)$$

Since $\mathfrak{c}_D = \bigoplus_i D^{\otimes 2i}$, this cohomology group vanishes by Riemann–Roch when

$$2 \deg(D) - 3r - 2 > 2g - 2$$

which can be rearranged to read

$$\deg(D) > g + \frac{3}{2}r.$$

Starting with this assumption, we see that (6.16) is a surjection. $\square$

We will use a local description of δ invariants.

Definition 6.35. For $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$, we denote by $\delta_v(a) = \dim \mathcal{M}_v^*(a)$ the dimension of the corresponding RASF at a .

Note that Proposition 6.28 gives a formula computing $\delta_v(a)$. Moreover, by the product formula (Proposition 6.24), for any $a \in \mathcal{A}^{\text{ani}}(\bar{k})$ we have an equality

$$\delta(a) = \sum_{v \in |\check{\Sigma}|} \delta_v(a),$$

where only finitely many terms in the right hand side are nonzero.

Let $\mathcal{A}_{(1^\delta)} \subset \mathcal{A}_\delta$ denote the locus of points a in $\mathcal{A}_\delta$ such that $\delta_x(a) \leq 1$ for all $x \in \check{\Sigma}(\bar{k})$. This subset was used in [Ngo10] alongside other locally closed sets subsets $\mathcal{A}_\lambda$ for partitions $\lambda \vdash \delta$ to give a stratification of $\mathcal{A}_\delta$. In particular, Ngô shows, in the proof of [Ngo10, Proposition 5.7.2] that:

Lemma 6.36. *The subset $\mathcal{A}_{(1^\delta)}$ is open and dense in $\mathcal{A}_\delta$.*

We will need the following refinement of $\mathcal{A}_{(1^\delta)}$. Let $\mathcal{A}_{(1^\delta)}^{\min}$ be the set of points a in $\mathcal{A}_{(1^\delta)}$ such that, for all x such that $\delta_x(a) = 1$, we also have $d_x^{\text{cent}}(a) = 2$. Note that by the dimension formula Proposition 6.28, we must have $d_x^{\text{cent}}(a) \geq 2\delta_x(a)$ for all x , so this subset should be thought of as points of $\mathcal{A}_\delta$ which meet the divisor $\mathfrak{D}_D^{\text{cent}}$ “as transversely as possible.”

Lemma 6.37. *Treat $H^0(\Sigma_v, \mathfrak{c}(\mathcal{O}_v/\varpi_v^3))$ as an affine space. There exists a closed subvariety $\mathfrak{Z} \subset H^0(\Sigma_v, \mathfrak{c}(\mathcal{O}_v/\varpi_v^3))$ and an open, dense subset $\mathfrak{U} \subset \mathfrak{Z}$ such that the condition $\delta_v(a) > 0$ is equivalent to the image of a under the local evaluation map*

$$\mathcal{A} \rightarrow H^0(\Sigma_v, \mathfrak{c}(\mathcal{O}_v/\varpi_v^3))$$

landing in $\mathfrak{Z}$ while the condition that both $\delta_v(a) = 1$ and $d_v^{\text{cent}}(a) = 2$ is equivalent to the image of a landing in $\mathfrak{U}$.

Proof. Since our δ -invariant coincides with the δ -invariant for the local Jacobian of $\bar{\Sigma}_{a,v}$ (see Lemma 5.29), the condition that $\delta_v(a) > 0$ is equivalent to the condition that the germ of the associated spectral curve is singular. This can be described by vanishing of partial derivatives with respect to x and ϖ_v and is hence a closed condition.

For the second claim, note that $d_v^{\text{cent}}(a) = \text{val}_v(a^* \mathfrak{D}_D^{\text{cent}})$ is an upper semicontinuous function, and so the locus with minimal valuation is open in the space where $\delta_v(a) > 0$. In particular, by the dimension formula (Theorem 6.28), $d_v^{\text{cent}}(a) = 2$ is the minimal valuation if we impose $\delta_v(a) > 0$. As having valuation 2 is detectable mod ϖ_v^3 , the lemma follows. $\square$

Fix δ distinct points $\underline{x} = (x_1, \dots, x_\delta)$ on $\check{\Sigma}$. For $\underline{\xi} = (\xi_1, \dots, \xi_\delta)$ with $\xi_i \in H^0(\Sigma, \mathcal{O}_{x_i})$ satisfying both $\delta_{x_i}(a) = 1$ and $d_{x_i}^{\text{cent}}(a) = 2$, we denote by $\mathcal{A}_{\underline{\xi}}$ the preimage of $\underline{\xi}$ under the local evaluation map

$$\mathcal{A} \rightarrow \prod_{i=1}^{\delta} H^0(\Sigma, \mathcal{O}_{x_i}/\varpi_{x_i}^3).$$

Let $\mathcal{A}_{\geq(1^\delta)}$ denote the subvariety of $a \in \mathcal{A}$ for which there exist at least δ distinct points $x_1, \dots, x_\delta$ of $\check{\Sigma}$ such that $\delta_{x_i}(a) \geq 1$. Then, by Lemma 6.37, we have

$$(6.17) \quad \mathcal{A}_{\geq(1^\delta)} = \bigcup_{\underline{\xi} \in \text{ev}^{-1}(\mathfrak{Z})} \mathcal{A}_{\underline{\xi}}, \quad \mathcal{A}_{(1^\delta)}^{\min} = \bigcup_{\underline{\xi} \in \text{ev}^{-1}(\mathfrak{U})} \mathcal{A}_{\underline{\xi}}$$

where $\mathfrak{Z}$ and $\mathfrak{U}$ are as in the statement of the lemma. In particular, we also deduce the following.

Corollary 6.38. *Suppose that $\deg(D) \geq g + \frac{3}{2}\delta$. The subset $\mathcal{A}_{(1^\delta)}^{\min}$ contains an open and dense subset of $\mathcal{A}_{(1^\delta)}$. In particular, it contains an open, dense subset of $\mathcal{A}_\delta$.*

Proof. By Lemma 6.34, the hypothesis on degrees implies that the local evaluation map (6.16) is surjective, and so $\text{ev}^{-1}(\mathfrak{U})$ is open and dense in $\text{ev}^{-1}(\mathfrak{Z})$. The result follows therefore from (6.17). $\square$

Proof of Proposition 6.33. Fix any points $x_1, \dots, x_n$ on $\check{\Sigma}$ which are pairwise distinct. For any fixed x, y (with $x \neq y$) distinct from each of the x_i , by Lemma 6.34 and our assumption on $\deg(D)$, the map

$$\mathcal{A}_{\underline{\xi}} \rightarrow H^0(\Sigma, \mathcal{O}_x/\varpi_x) \oplus H^0(\Sigma, \mathcal{O}_y/\varpi_y)$$

remains surjective. Therefore, if we consider the variety $Z_1 \subset \mathfrak{D}_D^{\text{cent}} \oplus \mathcal{A}_{\underline{\xi}}$ of pairs (c, a) such that $a(\check{\Sigma})$ meets $\mathfrak{D}_D^{\text{cent}}$ at c and $d_c^{\text{cent}}(a) > 1$, we can show that $\dim(Z_1) \leq \dim(\mathcal{A}_{\underline{\xi}}) - 1$, and similarly for $Z_2 \subset \mathfrak{D}_D^{\text{br}} \oplus \mathcal{A}_{\underline{\xi}}$ consisting of pairs (c, a) such that $a(\check{\Sigma})$ meets $\mathfrak{D}_D^{\text{br}}$ at c and $d_c^{\text{br}}(a) > 1$. Since $\mathcal{A}_{\underline{\xi}, \delta}^\dagger := \mathcal{A}_{\underline{\xi}} \cap \mathcal{A}_\delta^\dagger$ contains the complement of the image of $Z_1 \cup Z_2$ in $\mathcal{A}_{\underline{\xi}}$, we deduce that $\mathcal{A}_{\underline{\xi}, \delta}^\dagger$ contains an open, dense subset of $\mathcal{A}_{\underline{\xi}}$. By the decomposition (6.17), it follows that $\mathcal{A}_\delta^\dagger$ is dense in $\mathcal{A}_{(1^\delta)}^{\min}$. The result now follows from Corollary 6.38. $\square$

7. POINT COUNTING

In this section, we connect the geometry of relative affine Springer fibers to the arithmetic of orbital integrals by counting points. In particular, this will permit us to access orbital integrals by studying cohomology of relative affine Springer fibers.

7.1. Orbital Integrals. We begin by recalling the definition of orbital integrals for each of the unitary and Siegel cases. Let $V^\star$ denote the representation of the group $H^\star$ over k , where $\star = \mathbf{S}$ or $\mathbf{U}$ as in §2.2. We may choose Haar measures dh_v on $H^\star(F_v)$, normalized so that $\text{vol}(H^\star(\mathcal{O}_v), dh_v) = 1$, and dt_v on $I_\gamma \subset H^\star(F_v)$. The orbital integral at γ is defined as

$$\text{Orb}^\star(\gamma, dt_v) = \int_{I_\gamma \backslash H^\star(F_v)} \mathbb{1}_{V^\star(\mathcal{O}_v)}(h^{-1} \cdot \gamma) \frac{dh_v}{dt_v}$$

where $\mathbb{1}_{V^\star(\mathcal{O}_v)}$ denotes the indicator function for the integral points $V^\star(\mathcal{O}_v)$.

Recall that an element $\gamma \in V^\star(F_v)$ is *geometrically conjugate* to γ_0 when $\gamma = h \cdot \gamma_0$ for $h \in H(\check{F}_v)$; in this setting, we have the Galois cocycle

$$\text{inv}(\gamma, \gamma_0) \in H^1(F_v, I_{\gamma_0}) = H^1(F_v, J_a),$$

where a is the common image of γ and γ_0 in $\mathfrak{c}(F_v)$, as follows: choose $h \in H(\check{F}_v)$ such that $h \cdot \gamma_0 = \gamma$. Then, for every $\varsigma \in \text{Gal}(\bar{F}_v/F_v)$, the element $h^{-1} \cdot \varsigma(h)$ fixes γ_0 and hence must lie in the centralizer I_{γ_0} . The class $\text{inv}(\gamma, \gamma_0)$ is the cocycle

$$\varsigma \mapsto h^{-1} \cdot \varsigma(h);$$

this is independent of the choice of h since h is well defined up to translation by the centralizer I_{γ_0} . When γ_0 is regular semi-simple (so that I_{γ_0} is abelian), this cocycle is well defined up to multiplication by a 1-coboundary $\varsigma \mapsto x^{-1} \cdot \varsigma(x)$ for $x \in I_{\gamma_0}$.

For $\kappa \in H^1(F_v, I_{\gamma_0})^D$ (where D here refers to the Pontryagin dual for finite abelian groups), we form the κ orbital integral by considering the weighted sum

$$\text{Orb}_a^{\star, \kappa}(dt_v) = \sum_{\gamma} \langle \kappa, \text{inv}(\gamma, \gamma_0) \rangle \text{Orb}^\star(\gamma, dt_v)$$

We will frequently omit the dependence on the Haar measure dt_v when it is understood from context. When $\kappa = \mathbf{1}$ is trivial, we denote $\text{SO}_a^\star = \text{Orb}_a^{\star, \mathbf{1}}$.

We first explain how a character on this cohomology group gives rise to an endoscopic datum of $X^\mathbf{S}$ in the sense of Definition 2.12. Recall that Tate–Nakayama duality gives $H^1(F_v, I_{\gamma_0})^D \simeq \pi_0(\hat{I}_{\gamma_0}^\Gamma)$. We may identify dual tori $\hat{I}_{\gamma_0} \simeq \hat{\mathbf{C}}$, which may not respect the Galois actions on the two dual tori. Nevertheless, [Les24, Lemma 7.14] implies that there exists a 1-cocycle $w : \Gamma \rightarrow W_X$ valued in the little Weyl group of $X^\star$ such that this isomorphism becomes Γ -equivariant if we endow $\hat{\mathbf{C}}$ with the twisted action

$$\sigma * t = w_\sigma(t).$$

Thus we may identify these two tori, and if $\kappa \in H^1(F_v, I_{\gamma_0})^D \simeq \pi_0(\hat{\mathbf{C}}^{\Gamma, *})$, we may choose a lift $\dot{\kappa} \in \hat{\mathbf{C}}^{\Gamma, *}$. Composing with the symmetrization map $s : \hat{\mathbf{X}} \rightarrow \hat{\mathbf{G}}$, we obtain an endoscopic datum via the classical construction (cf. [Ngo10, §1.8]).

For an endoscopic datum $\mathfrak{e} = (\kappa, \rho)$ for $X^\mathbf{S}$, recall from Lemma 2.21 that we have a reduced divisor $\mathfrak{R}_\mathfrak{e}$ such that

$$(7.1) \quad \nu_\mathfrak{e}^* \mathcal{D}^{\text{cent}} = \mathcal{D}_\mathfrak{e}^{\text{cent}} + 2\mathfrak{R}_\mathfrak{e}, \quad \nu_\mathfrak{e}^* \mathcal{D}^{\text{br}} = \mathcal{D}_\mathfrak{e}^{\text{br}}.$$

For $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$, we denote

$$(7.2) \quad r_\mathfrak{e}(a) = \deg(a^* \mathfrak{R}_\mathfrak{e}), \quad d^{\text{cent}}(a) = \deg(a^* \mathcal{D}^{\text{cent}}), \quad d^{\text{br}}(a) = \deg(a^* \mathcal{D}^{\text{br}})$$

We similarly denote $d_\mathfrak{e}^{\text{cent}}(a_\mathfrak{e})$ and $d_\mathfrak{e}^{\text{br}}(a_\mathfrak{e})$ for $a_\mathfrak{e} \in \mathfrak{c}_\mathfrak{e}^\heartsuit(\mathcal{O}_v)$. Note that the identities (7.1) imply that, if $a_\mathfrak{e}$ maps to a under $\nu_\mathfrak{e}$, then $d^{\text{br}}(a) = d_\mathfrak{e}^{\text{br}}(a_\mathfrak{e})$ and $d^{\text{cent}}(a) = d_\mathfrak{e}^{\text{cent}}(a_\mathfrak{e}) + 2r_\mathfrak{e}(a_\mathfrak{e})$.

Note that if $\nu_\mathfrak{e}(a_\mathfrak{e}) = a$ for $a_\mathfrak{e} \in \mathfrak{c}_\mathfrak{e}(\mathcal{O}_v)$ and $a \in \mathfrak{c}^\heartsuit$, then $J_a(F_v) \simeq J_{a_\mathfrak{e}}^\mathfrak{e}$ by Proposition 2.14. Hence, for a choice of Haar measure dt_v on $J_a(F_v)$, we also obtain such a measure on $J_{a_\mathfrak{e}}^\mathfrak{e}$. The main theorem of this paper is the following.

Theorem 7.1. *Let $\mathfrak{e}$ be an elliptic endoscopic datum, and suppose that $a \in \mathfrak{e}^\vee(\mathcal{O}_v)$ and $a_\mathfrak{e} \in \mathfrak{e}_\mathfrak{e}^\vee(\mathcal{O}_v)$ are matched in the sense that $\nu_\mathfrak{e}(a_\mathfrak{e}) = a$. Then, we have*

$$\mathrm{Orb}_a^{\mathbf{S}, \kappa}(dt_v) = q^{r_\mathfrak{e}(a)} \mathrm{SO}_{a_\mathfrak{e}}^\mathfrak{e}(dt_v).$$

We will drop the measure dt_v from the notation for orbital integrals where it is understood. This result will be reinterpreted geometrically in terms of RASFs in Theorem 7.9 after we've developed some point counting techniques for affine Springer fibers. The proof of Theorem 7.1 is completed in §9.

7.2. Point Counting on RASFs. We now interpret these orbital integral as weighted point counts of groupoids of k -points associated to relative affine Springer fibers. We first recall the classical setting in [Ngo10, Chapter 8], which will apply to our relative Hitchin systems verbatim.

7.2.1. General framework. Let P be a smooth commutative group acting on a scheme M over k . We assume that $\pi_0(P)$ is a finite-type abelian group, the stabilizer $\mathrm{Stab}_P(m)$ is finite type for all points $m \in M$, and there exists a discrete subgroup $\Lambda \subset P$ such that P/Λ and M/Λ are finite type. This will apply to our cases of interest by Proposition 6.16.

For P and M satisfying the above properties, we put $\mathcal{X} = [M/P]$ the stack over k whose set of k -points is given by

$$\mathcal{X}(k) = \{(m, p) \in M(\bar{k}) \times P(\bar{k}) \mid p \cdot \sigma(m) = m\},$$

where we recall that σ denotes the Frobenius element of k . The k -points $\mathcal{X}(k)$ naturally have the structure of a groupoid. For any $x \in \mathcal{X}(k)$, we denote by $\mathrm{Aut}_{\mathcal{X}(k)}(x)$ the automorphism group of x . Let P_σ denote the set of σ -twisted conjugacy classes in $P(\bar{k})$, i.e. elements of P_σ are sets

$$\{gp\sigma(g)^{-1} \mid g \in P(\bar{k})\}$$

for $p \in P(\bar{k})$. The Galois cohomology $H^1(k, P)$ is the maximal torsion subgroup contained in P_σ . We have a class map

$$\mathrm{cl}: \mathcal{X}(k) \rightarrow H^1(k, P)$$

defined by taking (m, p) to the σ -twisted conjugacy class of p .

Definition 7.2. For any character $\kappa: H^1(k, P) \rightarrow \mathbb{Q}^\times$, we define

$$\#\mathcal{X}(k)_\kappa = \sum_{x \in |\mathcal{X}(k)|} \frac{\kappa(x)}{\#\mathrm{Aut}_{\mathcal{X}(k)}(x)}$$

where the sum above ranges over isomorphism classes in $\mathcal{X}(k)$. Here we abuse notation and set $\kappa(x) = \kappa(\mathrm{cl}(x))$. When $\kappa = 1$ is the trivial character, we abbreviate this sum simply by $\#\mathcal{X}(k)$.

We now specialize to our cases of interest. Let $a \in \mathfrak{e}^\vee(\mathcal{O}_v)$, and let $\mathcal{X} = [\mathcal{M}_v^*(a)/\mathcal{P}_v(J'_a)]$ for $J'_a \rightarrow J_a$ a fixed group scheme with connected fibers over $\mathcal{O}_v$ which induces an isomorphism on the generic fiber. We will deduce a formula relating the point counts $\#[\mathcal{M}_v^*(a)/\mathcal{P}_v(J'_a)](k)_\kappa$ with the κ orbital integral $\mathrm{Orb}_a^{\mathbf{S}, \kappa}$. Note that by Theorem 2.9, J_a has connected fibers for all $a \in \mathfrak{e}^\vee(\mathcal{O}_v)$, and hence all results that follow will hold for $J'_a = J_a$. Recall the notation of $\mathfrak{X}_\gamma^*$ from (6.3) and Proposition 6.7.

We begin by recalling a lemma from [Ngo10].

Lemma 7.3. ([Ngo10, Lemma 8.2.4]) *Let J'_a be a smooth group scheme over $\mathcal{O}_v$ with connected fibers, and let $J'_a \rightarrow J_a$ be an homomorphism of group schemes over $\mathcal{O}_v$ which is an isomorphism over F_v . Then, there is an isomorphism $H^1(k, \mathcal{P}_v(J'_a)) \simeq H^1(F_v, J_a)$.*

When dealing with endoscopic cases, these results will also apply to the morphism $J_a \rightarrow J_{\mathfrak{e}, a_\mathfrak{e}}$ with the unfortunate notational mishap of considering the case when $J = J_{\mathfrak{e}, a_\mathfrak{e}}$.

The class function now gives a map

$$(7.3) \quad \text{cl}: [\mathcal{M}_v^*(a)/\mathcal{P}_v(J'_a)](k) \rightarrow H^1(F_v, J_a).$$

This map is not surjective; we characterize its image in the following lemma.

Lemma 7.4. *Let $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$, and choose $\gamma \in V^*(F_v)$ lifting a . The canonical isomorphism $J_a \simeq I_\gamma \subset H^*(F_v)$ from Proposition 2.4 induces a morphism*

$$(7.4) \quad H^1(F_v, J_a) \rightarrow H^1(F_v, H^*)$$

Let $\ker_1^{*,\gamma}$ denote the kernel of (7.4). Then, for any $x \in [\mathfrak{X}_\gamma^*/\mathcal{P}_v(J'_a)](k)$, we have $\text{cl}(x) \in \ker_1^{*,\gamma}$.

Proof. Let $x = (m, p)$. We write $m = gH^*(\check{\mathcal{O}}_v)$ for $g \in H^*(\check{F}_v)$ and $p = jJ_a(\check{\mathcal{O}}_v)$ for $j \in J_a(\check{F}_v)$. The condition $p \cdot \sigma(m) = m$ is equivalent to $g^{-1} \cdot j \cdot \sigma(g) \in H^*(\check{\mathcal{O}}_v)$. Since every element of $H^*(\check{\mathcal{O}}_v)$ is σ -conjugate to the identity, we conclude that j is σ -conjugate to the identity under $H^*(\check{F}_v)$, and $\text{cl}(x) \in \ker_1^{*,\gamma}$. $\square$

Proposition 7.5. *For $\kappa \in H^1(F_v, J_a)^D$ a character, we have an equality*

$$\#[\mathcal{M}_v^*(a)/\mathcal{P}_v(J'_a)](k)_\kappa = \text{vol}(J'_a(\mathcal{O}_v), dt_v) \cdot \text{Orb}_a^{*,\kappa}$$

Proof. We may decompose

$$[\mathcal{M}_v^*(a)/\mathcal{P}_v(J'_a)](k) = \bigsqcup_{\xi \in \ker_1^{*,\gamma}} [\mathfrak{X}_\gamma^*/\mathcal{P}_v(J'_a)](k)_\xi$$

where the disjoint union is over the image under the class function. That is, if $j_\xi \in \mathcal{P}_v(J'_a)$ denotes a representative for the σ -twisted conjugacy class $\xi \in H^1(F_v, J_a) \simeq H^1(k, \mathcal{P}_v(J'_a))$, then we have

$$[\mathfrak{X}_\gamma^*/\mathcal{P}_v(J'_a)](k)_\xi = \{h \in \text{Gr}_H(\bar{k}) \mid h^{-1}j_\xi\sigma(h) = 1 \text{ and } \text{ad}(h)^{-1}\gamma \in V^*(\check{\mathcal{O}}_v)\}$$

Since the image of j_ξ lies in $\ker_1^{*,\gamma}$, there exists some $h_\xi \in H^*(\check{F})$ such that $h_\xi^{-1}j_\xi\sigma(h_\xi) = 1$. For any $h \in [\mathfrak{X}_\gamma^*/\mathcal{P}_v(J'_a)](k)_\xi$, the map sending h to $h' = h_\xi^{-1}h \in \text{Gr}_H(k)$ gives an isomorphism of groupoids

$$(7.5) \quad [\mathfrak{X}_\gamma^*/\mathcal{P}_v(J'_a)](k)_\xi = \{h' \in \text{Gr}_H(k) \mid \text{ad}(h')^{-1}\gamma_\xi \in V^*(\mathcal{O}_v)\}$$

where $\gamma_\xi = \text{ad}(h_\xi)^{-1}\gamma$. Note the groupoid structure on the above is realized through the action of I_{γ_ξ} on the right-hand side.

It is evident that the isomorphism classes of the groupoid on the right-hand side of (7.5) are given by the double coset space

$$\{h' \in I_{\gamma_\xi} \backslash H^*(F_v)/H^*(\mathcal{O}_v) \mid \text{ad}(h')^{-1}\gamma_\xi \in V^*(\mathcal{O}_v)\}$$

with automorphism groups

$$\text{Aut}(h') = (I_{\gamma_\xi} \cap h'H^*(\mathcal{O}_v)(h')^{-1})/J'_a(\mathcal{O}_v)$$

Thus, we compute that

$$\begin{aligned} \#[\mathfrak{X}_\gamma^*/\mathcal{P}_v(J'_a)](k)_\xi &= \text{vol}(J'_a(\mathcal{O}_v), dt_v) \cdot \int_{I_{\gamma_\xi} \backslash \text{Gr}_H} \mathbb{1}_{V^*(\mathcal{O}_v)}(\text{ad}(h')^{-1}\gamma_\xi) \cdot dt_v \\ &= \text{vol}(J'_a(\mathcal{O}_v), dt_v) \cdot \text{Orb}(\gamma_\xi) \end{aligned}$$

We conclude by summing over $\xi \in \ker_1^{*,\gamma}$ according to the appropriate κ weights. $\square$

Since J has connected fibers, we have the following special case.

Corollary 7.6. *Fix $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$. Let κ be a character of $H^1(k, \mathcal{P}_v(J_a)) \simeq H^1(F_v, J_a)$. Then, we have*

$$\#[\mathcal{M}_v^*(a)/\mathcal{P}_v(J_a)](k)_\kappa = \text{vol}(J_a(\mathcal{O}_v), dt_v) \cdot \text{Orb}_a^{*,\kappa}$$

7.3. Point counting along the relative Hitchin fibration. We now relate the local point-counting formula in Proposition 7.5 to point counts along the anisotropic locus in the relative Hitchin fibration.

Let $a \in \mathcal{A}^{\text{ani}}(k)$, and let $J'_a \rightarrow J_a$ be a map of group schemes over Σ . As always, we demand in the sequel that J'_a have connected geometric fibers and that the map be an isomorphism on the generic fiber. Define $\mathcal{P}'_a$ to be the space of J'_a -torsors on Σ , equipped with the natural map $\mathcal{P}'_a \rightarrow \mathcal{P}_a$.

We form the stacky quotient $[\mathcal{M}'_a/\mathcal{P}'_a]$. The class map for x the groupoid $[\mathcal{M}'_a/\mathcal{P}'_a](k)$ lands in $\text{cl}(x) \in H^1(k, \mathcal{P}'_a) \simeq \pi_0(\mathcal{P}'_a)_\sigma$, so we may consider the weighted point count $\#[\mathcal{M}'_a/\mathcal{P}'_a](k)_{\kappa'}$ as in §7.2 for $\kappa' \in H^1(k, \pi_0(\mathcal{P}'_a))^D$. We consider κ' that arise as follows. Recall that, since $\pi_0(\mathcal{P}_a)$ is finite, we have $H^1(k, \pi_0(\mathcal{P}_a)) \simeq \pi_0(\mathcal{P}_a)_\sigma$ is naturally identified with the space of Frobenius-twisted conjugacy classes in $\pi_0(\mathcal{P}_a)$, and similarly for $\mathcal{P}'_a$. Given a σ -invariant character

$$\kappa: \pi_0(\mathcal{P}_a) \rightarrow \overline{\mathbb{Q}}_\ell^\times,$$

we obtain a character κ' on $H^1(k, \pi_0(\mathcal{P}'_a))$ by pre-composing with the map $\pi_0(\mathcal{P}'_a) \rightarrow \pi_0(\mathcal{P}_a)$. We denote the corresponding point count by $\#[\mathcal{M}'_a/\mathcal{P}'_a](k)_\kappa$.

Similarly, for any $v \in |\Sigma|$, we have the Beauville–Laszlo gluing map $\mathcal{P}_v(J'_a) \rightarrow \mathcal{P}'_a$. Pre-composing κ as above with the map $\pi_0(\mathcal{P}_v(J'_a))_\sigma \rightarrow \pi_0(\mathcal{P}'_a)_\sigma$ gives a character κ'_v of $H^1(k, \pi_0(\mathcal{P}_v(J'_a))) \simeq H^1(F_v, J'_a)$. We again denote the corresponding weighted point count of the RASF at v by $\#[\mathcal{M}'_v(a)/\mathcal{P}_v(J'_a)](k)_\kappa$. We can relate the global and local point counts now as follows.

Proposition 7.7. *Let $a \in \mathcal{A}^{\text{ani}}(k)$, and set $U_a = a^{-1}(\mathfrak{c}_D^{\text{rs}})$ where $\mathfrak{c}^{\text{rs}} = \mathfrak{c} - (\mathfrak{D}^{\text{cent}} \cup \mathfrak{D}^{\text{br}})$. For any σ -invariant character $\kappa: \pi_0(\mathcal{P}_a) \rightarrow \overline{\mathbb{Q}}_\ell^\times$, we have*

$$(7.6) \quad \#[\mathcal{M}'_a/\mathcal{P}'_a](k)_\kappa = \prod_{v \in |\Sigma - U_a|} \#[\mathcal{M}'_v(a)/\mathcal{P}_v(J'_a)](k_v)_{\kappa}$$

Proof. Recall that by the product formula (Proposition 6.24), we have an equivalence of groupoids

$$[\mathcal{M}'_a/\mathcal{P}'_a](k) \simeq \prod_{v \in |\Sigma - U_a|} [\mathcal{M}'_v(a)/\mathcal{P}_v(J'_a)](k_v)$$

Consider the class functions cl defined in (7.3) on the groupoids $[\mathcal{M}'_a/\mathcal{P}'_a](k)$ and $[\mathcal{M}'_v(a)/\mathcal{P}_v(J'_a)](k_v)$. Then, by the definition of κ' and κ'_v , the above equivalence preserves the class function; that is,

$$\langle \text{cl}(x), \kappa' \rangle = \prod_{v \in |\Sigma - U_a|} \langle \text{cl}(x_v), \kappa'_v \rangle.$$

for any $x \in [\mathcal{M}'_a/\mathcal{P}'_a](k)$ and $x_v \in [\mathcal{M}'_v(a)/\mathcal{P}_v(J'_a)](k_v)$ corresponding to one another via the product formula. The proposition then follows. $\square$

By computing the lefthand side of (7.6) using the Grothendieck–Lefschetz trace formula, we now deduce:

Corollary 7.8. *For any σ -invariant character $\kappa: \pi_0(\mathcal{P}_a) \rightarrow \overline{\mathbb{Q}}_\ell^\times$, we have*

$$\sum_{i \geq 0} (-1)^i \text{tr} \left(\sigma, H^i(\mathcal{M}'_a)_\kappa \right) = \#(\mathcal{P}'_a)^\circ(k) \cdot \prod_{v \in |\Sigma - U_a|} \#[\mathcal{M}'_v(a)/\mathcal{P}_v(J'_a)](k_v)_\kappa$$

Proof. Follows directly from Proposition 7.7 and [Ngo10, Proposition 8.1.6]. $\square$

7.4. The geometric relative Fundamental Lemma and cases. In this section, we state the precise form of our main result using the language of RASFs. We then establish this result for a few basic cases, which will serve as base cases for the global argument in §9.

By Proposition 7.5, Theorem 7.1 is equivalent to the (local) geometric incarnation below.

Theorem 7.9. *Let $\mathfrak{e}$ be an elliptic endoscopic datum, and suppose that $a \in \mathfrak{e}^\vee(\mathcal{O}_v)$ and $a_{\mathfrak{e}} \in \mathfrak{e}_{\mathfrak{e}}^\vee(\mathcal{O}_v)$ are such that $\nu_{\mathfrak{e}}(a_{\mathfrak{e}}) = a$. Let $J'_a \rightarrow J_a$ be a smooth commutative group scheme over $\mathcal{O}_v$ with connected fibers which is isomorphic to J_a on the generic fiber $\Sigma_v^\bullet = \text{Spec}(F_v)$. Then, for all finite extensions K/k_v , we have*

$$(7.7) \quad \#[\mathcal{M}_v^{\mathbf{S}}(a)/\mathcal{P}_v(J'_a)](K)_\kappa = q^{r_{\mathfrak{e}}(a) \deg(v)[K:k_v]} \cdot \#[\mathcal{M}_v^{\mathfrak{e}}(a_{\mathfrak{e}})/\mathcal{P}_v(J'_a)](K)$$

Remark 7.10. We note that since J_a has connected fibers, Theorem 7.9 remains true for $J'_a = J_a$. It is important to note, however, that the quotient on either side of (7.7) is by the *same* symmetry group.

The full proof of Theorem 7.9 will be contained in §9. For now, we consider some particularly simple cases. As a first example, we consider the case of nontrivial obstruction class, where the theorem is a consequence of a functional equation; see also [Yun11, Lem 2.5.3] and [WZ24, Lem 3.6.4].

Proposition 7.11. *Suppose that $a, a_{\mathfrak{e}}$ are as in Theorem 7.9, and $\kappa \in H^1(F_v, J_a)^D$ is a nontrivial character. If $\text{Ob}_v(a_{\mathfrak{e}}) \neq 0$ (i.e., $\text{val}(\det(a))$ is odd), then $\text{Orb}_a^{\mathbf{S}, \kappa} = 0$. In particular, Theorem 7.9 holds in this case with both sides being equal to zero.*

Proof. Note that since $\text{Ob}_v(a) \neq 0$, we have $\mathcal{M}_v^{\mathfrak{e}}(a)(k) = \emptyset$, and hence the stable point count is $\#[\mathcal{M}_v^{\mathfrak{e}}(a)/\mathcal{P}_v(J_a)](k) = 0$. To show the κ -twisted point count on $\mathcal{M}_v^{\mathbf{S}}(a)$ vanishes, we construct a duality $\mathbb{D}$ on $\mathcal{M}_v^{\mathbf{S}}(a)(\bar{k})$, which will produce a functional equation for the point count at a . Recall from Proposition 6.7 that $\mathcal{M}_v^{\mathbf{S}}(a)(\bar{k})$ is the moduli of tuples $(\mathcal{E}', h, \iota, \phi'; \xi)$ where $(\mathcal{E}', h, \iota, \phi')$ is a Higgs object consisting of:

- (1) A free rank $2n$ $\mathcal{O}' \otimes_k \bar{k}$ -module $\mathcal{E}'$;
- (2) A nondegenerate Hermitian form $h: \mathcal{E}' \rightarrow \sigma^* \mathcal{E}'^\vee$;
- (3) An $\mathcal{O}' \otimes_k \bar{k}$ -linear involution $\iota: \mathcal{E}' \rightarrow \mathcal{E}'$ such that $h \circ \iota = -h$;
- (4) An $\mathcal{O}' \otimes_k \bar{k}$ -linear endomorphism $\phi': \mathcal{E}' \rightarrow \mathcal{E}'$ such that $\iota \circ \phi' = -\phi' \circ \iota$, $h(\phi'v, w) + h(v, \phi'w) = 0$, and ϕ' has characteristic polynomial a ;

and ξ is a trivialization on the punctured infinitesimal disk $\check{\Sigma}_v^\bullet = \text{Spec } \check{F}'_v$, i.e.

$$\xi: (\mathcal{E}', h, \iota, \phi')|_{\check{\Sigma}_v^\bullet} \xrightarrow{\sim} (R_a(\check{F}'_v), h^{\mathbf{S}}, \iota_R, [x]).$$

There is a natural duality on Higgs objects given by

$$\mathbb{D}(\mathcal{E}', h, \iota, \phi') = (\mathcal{E}'^\vee, (h^{-1})^\vee, \iota^\vee, \phi'^\vee)$$

Note that over any extension of F_v where F'_v splits, there exists an isomorphism

$$\theta: \mathbb{D}(R_a(\check{F}'_v), h^{\mathbf{S}}, \iota_R, [x]) \rightarrow (R_a(\check{F}'_v), h^{\mathbf{S}}, \iota_R, [x]).$$

In fact, the space of isomorphisms is a torsor $\mathcal{T}_a$ under the action of J_a , and so determines a class $[\mathcal{T}_a] \in H^1(F_v, J_a)$. For any choice of θ , we have an explicit cocycle representing $[\mathcal{T}_a]$ given by

$$d_\theta = \theta \cdot \sigma(\theta)^{-1},$$

where $\sigma(\theta)$ is the Frobenius-twisted isomorphism $\sigma \circ \theta \circ \sigma^{-1}$.

Fix a choice of such a θ , and now define a duality $\mathbb{D} = \mathbb{D}_\theta$ on $\mathcal{M}_v(a)(\bar{k})$ by

$$\mathbb{D}(\mathcal{E}', h, \iota, \phi'; \xi) = (\mathbb{D}(\mathcal{E}', h, \iota, \phi'); \theta \circ (\xi^{-1})^\vee).$$

This is compatible with inversion on $\mathcal{P}_v(J_a)$ in the sense that for every $m \in \mathcal{M}_v(a)(\bar{k})$ and $p \in \mathcal{P}_v(J_a)(\bar{k})$, we have

$$\mathbb{D}(p \cdot m) = p^{-1} \cdot \mathbb{D}(m).$$

The duality $\mathbb{D}$ satisfies the compatibility with Frobenius given by the cocycle d_θ ; that is, for any $m \in \mathcal{M}_v(a)(\bar{k})$,

$$\mathbb{D}(\sigma(m)) = d_\theta \cdot \sigma(\mathbb{D}(m))$$

Therefore, $\mathbb{D}$ induces an involution (which we also call $\mathbb{D}$) on the groupoid $[\mathcal{M}_v^{\mathbf{S}}(a)/\mathcal{P}_v(J_a)](k)$ by $\mathbb{D}(m, p) = (\mathbb{D}(m), d_\theta \cdot p^{-1})$. For any $x = (m, p)$, we compute

$$\text{cl}(\mathbb{D}(x)) = \text{cl}(\mathbb{D}(m), d_\theta \cdot p^{-1}) = d_\theta \cdot p^{-1} \in \pi_0(\mathcal{P}_v(J_a))_\sigma = H^1(F, J_a)$$

and hence $\text{cl}(\mathbb{D}(x)) = [\mathcal{T}_a] \cdot \text{cl}(x)^{-1}$ in $H^1(F_v, J_a)$.

Suppose now that $\kappa \in H^1(F_v, J_a)^D$ and $\nu_\epsilon(a_\epsilon) = a$. For any $x \in [\mathcal{M}_v^{\mathbf{S}}(a)/\mathcal{P}_v(J_a)](k)$, by our formula for the class function of $\mathbb{D}(x)$, we have

$$\langle \kappa, \text{cl}(\mathbb{D}(x)) \rangle = \kappa([\mathcal{T}_a]) \cdot \kappa(\text{cl}(x))^{-1}.$$

Since $H^1(F_v, J_a)$ is 2-torsion, $\kappa(\text{cl}(x))^{-1} = \kappa(\text{cl}(x))$. On the other hand, we prove the following lemma to compute $\kappa([\mathcal{T}_a])$.

Lemma 7.12. *With notation as above, write $X^\epsilon = X_r^{\mathbf{S}} \times X_s^{\mathbf{U}}$ and decompose $a_\epsilon = (a_\epsilon^{\mathbf{S}}, a_\epsilon^{\mathbf{U}})$. Then*

$$\kappa([\mathcal{T}_a]) = (-1)^{\text{val}(a_\epsilon^{\mathbf{U}})_{2s}} = \begin{cases} 1 & \text{if } \text{Ob}(a_\epsilon) = 0 \\ -1 & \text{if } \text{Ob}(a_\epsilon) \neq 0 \end{cases}$$

Proof. By the description given in Theorem 2.9, we compute

$$H^1(F_v, J_a) = R_a(F_v)^{\iota_R} / \text{Nm}(R_a(F_v')^{\iota_R})$$

where Nm denotes the norm map

$$\text{Nm}: R_a(F_v')^{\iota_R} \rightarrow R_a(F_v)^{\iota_R}.$$

Recall that $R_a(F_v')^{\iota_R} = F_v[x^2]/(Q_a(x^2))$ for the polynomial Q_a satisfying $Q_a(x^2) = P_a(x)$. By Proposition 2.14, we see that the endoscopic character κ determines an ordered factorization $Q_a = Q^{\mathbf{S}} \cdot Q^{\mathbf{U}}$ and hence a decomposition

$$R_a(F_v)^{\iota_R} / \text{Nm}(R_a(F_v')^{\iota_R}) \simeq R^{\mathbf{S}} \oplus R^{\mathbf{U}}.$$

Note that the coefficients of $Q^{\mathbf{S}}$ (resp., $Q^{\mathbf{U}}$) are precisely those given by the tuple $a_\epsilon^{\mathbf{S}}$ (resp., $a_\epsilon^{\mathbf{U}}$). Pairing a class in $(r^{\mathbf{S}}, r^{\mathbf{U}}) \in R^{\mathbf{S}} \oplus R^{\mathbf{U}} \simeq H^1(F_v, J_a)$ with κ is given by (compare [Les22, Lemma A.2])

$$\kappa(r^{\mathbf{S}}, r^{\mathbf{U}}) = (-1)^{\text{val}(\text{Nm}_{R^{\mathbf{U}}/F_v}(r^{\mathbf{U}}))}.$$

We now compute the class of the torsor $\mathcal{T}_a$ in this group explicitly. If we express the multiplication-by- x map $[x]$ and its dual $[x]^\vee$ with respect to the decomposition $R_a(F_v') = R_a(F_v')^{\iota_R} \oplus x \cdot R_a(F_v')^{\iota_R}$, we get

$$[x] = \begin{pmatrix} 0 & x^2 \\ 1 & 0 \end{pmatrix}, \quad [x]^\vee = \begin{pmatrix} 0 & 1 \\ x^2 & 0 \end{pmatrix}.$$

If an isomorphism $\theta: R_a(F_v')^{\iota_R} \rightarrow R_a(F_v')^{\iota_R}$ preserves ι_R and intertwines $[x]$ and $[x]^\vee$, then θ must have the form

$$\theta = \begin{pmatrix} x^2 t & 0 \\ 0 & t \end{pmatrix}$$

for some $t \in (R_a(F_v')^{\iota_R})^\times$. To preserve h , we obtain the further condition on t that $t \cdot \sigma(t)^{-1} = x^{-2}$. Hence, we conclude that

$$[\mathcal{T}_a] = x^{-2} = x^2 \in R_a(F_v)^{\iota_R} / \text{Nm}(R_a(F_v')^{\iota_R})$$

Since $Q^{\mathbf{U}}$ is monic of degree s , we have $\text{Nm}_{R^{\mathbf{U}}/F_v}(x^2) = (-1)^s Q^{\mathbf{U}}(0)$ is the constant term of $Q^{\mathbf{U}}$ up to sign, which is exactly $(a_\epsilon^{\mathbf{U}})_{2s}$. We therefore compute

$$\kappa([\mathcal{T}_a]) = (-1)^{\text{val}(a_\epsilon^{\mathbf{U}})_{2s}}.$$

The final line of the lemma follows from the computation of $\text{Ob}(a_\epsilon)$ given in Lemma 6.13. $\square$

Hence, we conclude from Corollary 7.6 the functional equation

$$\mathrm{Orb}_a^{\mathbf{S}, \kappa} = (-1)^{\mathrm{Ob}(a_\epsilon)} \mathrm{Orb}_a^{\mathbf{S}, \kappa}$$

from which the Proposition 7.11 follows. $\square$

We conclude this section by proving Theorem 7.9 for some special cases of split elements for which the dimension of the corresponding relative affine Springer fiber is at most one. We note that the geometry of all but the first of these cases appear in [Ngo10, Lem 8.5.7].

Proposition 7.13. *Let ϵ be an elliptic endoscopic datum. Theorem 7.9 holds for all $a_\epsilon \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$ with image $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$ satisfying either (c.f. (7.2))*

$$d^{\mathrm{br}}(a) = d^{\mathrm{cent}}(a) = 0$$

or

$$d^{\mathrm{br}}(a) = 0, \quad d^{\mathrm{cent}}(a) = d_\epsilon^{\mathrm{cent}}(a_\epsilon) = 1.$$

Moreover, in this case, both sides of (7.7) are nonzero rational numbers.

Proof. By the dimension formula (Theorem 6.28), we must have $\delta_v(a) = \delta_{\epsilon, v}(a) = 0$, so that by the proof of Proposition 6.20, the affine Springer fiber $\mathcal{M}_v^{\mathbf{S}}(a)$, respectively $\mathcal{M}_v^\epsilon(a)$, is a torsor for the action of the lattice $\mathcal{P}_v(J_a) = \mathcal{P}_v(J_{\epsilon, a_\epsilon}) = \Gamma_a$. Therefore, on the endoscopic side, we have

$$1 = \#[\mathcal{M}_v^\epsilon(a_\epsilon)/\mathcal{P}_v(J_{\epsilon, a_\epsilon})](k) = \mathrm{vol}(J_{\epsilon, a_\epsilon}(\mathcal{O}_v)) \cdot \mathrm{SO}_{a_\epsilon}$$

by Corollary 7.6 as well as

$$\#[\mathcal{M}_v^\epsilon(a_\epsilon)/\mathcal{P}_v(J'_a)](k) = \mathrm{vol}(J'_a(\mathcal{O}_v)) \cdot \mathrm{SO}_{a_\epsilon}$$

by Proposition 7.5. Combining these statements, we get that

$$\#[\mathcal{M}_v^\epsilon(a_\epsilon)/\mathcal{P}_v(J'_a)](k) = \frac{\mathrm{vol}(J'_a(\mathcal{O}_v))}{\mathrm{vol}(J_{\epsilon, a_\epsilon}(\mathcal{O}_v))}.$$

By an identical argument, we also get

$$\#[\mathcal{M}_v^{\mathbf{S}}(a)/\mathcal{P}_v(J'_a)](k) = \frac{\mathrm{vol}(J'_a(\mathcal{O}_v))}{\mathrm{vol}(J_a(\mathcal{O}_v))}.$$

To conclude, we note in these cases that the morphism $J_a \rightarrow J_{\epsilon, a_\epsilon}$ is an isomorphism so that the two denominators above agree. $\square$

Proposition 7.14. *Let ϵ be an elliptic endoscopic datum. Theorem 7.9 holds for $a \in \mathfrak{c}^\heartsuit(\mathcal{O}_v)$ such that $d^{\mathrm{br}}(a) = 0$, $d^{\mathrm{cent}}(a) = 2$ and $d_\epsilon^{\mathrm{cent}}(a) = 0$. Moreover, in this case, both sides of (7.7) are nonzero rational numbers.*

Proof. Since $d_\epsilon^{\mathrm{cent}}(a) = 0$, by the dimension formula (Theorem 6.28) we have $c_{\epsilon, v}(a_\epsilon) = 0$. Since $c_v(a)$ depends only on the generic fiber and $J_a \rightarrow J_{\epsilon, a_\epsilon}$ is an isomorphism on the generic fiber (see Proposition 2.14), it also follows that $c_v(a) = 0$. Therefore, we have $\delta_{\epsilon, v}(a_\epsilon) = 0$ while $\delta_v(a) = 1$, and a is split.

On the endoscopic side, an identical argument to the proof of Proposition 7.13 implies that

$$(7.8) \quad \#[\mathcal{M}_v^\epsilon(a)/\mathcal{P}_v(J'_a)](k) = \frac{\mathrm{vol}(J'_a(\mathcal{O}_v))}{\mathrm{vol}(J_{\epsilon, a_\epsilon}(\mathcal{O}_v))}.$$

We now consider the Siegel case. Consider the polynomial $Q_a(x^2) = P_a(x)$ (cf. Proposition 6.16). Since a is split, it follows that $Q_a(y)$ splits as a product of linear factors $Q_a(z) = \prod_{i=1}^n (z - u_i)$ over $\check{F}_v$. Furthermore, by Gauss's Lemma, $u_i \in \check{\mathcal{O}}_v$ for all i , and because $d^{\mathrm{br}}(a) = \mathrm{val}(a_{2n}) = 0$ and $a_{2n} = \prod_i u_i$, the u_i are units in $\check{\mathcal{O}}_v$. Since $d^{\mathrm{cent}}(a) = \mathrm{val} \mathrm{Disc}(Q_a) = 2$, there exists exactly one pair of roots, u_i, u_j , such that $\mathrm{val}(u_i - u_j) = 1$, and all other differences $u_{i'} - u_{j'}$ are units. By the assumption that $d_\epsilon^{\mathrm{cent}}(a_\epsilon) = 0$, the κ factorization $Q_a = Q^{\mathbf{S}} Q^{\mathbf{U}}$ places u_i and u_j in different factors.

We reuse notation $R = R_a(\check{\mathcal{O}}_v)$ and R^ι from Section 6.4. We also use $\widetilde{R}^\iota$ to denote the normalization of R^ι . By Theorem 2.9 and Proposition 2.14, we have identifications

$$J_a(\check{\mathcal{O}}_v) \simeq \check{\mathcal{O}}_v[y]/Q_a(y) = R^\iota,$$

and of $J_{\mathfrak{e},a_\mathfrak{e}}(\check{\mathcal{O}}_v)$ with the normalization

$$J_a(\check{\mathcal{O}}_v) \simeq \check{\mathcal{O}}_v[y]/Q^{\mathbf{S}}(y) \times \check{\mathcal{O}}_v[y]/Q^{\mathbf{U}}(y) = \widetilde{R}^\iota.$$

Define a torus $A_a := J_a(\check{\mathcal{O}}_v)/J_{\mathfrak{e},a_\mathfrak{e}}(\check{\mathcal{O}}_v) = \widetilde{R}^\iota/R^\iota$, with the natural k structure defined by the Galois actions on $J_a(\check{\mathcal{O}}_v)$ and $J_{\mathfrak{e},a_\mathfrak{e}}(\check{\mathcal{O}}_v)$. By our descriptions of $J_a(\check{\mathcal{O}}_v)$ and $J_{\mathfrak{e},a_\mathfrak{e}}(\check{\mathcal{O}}_v)$ above, the torus A_a is of rank 1.

Lemma 7.15. *There is a short exact sequence*

$$(7.9) \quad 0 \rightarrow A_a(\bar{k}) \rightarrow \mathcal{P}_v(J_a)(\bar{k}) \rightarrow \Gamma_a \rightarrow 0,$$

where Γ_a is the lattice of Section 6.4. Moreover, all morphisms are equivariant with respect to the Galois action of $\text{Gal}(\bar{k}/k_v)$.

Proof. Explicitly, the lattice Γ_a is given by

$$\Gamma_a = (R^\iota \otimes_{\check{\mathcal{O}}_v} \check{F}_v)/\widetilde{R}^\iota$$

Hence, the sequence (7.9) is explicitly

$$0 \rightarrow \widetilde{R}^\iota/R^\iota \rightarrow (R^\iota \otimes_{\check{\mathcal{O}}_v} \check{F}_v)/R^\iota \rightarrow (R^\iota \otimes_{\check{\mathcal{O}}_v} \check{F}_v)/\widetilde{R}^\iota \rightarrow 0,$$

whose exactness is evident. $\square$

Note in particular, that Γ_a is of rank n . We may compute

$$\#A_a(k_v) = \frac{\text{vol } J_{\mathfrak{e},a_\mathfrak{e}}(\mathcal{O}_v)}{\text{vol } J_a(\mathcal{O}_v)}.$$

The relative affine Springer fiber $\mathcal{M}_v^{\mathbf{S}}(a)$ was implicitly computed in Proposition 6.20; we make this computation more explicit here. Over $\bar{k}$, we have a lift of a of the form

$$\gamma = \begin{pmatrix} 0 & D \\ I_n & 0 \end{pmatrix} \in V^\epsilon(\bar{k}), \quad D = \text{diag}(u_1, \dots, u_n).$$

By Lemma 6.9, we compute that $\mathcal{M}_v^{\mathbf{S}}(a) \otimes_k \bar{k} \simeq \mathfrak{X}_\gamma$ is represented by a pair (g_1, g_2) with $g_1^{-1}Dg_2$ and $g_2^{-1}g_1$ having integral entries. The determinants multiply to the unit $\det D$, so both $g_1^{-1}Dg_2$ and $g_2^{-1}g_1$ are invertible over $\check{\mathcal{O}}_v$. In particular, $g_1 = g_2$ in the affine Grassmannian for GL_n , and we can identify $\mathcal{M}_v(a) \otimes_k \bar{k}$ with the GL_n affine Springer fiber of the diagonal matrix D . With the condition on D , this is precisely the case considered in [Ngo10, §8.3], with our torus A_a coinciding with Ngô's torus $A_{\pm\alpha}$. Namely, let

$$M = \bigcup_{j \in \mathbb{Z}} C_j, \quad C_j \simeq \mathbf{P}^1 \text{ for all } j$$

be an infinite chain of projective lines over $\bar{k}$ with a Galois action swapping components $\sigma(C_i) = C_{1-i}$ and a unique fixed point. There is an action of $P = A_a \times \mathbb{Z}$ on M with the torus A_a acting on each irreducible component C_j by fixing the two intersection points with consecutive components C_{j-1} and C_{j+1} , and $\mathbb{Z}$ acting by translating along the chain. There is a natural inclusion $P \subset \mathcal{P}_v(J_a)$, and the elementary computations in *loc. cit.* show that

$$\mathcal{M}_v(a) \simeq M \times^P \mathcal{P}_v(a),$$

with the natural action of $\mathcal{P}_v(a)$ through the second factor. Therefore, we have

$$\#[\mathcal{M}_v(a)/\mathcal{P}_v(J_a)](k)_\kappa = \#[M/P](k)_{\kappa'}$$

for the induced character $\kappa': H^1(k, P) \hookrightarrow H^1(k, \mathcal{P}_v(J_a)) \rightarrow \overline{\mathbb{Q}}_\ell^\times$. By [Ngo10, Example 8.1.15], we deduce that

$$\#[\mathcal{M}_v^{\mathbf{S}}(a)/\mathcal{P}_v(J_a)](k)_\kappa = \frac{q_v}{\#A_a(k_v)} = q_v \cdot \frac{\text{vol } J'_a(\mathcal{O}_v)}{\text{vol } J_{\mathfrak{e}, a_{\mathfrak{e}}}(\mathcal{O}_v)}.$$

Therefore, $\#[\mathcal{M}_v^{\mathbf{S}}(a)/\mathcal{P}_v(J'_a)](k)_\kappa = q \cdot \#[\mathcal{M}_v^{\mathfrak{e}}(a)/\mathcal{P}_v(J'_a)](k)$, and Theorem 7.9 follows in this case since $r_{\mathfrak{e}}(a) = 1$. $\square$

8. THE SUPPORT THEOREM AND COMPUTATION OF COHOMOLOGY

In this section, we recall Ngô's support theorem. In §8.2, we will apply this support theorem by analyzing top degree cohomology over $\mathcal{A}^\dagger$, which will demonstrate the importance of the regular quotient in relative endoscopy.

8.1. The support theorem. We begin by introducing some notions needed to state the support theorem.

Definition 8.1. Let S be a scheme. A *weak abelian fibration* over S is the data of a proper morphism $f: M \rightarrow S$, with M a Deligne–Mumford stack, together with a smooth commutative group scheme $g: P \rightarrow S$ acting on M over S such that

- (1) the morphisms f and g are of the same relative dimension d ;
- (2) for every $m \in M(k)$ mapping to $s \in S(k)$, the stabilizer of m in P_s is affine;
- (3) the Tate module $T_{\overline{\mathbb{Q}}_\ell}(P^0)$ of the fiberwise neutral component P^0 admits a polarization. More precisely, étale-locally on S there exists a pairing

$$(8.1) \quad T_{\overline{\mathbb{Q}}_\ell}(P^0) \times T_{\overline{\mathbb{Q}}_\ell}(P^0) \rightarrow \overline{\mathbb{Q}}_\ell(1)$$

such that, for any $s \in S(k)$, if

$$0 \rightarrow R_s \rightarrow P_s^0 \rightarrow A_s \rightarrow 0$$

is the Chevalley decomposition of P_s^0 —with R_s the maximal connected affine subgroup of P_s^0 and A_s an abelian variety—then when restricted to s , the bilinear map (8.1) factors through $T_{\overline{\mathbb{Q}}_\ell}(A_s)$ and induces a perfect pairing on $T_{\overline{\mathbb{Q}}_\ell}(A_s)$.

Given a weak abelian fibration, we have a δ -stratification on S determined by the function $\delta(s) = \dim R_s$, where $R_s \subset P_s$ is the maximal connected affine subgroup of P_s as above.

Definition 8.2. A weak abelian fibration (f, g) on S is δ -regular if for all $\delta \geq 0$, we have $\text{codim}_S(S_\delta) \geq \delta$.

We now can state Ngô's support theorem.

Theorem 8.3. ([Ngo10, Theorem 7.2.1]) *Let S be a geometrically irreducible scheme over k , and let $f: M \rightarrow S$ and $g: P \rightarrow S$ form a δ -regular weak abelian fibration of relative dimension d . Let $\mathcal{F}$ be a pure and self-dual constructible sheaf on M^1 .*

Let K be a simple constituent of the perverse cohomology sheaf ${}^p\mathbf{H}^n(f_\mathcal{F})$ with support Z . Then Z satisfies the following two properties.*

- (1) *Z is δ -critical, i.e. $\text{codim}_S(Z) = \delta_Z$ where δ_Z is the minimum value of the δ -invariant on Z .*
- (2) *There exists a non-empty open set $U \subset S_{\bar{k}}$ with $U \cap Z \neq \emptyset$ and a local system L on $U \cap Z$ such that i_*L is a direct factor of $R^{2d}f_*\mathcal{F}|_U$, the top degree ordinary cohomology sheaf restricted to U .*

¹For example, we will take $\mathcal{F} = \overline{\mathbb{Q}}_{\ell, M}$ when M is smooth.

8.2. Application to relative Hitchin systems. We now apply Theorem 8.3 to our cases of interest. We begin by showing that $(f^{\star, \text{ani}}, g^{\text{ani}})$ forms a weak abelian fibration.

Proposition 8.4. *The action of the group scheme $g^{\text{ani}}: \mathcal{P}^{\text{ani}} \rightarrow \mathcal{A}^{\text{ani}}$ on $f^{\star, \text{ani}}: \mathcal{M}^{\star, \text{ani}} \rightarrow \mathcal{A}^{\text{ani}}$ gives $(f^{\star, \text{ani}}, g^{\text{ani}})$ the structure of a weak abelian fibration.*

Proof. Proposition 5.8 shows that $f^{\star, \text{ani}}$ is a proper and flat morphism whose (equidimensional) fibers are of the same relative dimension as g^{ani} (see Corollary 6.29). Moreover, the centralizers of the action of $\mathcal{P}_a$ on $\mathcal{M}_a$ are identified with $H^0(\Sigma, J_a)$, which is affine since J_a is commutative. Finally, we note that $\mathcal{P}$ has polarizable Tate module via an identical argument to [Ngo10, §4.12]. $\square$

As the property of being a weak abelian fibration is stable under base change, the above also implies that $(\underline{f}^{\star, \text{ani}}, \underline{g}^{\text{ani}})$ is a weak abelian fibration, where the underline denotes the base change to the cover $\underline{\mathcal{A}}^{\text{ani}}$ defined in §5.3.

The property of δ -regularity is more delicate. While we expect $(\underline{f}^{\star, \text{ani}}, \underline{g}^{\text{ani}})$ is δ -regular, do not prove this statement. Rather we will skirt the issue of δ -regularity over the entire anisotropic locus by applying Corollary 5.30 and arguing for $\deg(D)$ large enough. For now, we restrict to the following locus.

Definition 8.5. Fix an effective divisor D on Σ . For any fixed positive integer $\delta \geq 0$, let $\underline{\mathcal{A}}_{<\delta}$ denote the union

$$\underline{\mathcal{A}}_{<\delta} = \bigsqcup_{\delta' < \delta} \underline{\mathcal{A}}_{\delta'}.$$

Let $\delta^{\text{bad}}(D)$ denote the minimal number δ such that the inequality

$$\text{codim}_{\underline{\mathcal{A}}^{\text{ani}}}(\underline{\mathcal{A}}_{\delta}) \geq \delta$$

fails; note that $\delta^{\text{bad}}(D) > 0$. The locus $\underline{\mathcal{A}}^{\text{good}}$ is defined as $\underline{\mathcal{A}}_{<\delta^{\text{bad}}(D)}$.

By definition, the restriction of $(\underline{f}^{\star, \text{ani}}, \underline{g}^{\text{ani}})$ to $\underline{\mathcal{A}}^{\text{good}}$ is δ -regular.

Recall from Definition 5.11 that the dagger locus $\mathcal{A}^{\dagger} \subset \mathcal{A}^{\text{ani}}$ denotes the subvariety of $\mathcal{A}^{\text{ani}}$ whose S -points $a: \Sigma \times S \rightarrow \mathfrak{c}_D$ satisfy that the image $a(\Sigma)$ intersects the branching divisor $\mathfrak{D}_D^{\text{br}}$ transversely. We now prove a density statement for $\mathcal{M}^{\star, \text{reg}}$ over $\mathcal{A}^{\dagger}$.

Lemma 8.6. *For any $a \in \mathcal{A}^{\dagger}(\bar{k})$, the regular locus $\mathcal{M}_a^{\star, \text{reg}}$ is dense in $\mathcal{M}_a^{\star}$. That is, the inclusion $\mathcal{M}^{\star, \text{reg}} \times_{\mathcal{A}} \mathcal{A}^{\dagger} \subset \mathcal{M}^{\star, \dagger}$ is fiberwise dense.*

Proof. Without loss of generality, we may assume that $k = \bar{k}$. By the product formula in Proposition 6.24, it suffices to check the above for RASFs $\mathcal{M}_v^{\star}(a)$ where $a \in \mathfrak{c}^{\heartsuit}(\mathcal{O}_v)$ is such that either a intersects transversely with $\mathfrak{D}^{\text{br}}$ or a does not intersect $\mathfrak{D}^{\text{br}}$ at all. In the latter case, $\mathcal{M}_v^{\star}(a)/\Gamma_a$ is connected, and so $\mathcal{M}_v^{\star}(a)$ contains a dense $\mathcal{P}_v(J_a)$ -orbit which is regular. In the former case, $\mathcal{M}_v^{\star}(a)/\Gamma_a$ has exactly two connected components. There are two non-conjugate choices of Kostant-Rallis section over $\bar{k}$, so each component of $\mathcal{M}_v^{\star}(a)/\Gamma_a$ contains the regular $\mathcal{P}_v(J_a)$ -orbit of a choice of Kostant-Rallis section. Each such orbit is dense in its connected component in $\mathcal{M}_v^{\star}(a)/\Gamma_a$, and so its preimage is dense in $\mathcal{M}_v^{\star}(a)$. $\square$

Recall from (5.5) that we have a decomposition

$$Rf_{\underline{\star}}^{\text{ani}} \overline{\mathbb{Q}}_{\ell} = \bigoplus_{\kappa \in \widehat{\mathbf{C}}} (Rf_{\underline{\star}}^{\text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa},$$

with $(Rf_{\underline{\star}}^{\text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}$ being the locus on which $\mathbb{X}^*(\widehat{\mathbf{C}})$ acts through $\underline{\mathcal{P}}$ by evaluation at $\kappa \in \widehat{\mathbf{C}}$. (See Proposition 5.24.) We denote the case $\kappa = 1$ by $(Rf_{\underline{\star}}^{\text{ani}} \overline{\mathbb{Q}}_{\ell})_{\text{st}}$ and call it the stable constituent.

Proposition 8.7. *In both the Siegel and unitary cases, the top degree stable cohomology over the anisotropic locus is a local system when restricted to $\underline{\mathcal{A}}^\dagger \subset \underline{\mathcal{A}}^{\text{ani}}$. That is,*

$$(R^{2d} f_{\star}^{\star, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\text{st}}|_{\underline{\mathcal{A}}^\dagger} = (R^{2d} f_{\star}^{\star, \dagger} \overline{\mathbb{Q}}_{\ell})_{\text{st}}$$

is a local system on $\underline{\mathcal{A}}^\dagger$.

Proof. By Lemma 8.6, we have $R^{2d} f_{\star}^{\star, \dagger} \overline{\mathbb{Q}}_{\ell} = R^{2d} (f_{\star}^{\star, \dagger}|_{\underline{\mathcal{M}}^{\text{reg}}})_! \overline{\mathbb{Q}}_{\ell}$, where in the latter we push forward the constant sheaf on $\underline{\mathcal{M}}^{\star, \text{reg}}$. Proposition 5.15 (cf. [HM25, Thm 1.1]) gives a factorization of $f^{\star}|_{\underline{\mathcal{M}}^{\star, \text{reg}}}$ into

$$\underline{\mathcal{M}}^{\star, \text{reg}} \xrightarrow{f^{\star, \text{reg}}} \underline{\mathcal{A}}^{\star, \text{reg}} \xrightarrow{\varphi^{\star}} \underline{\mathcal{A}},$$

where $f^{\star, \text{reg}}$ is a torsor for the smooth, commutative group scheme $\underline{\mathcal{P}} \times_{\underline{\mathcal{A}}} \underline{\mathcal{A}}^{\star, \text{reg}}$ over $\underline{\mathcal{A}}^{\star, \text{reg}}$, and $\varphi^{\star}$ is finite and étale. On top degree cohomology, this implies

$$(8.2) \quad R^{2d} (f_{\star}^{\star, \dagger}|_{\underline{\mathcal{M}}^{\text{reg}}})_! \overline{\mathbb{Q}}_{\ell} \simeq \varphi_{\star}^{\star} \left(R^{2d} f_{\dagger}^{\star, \text{reg}} \overline{\mathbb{Q}}_{\ell}|_{\underline{\mathcal{A}}^\dagger \times_{\underline{\mathcal{A}}} \underline{\mathcal{A}}^{\star, \text{reg}}} \right).$$

The group scheme $(\varphi^{\star})^* \underline{\mathcal{P}}$ acts on $\underline{\mathcal{M}}^{\star, \text{reg}}$ over the regular base $\underline{\mathcal{A}}^{\star, \text{reg}}$ and hence on the sheaf $R^{2d} f_{\dagger}^{\star, \text{reg}} \overline{\mathbb{Q}}_{\ell}$. This gives an action of $\underline{\mathcal{P}}$ on $\varphi_{\star}^{\star} (R^{2d} f_{\dagger}^{\star, \text{reg}} \overline{\mathbb{Q}}_{\ell}|_{\underline{\mathcal{A}}^\dagger})$ via the unit $\underline{\mathcal{P}} \rightarrow \varphi_{\star}^{\star} \varphi^{\star} \underline{\mathcal{P}}$. With this action, the isomorphism (8.2) intertwines the action of $\underline{\mathcal{P}}$. In particular, we have an equality of stable components

$$(R^{2d} f_{\star}^{\star, \dagger} \overline{\mathbb{Q}}_{\ell})_{\text{st}} \simeq \varphi_{\star}^{\star} ((R^{2d} f_{\dagger}^{\star, \text{reg}} \overline{\mathbb{Q}}_{\ell}|_{\underline{\mathcal{A}}^\dagger})_{\text{st}})$$

As $\underline{\mathcal{M}}^{\star, \text{reg}}$ is a torsor for the action of $\underline{\mathcal{P}} \times_{\underline{\mathcal{A}}} \underline{\mathcal{A}}^{\star, \text{reg}}$, the complex $(R^{2d} f_{\dagger}^{\star, \text{reg}} \overline{\mathbb{Q}}_{\ell}|_{\underline{\mathcal{A}}^\dagger})_{\text{st}}$ is a local system on $\underline{\mathcal{A}}^{\star, \text{reg}} \times_{\underline{\mathcal{A}}} \underline{\mathcal{A}}^\dagger$. Finally, since $\varphi^{\star}$ is finite étale (see Proposition 5.17), the right-hand side is also a local system on $\underline{\mathcal{A}}^\dagger$, and the proposition follows. $\square$

Corollary 8.8. *Fix an integer δ and choose a divisor D such that $\delta^{\text{bad}}(D) > \delta$ and $\deg(D) > g + \frac{3}{2}\delta$. Then, any simple constituent of the stable cohomology ${}^p\mathbf{H}^n(f_{\star}^{\star, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\text{st}}$ whose support intersects $\underline{\mathcal{A}}_{<\delta}$ has full support.*

Proof. Let Z be the support of a simple constituent of ${}^p\mathbf{H}^n(f_{\star}^{\star, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\text{st}}$ which intersects $\underline{\mathcal{A}}_{<\delta}$. By Theorem 8.3(1) combined with δ -regularity on $\underline{\mathcal{A}}_{<\delta} \subset \underline{\mathcal{A}}^{\text{good}}$, Z must be the closure of a component of $\underline{\mathcal{A}}_{\delta_Z}$. The claim is that $\delta_Z = 0$.

To see this, consider the open set $\underline{\mathcal{A}}^\dagger \subset \underline{\mathcal{A}}$. By Proposition 6.33 and the assumption on $\deg(D)$, $\underline{\mathcal{A}}^\dagger \cap \underline{\mathcal{A}}_{\delta}$ is dense in each component of $\underline{\mathcal{A}}_{\delta}$. Moreover, Proposition 8.7 shows that the top ordinary cohomology $(R^{2d} f_{\star}^{\star, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\text{st}}|_{\underline{\mathcal{A}}^\dagger}$ is a local system. Now by condition (2) of Theorem 8.3, there exists some $U \subset \underline{\mathcal{A}}^\dagger$ such that $U \cap Z \neq \emptyset$ and the local system $(R^{2d} f_{\star}^{\star, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\text{st}}|_{\underline{\mathcal{A}}^\dagger}$ admits a nontrivial direct factor supported in $U \cap \underline{\mathcal{A}}_{\delta}$. This is clearly a contradiction if $\delta \geq 1$, forcing Z to be the closure of $\underline{\mathcal{A}}_0$, which is the whole of $\underline{\mathcal{A}}^{\text{ani}}$. $\square$

We now turn to κ -isotypic components in the case $\star = \mathbf{S}$. Recall from Proposition 5.27 that for a 2-torsion element $\kappa \in \widehat{\mathbf{C}}$, we have a closed set $\underline{\mathcal{A}}_{\kappa} = \nu_{\mathfrak{e}}(\underline{\mathcal{A}}_{\mathfrak{e}})$, where $\mathfrak{e}$ is the (unique) pointed elliptic endoscopic datum containing κ , which is the support of the sheaf $(Rg_{\dagger}^{\text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}$. We begin with a lemma about the density of the dagger locus in $\underline{\mathcal{A}}_{\kappa}$.

Lemma 8.9. *Assume that $\deg(D) > 2g$. The subset $\underline{\mathcal{A}}_{\kappa}^{\dagger} := \underline{\mathcal{A}}_{\kappa} \cap \underline{\mathcal{A}}^{\dagger}$ is open and dense in every component of $\underline{\mathcal{A}}_{\kappa}$.*

Proof. By Lemma 2.21, we have $\nu_{\mathfrak{e}}(\underline{\mathcal{A}}_{\mathfrak{e}}) \cap \underline{\mathcal{A}}^{\dagger} = \nu_{\mathfrak{e}}(\underline{\mathcal{A}}_{\mathfrak{e}}^{\dagger})$. By Lemma 5.12, $\underline{\mathcal{A}}_{\mathfrak{e}}^{\dagger}$ is open and dense in $\underline{\mathcal{A}}_{\mathfrak{e}}$, so since $\nu_{\mathfrak{e}}$ is a closed embedding, we conclude that $\underline{\mathcal{A}}_{\kappa}^{\dagger}$ is dense in each component of $\underline{\mathcal{A}}_{\kappa}$. $\square$

In this case, the crucial calculation needed to apply the support theorem is the following.

Proposition 8.10. *The sheaf $(Rf_{\ast}^{\mathbf{S}, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}$ has support contained in $\mathcal{A}_{\kappa}$. Moreover, the top degree cohomology $(R^{2d} f_{\ast}^{\mathbf{S}, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}$ is a local system when restricted to the locally closed set $\mathcal{A}_{\kappa}^{\dagger}$.*

Proof. By Proposition 5.24, the action of $\mathbb{X}_{\ast}(\mathbf{C}) \times \mathcal{A}^{\text{ani}}$ on $Rf_{\ast}^{\mathbf{S}, \text{ani}} \overline{\mathbb{Q}}_{\ell}$ is through $\pi_0(\mathcal{P}/\mathcal{A}^{\text{ani}})$. We have a $\pi_0(\mathcal{P}/\mathcal{A}^{\text{ani}})$ equivariant isomorphism

$$R^{2d} \underline{g}_!^{\text{ani}} \overline{\mathbb{Q}}_{\ell} \simeq \overline{\mathbb{Q}}_{\ell}[\pi_0(\mathcal{P}/\mathcal{A}^{\text{ani}})](-d),$$

where $(-d)$ denotes Tate twist. Therefore, by Proposition 5.27, $\overline{\mathbb{Q}}_{\ell}[\pi_0(\mathcal{P}/\mathcal{A}^{\text{ani}})]_{\kappa}$ is supported on $\mathcal{A}_{\kappa}$, and hence, the support of $(Rf_{\ast}^{\mathbf{S}, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}$ is contained in $\mathcal{A}_{\kappa}$. Over $\mathcal{A}_{\kappa}^{\dagger}$, an identical argument to that of Proposition 8.7 works: Namely, we use the density of the regular locus in Hitchin fibers over $\mathcal{A}^{\dagger}$ (Lemma 8.6) to reduce to the study of the top degree cohomology over the regular locus. Over this space, we use the factorization

$$\mathcal{M}^{\mathbf{S}, \text{reg}} \xrightarrow{f^{\mathbf{S}, \text{reg}}} \mathcal{A}^{\mathbf{S}, \text{reg}} \xrightarrow{\varphi^{\mathbf{S}}} \mathcal{A}$$

from Proposition 5.15, noting that the κ -isotypic piece of top degree cohomology is then a pushforward $\varphi_{\ast}^{\mathbf{S}}(\mathcal{L}_{\kappa})$, where by Proposition 5.27, $\mathcal{L}_{\kappa}$ is a local system on $(\varphi^{\mathbf{S}})^{-1}(\mathcal{A}_{\kappa}^{\dagger})$ determined by the isomorphism class of the $\varphi^{\mathbf{S}, \ast}(\mathcal{P})$ -torsor $\mathcal{M}^{\mathbf{S}, \text{reg}}$. $\square$

Corollary 8.11. *Fix an integer δ , and choose the divisor D on Σ so that $\deg(D) > g + \frac{3}{2}\delta$ and $\delta^{\text{bad}}(D) > \delta$. Let Z be the support of an irreducible summand of ${}^p\mathbf{H}^n(f_{\ast}^{\mathbf{S}, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}$. If $Z \cap \nu_{\epsilon}(\mathcal{A}_{\epsilon, < \delta}) \neq \emptyset$, then $Z = \nu_{\epsilon, \ast}(\mathcal{A}_{\epsilon}^{\text{ani}})$ is the image of the transfer map for the unique pointed *still there?* endoscopic datum $\epsilon = (\kappa, \rho)$ extending κ .*

Proof. First, we relate δ -regularity in $\mathcal{A}_{\epsilon}$ to δ -regularity in $\mathcal{A}$. Suppose that $Z \subset \mathcal{A}_{\kappa}$ has preimage $Z_{\epsilon} = \nu_{\epsilon}^{-1}(Z)$ which satisfies the δ -regularity condition

$$\text{codim}_{\mathcal{A}_{\epsilon}}(Z_{\epsilon}) \geq \delta_{\epsilon, Z_{\epsilon}}.$$

Recall the invariant $r_v(D) = \sum_{v \in |\Sigma|} \deg(v) r_v(a_{\epsilon})$ from Corollary 6.32. Then, as ν_{ϵ} is finite, we compute

$$\text{codim}_{\mathcal{A}}(Z) = \text{codim}_{\mathcal{A}_{\epsilon}}(Z_{\epsilon}) + r_v(D) \geq \delta_{\epsilon, Z_{\epsilon}} + r_v(D) = \delta_Z$$

where the last equality follows from Corollary 6.32. Hence, $\nu_{\epsilon}(\mathcal{A}_{\epsilon}^{\text{good}}) \subset \mathcal{A}^{\text{good}}$, and we can apply Theorem 8.3.

The result now follows from Proposition 8.10 by an identical argument to that of Corollary 8.8, the key ingredient being Proposition 6.33, which shows in particular that, given the assumptions on $\deg(D)$, $\mathcal{A}_{\kappa} \cap \mathcal{A}_{\delta}^{\dagger} \subset \mathcal{A}_{\kappa} \cap \mathcal{A}_{\delta}$ is open and dense. $\square$

9. PROOF OF THE MAIN THEOREMS

In this section, we prove the main results of the paper, beginning with the geometric stabilization over the locus $\mathcal{A}^{\text{good}}$ (Theorem 9.1), and then using this result to obtain the local geometric statement Theorem 7.9.

9.1. Global Stabilization. We prove the main global result:

Theorem 9.1. *Fix a divisor D on Σ , and put $\delta = \min\{\delta^{\text{bad}}(D), \frac{2}{3}(\deg(D) - g)\}$; we choose D so that $\delta > 0$. For any 2-torsion element $\kappa \in \widehat{\mathbf{C}}[2]$, we have an isomorphism of complexes in $D^b(\mathcal{A}_{\epsilon, < \delta})$ over $\bar{k}$*

$$\nu_{\epsilon}^*(Rf_{\ast}^{\mathbf{S}, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}[2r_{\epsilon}(D)](r_{\epsilon}(D)) = (Rf_{\ast}^{\epsilon, \text{ani}} \overline{\mathbb{Q}}_{\ell})_{\text{st}}$$

(restricted to the locus $\mathcal{A}_{\epsilon, < \delta}$) which induces an isomorphism of semi-simplifications over k .

Proof. Since $f^{\mathbf{S}}$ and $f^{\mathbf{e}}$ are proper over $\mathcal{A}^{\text{ani}}$ and the spaces $\mathcal{M}^{\mathbf{S},\text{ani}}, \mathcal{M}^{\mathbf{e},\text{ani}}$ are smooth by Proposition 5.8, both sides of the proposed identity are pure and hence split as a direct sum of their perverse cohomology sheaves. The theorem reduces to the identity

$$\bigoplus_n \nu_{\mathbf{e}}^*({}^p\mathbf{H}^n(\mathbf{R}f_{\mathbf{e}}^{\mathbf{S},\text{ani}}\overline{\mathbb{Q}}_{\ell})_{\kappa}[2r_{\mathbf{e}}(D)](r_{\mathbf{e}}(D)))[-n] = \bigoplus_n {}^p\mathbf{H}^n(\mathbf{R}f_{\mathbf{e}}^{\mathbf{e},\text{ani}}\overline{\mathbb{Q}}_{\ell})_{\text{st}}[-n]$$

Moreover, by the decomposition theorem of Beilinson, Bernstein, Deligne, and Gabber [?], the perverse cohomology groups on either side are semisimple. By Corollary 8.11 and our assumption on $\deg(D)$, it is enough to show there is an isomorphism after restricting both sides to an open set $\mathcal{U} \subset \mathcal{A}_{\mathbf{e}}^{\dagger}$, where both sides are local systems. Namely, let $\mathcal{U}$ denote the open subset of $\mathcal{A}_{\mathbf{e}}^{\dagger}$ whose $\bar{k}$ -points are given by $a \in \mathcal{A}_{\mathbf{e}}^{\dagger}(\bar{k})$ for which $a(\Sigma)$ intersects $\mathfrak{D}_{\mathbf{e}}^{\text{cent}}$ transversely, and $a^{-1}(\mathfrak{D}_{\mathbf{e}}^{\text{cent}})$ and $a^{-1}(\mathfrak{D}_{\mathbf{e}}^{\text{br}})$ are disjoint divisors in Σ . Both sides are local systems on $\mathcal{A}_{\mathbf{e}}^{\dagger}$ by Propositions 8.7 and 8.10.

Denote

$$L_{\kappa}^n = \nu_{\mathbf{e}}^*({}^p\mathbf{H}^n(\mathbf{R}f_{\mathbf{e}}^{\mathbf{S}}\overline{\mathbb{Q}}_{\ell})_{\kappa}[2r_{\mathbf{e}}(D)](r_{\mathbf{e}}(D)))|_{\mathcal{U}} \quad \text{and} \quad L_{\mathbf{e},\text{st}}^n = {}^p\mathbf{H}^{n-2r_{\mathbf{e}}(D)}(\mathbf{R}f_{\mathbf{e}}^{\mathbf{e}}\overline{\mathbb{Q}}_{\ell})_{\text{st}}|_{\mathcal{U}}$$

which are local systems by Propositions 8.7 and 8.11. By Chebotarev density theorem and purity of the pushforwards $\mathbf{R}f_{\mathbf{e}}^{\mathbf{S},\text{ani}}\overline{\mathbb{Q}}_{\ell}$, it suffices to show that for all finite extensions K over k with Frobenius σ_K and for all $a \in \mathcal{U}(K)$,

$$(9.1) \quad q^{-r_{\mathbf{e}}(D) \cdot [K:k]} \cdot \sum_n (-1)^n \text{tr}(\sigma_K, (L_{\kappa}^n)_a) = \sum_n (-1)^n \text{tr}(\sigma_K, (L_{\mathbf{e},\text{st}}^n)_a)$$

Without loss of generality, we assume that $K = k$. Choose a smooth group scheme J'_a with connected fibers equipped with a map $J'_a \rightarrow J_a$ which is an isomorphism on generic fibers, and let $\mathcal{P}'_a$ be the commutative group stack classifying J'_a -torsors on $\mathcal{A} \times \Sigma$. Recall that $U_a \subset \Sigma$ denotes the preimage of $\mathfrak{c}_D^{\text{rs}}$ under a . Then, using Corollary 7.8, we compute that the left hand side of (9.1) is

$$\#(\mathcal{P}'_a)^{\circ}(k) \prod_{v \in |\Sigma - U_a|} \#[\mathcal{M}_v^{\mathbf{S}}(a_v)/\mathcal{P}_v(J'_{a_v})](k_v)_{\kappa},$$

while the right hand side is

$$q^{r_{\mathbf{e}}(D)} \cdot \#(\mathcal{P}'_a)^{\circ}(k) \prod_{v \in |\Sigma - U_a|} \#[\mathcal{M}_v^{\mathbf{e}}(a_{\mathbf{e},v})/\mathcal{P}_v(J'_a)](k_v)_{\text{st}}.$$

Note that the extra power of q in the right hand side is from the Tate twist. Since $r_{\mathbf{e}}(D) = \sum_v \deg(v)r_{\mathbf{e},v}(a_{\mathbf{e}})$ can be computed locally, the identity will therefore follow from the cases of Theorem 7.9 given in Propositions 7.11, 7.13, and 7.14; namely, those $a_{\mathbf{e}} \in \mathfrak{c}_{\mathbf{e}}^{\heartsuit}(\mathcal{O}_v)$ and $a \in \mathfrak{c}^{\heartsuit}(\mathcal{O}_v)$ which intersect the divisor $\mathfrak{D}^{\text{br}}$ transversely but do not intersect $\mathfrak{D}^{\text{cent}}$ ($d_v^{\text{br}}(a) \leq 1$ for all v and $d_v^{\text{cent}}(a) = d_v^{\text{cent}}(a_{\mathbf{e}}) = 0$) and those $a_{\mathbf{e}}, a$ which do not intersect the divisor $\mathfrak{D}^{\text{br}}$ and for which $d_v^{\text{cent}}(a_{\mathbf{e}}) \leq 1$. Theorem 9.1 therefore follows. $\square$

9.2. Proof of Theorem 7.9. In this section, we prove Theorem 7.9, which immediately implies Theorem 7.1 via Corollary 7.6.

Fix a place $v \in |\Sigma|$ with residue field $k_v = k$ such that there is a unique place $v' \in |\Sigma'|$ over v . Let $\mathbf{e} = (\kappa, \rho)$ be an endoscopic datum over $F_v = k((\varpi_v))$ and $a_{\mathbf{e},v} \in \mathfrak{c}_{\mathbf{e}}^{\heartsuit}(\mathcal{O}_v)$ with image under the transfer map $\nu_{\mathbf{e}}$ given by a_v . We wish to prove Theorem 7.9 for the pair $a_{\mathbf{e},v}, a_v$. By Proposition 7.11, we can assume that $\text{Ob}_v(a_{\mathbf{e},v}) = 0$. Fix $N \gg 0$ such that Proposition 6.26 holds for both a_v and $a_{\mathbf{e},v}$, and fix a line bundle D on Σ with

$$\deg(D) \geq 2g + nN + 1$$

Proposition 9.2. *There exists an integer N' such that, for all finite extensions K of k of degree greater than N' , there exists a point $a_{\mathbf{e}} \in \mathcal{A}_{\mathbf{e}}(K)$ such that*

$$(1) \ a_{\mathbf{e}} \equiv a_{\mathbf{e},v} \pmod{\varpi_v^N};$$

- (2) the support of the divisors $a_{\mathfrak{c}}^{-1}(\mathfrak{D}_D^{\text{cent}})$ and $a_{\mathfrak{c}}^{-1}(\mathfrak{D}_D^{\text{br}})$ intersects non-trivially at most at the place v in Σ ;
- (3) $\text{Ob}_u(a_{\mathfrak{c}})$ is trivial for all places $u \in |\Sigma|$;
- (4) $a_{\mathfrak{c}} \in \mathcal{A}_{\mathfrak{c}}^{\text{ani}}(K)$;
- (5) $\delta_{\mathfrak{c}}(a_{\mathfrak{c}}) = \delta_{\mathfrak{c},v}(a_{\mathfrak{c},v})$.

Proof. We recall how the obstruction is computed on the endoscopic base. We denote ranks of symmetric varieties by subscripts. Write $X^{\mathfrak{c}} = X_r^{\mathbf{S}} \times X_s^{\mathbf{U}}$ with $r + s = n$, so that $\mathcal{A}_{\mathfrak{c}} = \mathcal{A}_r \times \mathcal{A}_s$, where the first factor is the Hitchin base of the Siegel factor and the second that of the unitary factor. By Lemma 6.12, for every non-split place u we have

$$\text{Ob}_u(a_{\mathfrak{c}}) = \text{val}_u((a_{\mathfrak{c}}^{\mathbf{U}})_{2s}) \bmod 2.$$

where $a_{\mathfrak{c}} = (a_{\mathfrak{c}}^{\mathbf{S}}, a_{\mathfrak{c}}^{\mathbf{U}})$ and $(a_{\mathfrak{c}}^{\mathbf{U}})_{2s}$ is the determinant coordinate of $a_{\mathfrak{c}}^{\mathbf{U}}$. In particular, condition (3) holds for *every* place u whenever $a_{2s}^{\mathbf{U}}$ is in the image of the norm map

$$(9.2) \quad \text{Nm}: H^0(\Sigma', \vartheta^* D^{\otimes s}) \rightarrow H^0(\Sigma, D^{\otimes 2s}), \quad t \mapsto t \cdot \sigma(t),$$

which motivates the following definition.

Definition 9.3. Let $\text{Nm} \subset H^0(\Sigma, D^{\otimes 2s})$ denote the image of the norm map (9.2). We define the *norm locus*

$$\mathcal{A}_s^{\text{Nm}} := \{ (a_2^{\mathbf{U}}, \dots, a_{2s}^{\mathbf{U}}) \mid a_{2s}^{\mathbf{U}} \in \text{Nm} \} \subset \mathcal{A}_s$$

and $Q := \mathcal{A}_r \times \mathcal{A}_s^{\text{Nm}} \subset \mathcal{A}_{\mathfrak{c}}$.

Lemma 9.4. Let $Q^{\text{ani}} = Q \cap \mathcal{A}_{\mathfrak{c}}^{\text{ani}}$. The codimension of $Q \setminus Q^{\text{ani}}$ in Q is greater than or equal to $\deg(D) + 1 - g$.

Proof. Note that if $X^{\mathfrak{c}} = X^{\mathbf{S}}$, then we are done by Lemma 5.9. Therefore, as $X^{\mathfrak{c}}$ splits as a product and Q only imposes conditions on the unitary factor, it suffices to prove the lemma for $X^{\mathfrak{c}} = X^{\mathbf{U}}$. We will drop subscripts $\mathfrak{c}$ and work with $X^{\mathbf{U}}$ throughout. Recall from §3 that we denote the Hitchin base for the unitary group U_{2n} by $\mathcal{B} = \mathcal{B}_{2n}$. We have a natural inclusion $\mathcal{A} \rightarrow \mathcal{B}$ which identifies $\mathcal{A}$ with the fixed points of the involution $\theta(a_i) = (-1)^i a_i$ on $\mathcal{B}$. By definition of the anisotropic locus, $\mathcal{A}^{\text{ani}} = \mathcal{A} \cap \mathcal{B}^{\text{ani}}$.

Fix a θ - and σ -stable torus T of U_{2n} which is maximally θ -split; that is, for which the dimension of the subtorus on which θ acts by inversion is maximal. Note that there is a unique $H^{\mathbf{U}}(F)$ -conjugacy class of such tori. For every Levi subgroup M of U_{2n} containing T , we associate a subvariety $\mathcal{B}_M \subset \mathcal{B}$ consisting of those points in $\mathcal{B} = H^0(\Sigma, (\mathfrak{u}_{2n} // \text{U}_{2n})_{\mathcal{L}(D)})$ which factor through $\mathfrak{m} // M \rightarrow \mathfrak{u}_{2n} // \text{U}_{2n}$. We note that the subvarieties $\mathcal{B}_M$ are independent of conjugacy class of M in U_{2n} . Since U_{2n} is a quasisplit group scheme over Σ , [Ngo10, Proposition 6.3.5] implies that

$$\mathcal{B} \setminus \mathcal{B}^{\text{ani}} = \bigcup_{T \subset M \subset \text{U}_{2n}} \mathcal{B}_M$$

where the union ranges over all proper Levi subgroups M containing T .

Put $\mathcal{A}_M = \mathcal{A} \cap \mathcal{B}_M$. It now suffices to show $\text{codim}_Q(Q \cap \mathcal{A}_M) > \deg(D) + 1 - g$. All Levi subgroups M are of the form

$$M = \text{U}_{2m_0} \oplus \prod_{i=1}^{\ell} \text{Res}_{\Sigma'/\Sigma}(\text{GL}_{m_i})$$

where $\sum_{i=0}^{\ell} m_i = n$. Note in particular, $\mathcal{A}_M$ is the image of a product of Hitchin bases for $U_{2m_0}/(U_{m_0} \times U_{m_0})$ and $\text{Res}_{\Sigma'/\Sigma}(\text{GL}_{m_i})/U_{m_i}$. In particular, using Riemann-Roch, one can compute

$$\begin{aligned} \dim(\mathcal{A}_M) &= m_0(m_0 + 1) \deg(D) + m_0(1 - g) + \sum_{i=1}^{\ell} \left[\frac{m_i(m_i + 1)}{2} \deg(D) + m_i(1 - g) \right] \\ &= \left[m_0(m_0 + 1) + \sum_{i=1}^{\ell} \frac{m_i(m_i + 1)}{2} \right] \deg(D) + n(1 - g). \end{aligned}$$

It is easy to check that, for fixed ℓ , the above quantity is maximized when $m_0 = n - \ell$ and $m_i = 1$ for all $i \geq 1$, in which case we obtain

$$(9.3) \quad \dim(\mathcal{A}_M) \leq \left[(n - \ell)^2 + n \right] \deg(D) + n(1 - g).$$

We also compute by Riemann-Roch that

$$(9.4) \quad \dim(Q) = \sum_{i=1}^{n-1} (2id + 1 - g) + 2nd + 2(1 - g) = n(n + 1) \deg(D) + (n + 1)(1 - g).$$

Hence, we have

$$\begin{aligned} \text{codim}_Q(Q \cap \mathcal{A}_M) &= \dim(Q) - \dim(\mathcal{A}_M \cap Q) \geq \dim(Q) - \dim(\mathcal{A}_M) \\ &= [2\ell n - \ell^2] \deg(D) + 1 - g \geq n^2 \deg(D) + 1 - g \\ &\geq \deg(D) + 1 - g \end{aligned}$$

where the second to last inequality follows since $2\ell n - \ell^2$ is maximized for $1 \leq \ell \leq n$ when $\ell = n$. $\square$

Now, again by Riemann–Roch and our assumption on $\deg(D)$, we have a surjective evaluation map

$$(9.5) \quad \mathcal{A}_{\mathfrak{c}} = H^0(\Sigma, \mathfrak{c}_{\mathfrak{c}, D}) \rightarrow H^0(\Sigma_v, \mathfrak{c}_{\mathfrak{c}}(\mathcal{O}_v/\varpi_v^N))$$

Let Z be the preimage of $a_{\mathfrak{c}, v}$, where we view $a_{\mathfrak{c}, v}$ as an element in $H^0(\Sigma_v, \mathfrak{c}(\mathcal{O}_v/\varpi_v^N))$. By surjectivity of the linear map (9.5), the codimension of Z in $\mathcal{A}$ is nN . As a consequence, we have

$$\text{codim}_Q(Q \cap Z) \leq nN.$$

We now impose condition (5) by using a transversality argument. Namely, let $(Q \cap Z)^{\min} \subset Q \cap Z$ denote the subvariety of points a satisfying $d_{\mathfrak{c}, u}^{\text{cent}}(a) \leq 1$ for all $u \neq v$. By an identical argument to Corollary 6.38, to prove $(Q \cap Z)^{\min}$ contains an open, dense subset of $Q \cap Z$, it suffices to prove the following.

Lemma 9.5. *Fix a place $u \neq v$, and let $Q_u^{(j)} \subset H^0(\Sigma_u, \mathfrak{c}(\mathcal{O}_u/\varpi_u^j))$ denote the subset of tuples $(a_2, \dots, a_{2n}) \in \mathfrak{c}(\mathcal{O}_u/\varpi_u^j)$ such that $a_{2n} \in \mathcal{O}_u/\varpi_u^j$ is the norm of an element in $\mathcal{O}_{u'}/\varpi_u^j$. With the assumptions on $\deg(D)$ above, the product of local evaluation maps at v and u gives a surjection*

$$Q \rightarrow Q_v^{(N)} \times Q_u^{(2)} \subset H^0(\Sigma_v, \mathfrak{c}(\mathcal{O}_v/\varpi_v^N)) \times H^0(\Sigma_u, \mathfrak{c}(\mathcal{O}_u/\varpi_u)).$$

Proof. The map $Q \rightarrow Q_v^{(N)} \times Q_u^{(2)}$ is identified with the map on H^0 induced by map

$$0 \rightarrow \left(\vartheta_* \vartheta^* D^{\otimes n} \oplus \bigoplus_{i=1}^{n-1} D^{\otimes 2i} \right) \otimes \mathcal{O}(-Nv - 2u) \rightarrow D^{\otimes n} \oplus \bigoplus_{i=1}^{n-1} D^{\otimes 2i} \rightarrow D^{\otimes n} \oplus \bigoplus_{i=1}^{n-1} D^{\otimes 2i} \Big|_{Nv+2u} \rightarrow 0$$

so the cokernel is

$$H^1\left(\Sigma, \left(D^{\otimes n} \oplus \bigoplus_{i=1}^{n-1} D^{\otimes 2i} \right) \otimes \mathcal{O}(-Nv - 2u)\right),$$

which vanishes for $\deg(D) \geq 2g + nN + 1$. $\square$

We now have

$$\mathrm{codim}_Q(Q \cap Z)^{\min} \leq nN.$$

In particular, by our assumptions on the degree of D and Lemma 9.4, we have

$$\mathrm{codim}_Q(Q \cap Z) < \mathrm{codim}_Q(Q \setminus Q^{\mathrm{ani}})$$

so that the locally closed set $Q^{\mathrm{ani}} \cap (Q \cap Z)^{\min}$ is nonempty. Since this is a nonempty locally closed subvariety, by Lang-Weil there is an integer N' such that it has K points for any finite extension K of k with $[K : k] \geq N'$. In particular, if we choose any $a_{\mathfrak{e}} \in (Q^{\mathrm{ani}} \cap (Q \cap Z)^{\min})(K)$, then by Proposition 6.26, its local δ invariant at v agrees with $\delta_{\mathfrak{e},v}(a_{\mathfrak{e},v})$ and by Proposition 6.28 transversality implies $\delta_{\mathfrak{e},u}(a_{\mathfrak{e},u}) = 0$. Hence, $a_{\mathfrak{e}}$ satisfies condition (5). The other conditions are clear from our choice, so $a_{\mathfrak{e}}$ is a K point satisfying the assumptions of the proposition. $\square$

Let $a_{\mathfrak{e}}$ be as in Proposition 9.2, and $a = \nu_{\mathfrak{e}}(a_{\mathfrak{e}})$. Let $\underline{a}, \underline{a}_{\mathfrak{e}}$ be lifts in $\underline{\mathcal{A}}^{\mathrm{ani}}(K)$, respectively $\underline{\mathcal{A}}_{\mathfrak{e}}^{\mathrm{ani}}(K)$. By condition (5), we may also assume that $\deg(D) > g + \frac{3}{2}\delta_{\mathfrak{e}}(a_{\mathfrak{e}})$ and that $\deg(D)$ is large enough so that

$$\delta_{\mathfrak{e}}(a_{\mathfrak{e}}) = \delta_{\mathfrak{e},v}(a_{\mathfrak{e},v}) < \delta^{\mathrm{bad}}(D)$$

Hence, we assume that $\underline{\mathcal{A}}_{\mathfrak{e}, < \delta(a)} \subset \underline{\mathcal{A}}_{\mathfrak{e}}^{\mathrm{good}}$. Therefore, Theorem 9.1 holds over $\underline{\mathcal{A}}_{\mathfrak{e}, < \delta_{\mathfrak{e}}(a_{\mathfrak{e}})}$. Then, by the point counting formula for the product formula (Proposition 7.7) combined with Theorem 9.1, we have

$$(9.6) \quad \#(\mathcal{P}'_a)^{\circ}(K) \cdot \prod_{v \in |\Sigma_K - U_a|} \#[\mathcal{M}_v^{\mathbf{S}}(a_v)/\mathcal{P}(J'_{a_v})](K_v)_{\kappa} = q^{r_{\mathfrak{e}}(D)[K:k]} \cdot \#(\mathcal{P}'_a)^{\circ}(K) \cdot \prod_{v \in |\Sigma_K - U_a|} \#[\mathcal{M}_v^{\mathfrak{e}}(a_{\mathfrak{e},v})/\mathcal{P}(J'_{a_v})](K_v).$$

Note here that the extra power of q appears in the Tate twist in Theorem 9.1. Moreover, by the cases in Propositions 7.13 and 7.14, for each of the places $u \neq v$ of Σ , we have identities

$$\#[\mathcal{M}_u^{\mathbf{S}}(a_u)/\mathcal{P}_u(J'_{a_u})](K_u)_{\kappa} = q^{r_{\mathfrak{e}}(a_{\mathfrak{e},u}) \cdot [K:k]} \#[\mathcal{M}_u^{\mathfrak{e}}(a_{\mathfrak{e},u})/\mathcal{P}_u(J'_{a_{\mathfrak{e},u}})](K_u) \neq 0.$$

By canceling these terms from (9.6), we get the identity

$$\#[\mathcal{M}_v^{\mathbf{S}}(a_v)/\mathcal{P}_v(J'_{a_v})](K)_{\kappa} = q^{r_{\mathfrak{e}}(a_{\mathfrak{e},v}) \cdot [K:k]} \#[\mathcal{M}_v^{\mathfrak{e}}(a_{\mathfrak{e},v})/\mathcal{P}_v(J'_{a_v})](K).$$

Since this holds for every finite extension K of k of degree greater than N' , we can conclude the same for k , which proves Theorem 7.9. $\square$

It is now simple to deduce Theorem 9.1 over $\underline{\mathcal{A}}_{\mathfrak{e}}^{\mathrm{ani}}$ (without the restriction to $\underline{\mathcal{A}}^{\mathrm{good}}$) by reversing the flow of information from local back to global.

Theorem 9.6. *Fix any divisor D on Σ with $\deg(D) \geq 2g$. For any 2-torsion element $\kappa \in \widehat{\mathbf{C}}[2]$, we have an isomorphism of complexes in $D^b(\underline{\mathcal{A}}_{\mathfrak{e}}^{\mathrm{ani}})$ over $\bar{k}$*

$$\nu_{\mathfrak{e}}^*(\mathrm{R}f_{\ast}^{\mathbf{S}, \mathrm{ani}} \overline{\mathbb{Q}}_{\ell})_{\kappa}[2r_{\mathfrak{e}}(D)](r_{\mathfrak{e}}(D)) = (\mathrm{R}f_{\ast}^{\mathfrak{e}, \mathrm{ani}} \overline{\mathbb{Q}}_{\ell})_{\mathrm{st}}$$

which induces an isomorphism of semisimplifications over k .

Proof. By an argument identical to Theorem 9.1, using Chebotarev density theorem and Corollary 7.8, it suffices to prove Theorem 7.9. Having proved this, the global isomorphism follows on all of $\underline{\mathcal{A}}^{\mathrm{ani}}$. $\square$

REFERENCES

- [Art69] M. Artin, *Algebraic approximation of structures over complete local rings*, Inst. Hautes Études Sci. Publ. Math. (1969), no. 36, 23–58. MR 268188
- [Bez96] Roman Bezrukavnikov, *The dimension of the fixed point set on affine flag manifolds*, Mathematische Research Letters **3** (1996), no. 2, 185–189.
- [BL95] Arnaud Beauville and Yves Laszlo, *Un lemme de descente*, C. R. Acad. Sci. Paris Sér. I Math. **320** (1995), no. 3, 335–340. MR 1320381

- [BZSV23] David Ben-Zvi, Yiannis Sakellaridis, and Akshay Venkatesh, *Relative Langlands duality*, arXiv preprint, arXiv:2409.04677 (2023).
- [Chi22] Jingren Chi, *Geometry of Kottwitz-Viehmann varieties*, J. Inst. Math. Jussieu **21** (2022), no. 1, 1–65. MR 4366333
- [FJ93] Solomon Friedberg and Hervé Jacquet, *Linear periods*, J. Reine Angew. Math. **443** (1993), 91–139. MR 1241129
- [GKM04] Mark Goresky, Robert Kottwitz, and Robert Macpherson, *Homology of affine Springer fibers in the unramified case*, Duke Math. J. **121** (2004), no. 3, 509–561. MR 2040285
- [GKM06] Mark Goresky, Robert Kottwitz, and Robert MacPherson, *Purity of equivalued affine Springer fibers*, Represent. Theory **10** (2006), 130–146. MR 2209851
- [GKM09] ———, *Codimensions of root valuation strata*, Pure Appl. Math. Q. **5** (2009), no. 4, 1253–1310. MR 2560317
- [GPPN23] Oscar García-Prada and Ana Peón-Nieto, *Abelianization of Higgs bundles for quasi-split real groups*, Transform. Groups **28** (2023), no. 1, 285–325. MR 4552177
- [GWZ20] Michael Groechenig, Dimitri Wyss, and Paul Ziegler, *Geometric stabilisation via p -adic integration*, J. Amer. Math. Soc. **33** (2020), no. 3, 807–873. MR 4127904
- [Hit87] Nigel Hitchin, *Stable bundles and integrable systems*, Duke Math. J. **54** (1987), no. 1, 91–114. MR 885778
- [HLM24] Thomas Hameister, Zhilin Luo, and Benedict Morrissey, *Relative Dolbeault geometric Langlands via the regular quotient*, 2024, arXiv preprint, arXiv:2409.15691.
- [HM25] Thomas Hameister and Benedict Morrissey, *The Hitchin fibration for symmetric pairs*, Advances in Mathematics **482** (2025), 110560.
- [JL85] H. Jacquet and K. F. Lai, *A relative trace formula*, Compositio Math. **54** (1985), no. 2, 243–310. MR 783512
- [KL88] David Kazhdan and George Lusztig, *Fixed point varieties on affine flag manifolds*, Israel Journal of Mathematics **62** (1988), no. 2, 129–168.
- [KR71] B. Kostant and S. Rallis, *Orbits and representations associated with symmetric spaces*, Amer. J. Math. **93** (1971), 753–809. MR 311837
- [Les21] Spencer Leslie, *An analogue of the Grothendieck-Springer resolution for symmetric spaces*, Algebra Number Theory **15** (2021), no. 1, 69–107. MR 4226983
- [Les22] ———, *On the stabilization of relative trace formulae: descent and the fundamental lemma*, Adv. Math. **394** (2022), Paper No. 108026, 68. MR 4355728
- [Les24] ———, *Symmetric varieties for endoscopic groups*, 2024, arXiv preprint, arXiv: 2401.09156.
- [Les25] ———, *The endoscopic fundamental lemma for unitary Friedberg–Jacquet periods*, Annals of Mathematics **201** (2025), no. 3, 551–645.
- [Lev07] Paul Levy, *Involutions of reductive Lie algebras in positive characteristic*, Adv. Math. **210** (2007), no. 2, 505–559. MR 2303231
- [LN08] Gérard Laumon and Bao Chau Ngo, *Le lemme fondamental pour les groupes unitaires*, Annals of Mathematics **168** (2008), no. 2, 477–573.
- [LXZ25a] Spencer Leslie, Jingwei Xiao, and Wei Zhang, *Unitary Friedberg–Jacquet periods and their twists: Fundamental lemmas*, arXiv preprint arXiv:2503.09500 (2025).
- [LXZ25b] ———, *Unitary Friedberg–Jacquet periods and their twists: Relative trace formulas*, arXiv preprint arXiv:2503.09664 (2025).
- [McN04] George J McNinch, *Nilpotent orbits over ground fields of good characteristic*, Mathematische Annalen **329** (2004), 49–85.
- [Ngo06] Bao Chau Ngo, *Fibration de Hitchin et endoscopie*, Invent. Math. **164** (2006), no. 2, 399–453. MR 2218781
- [Ngo10] ———, *Le lemme fondamental pour les algèbres de Lie*, Publ. Math. Inst. Hautes Études Sci. (2010), no. 111, 1–169. MR 2653248
- [Pol15] Jason K. C. Polák, *Dimensions of affine Springer fibers for some GL_n symmetric spaces*, 2015, <https://arxiv.org/abs/1506.04703>.
- [Rom05] Matthieu Romagny, *Group actions on stacks and applications*, Michigan Mathematical Journal **53** (2005), no. 1, 209–236.
- [Rom22] ———, *Algebraicity and smoothness of fixed point stacks*, 2022.
- [Sch13] Laura P. Schaposnik, *Spectral data for G -Higgs bundles*, Ph.D. thesis, 2013, Thesis (D.Phil.)—University of Oxford (United Kingdom), p. 1. MR 3389247
- [Sch15] ———, *Spectral data for $U(m, m)$ -Higgs bundles*, Int. Math. Res. Not. IMRN (2015), no. 11, 3486–3498. MR 3373057
- [Sch18a] ———, *A geometric approach to orthogonal Higgs bundles*, Eur. J. Math. **4** (2018), no. 4, 1390–1411. MR 3866701
- [Sch18b] ———, *An introduction to spectral data for Higgs bundles*, The geometry, topology and physics of moduli spaces of Higgs bundles, Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap., vol. 36, World Sci. Publ., Hackensack, NJ, 2018, pp. 65–101. MR 3837869

- [SV17] Yiannis Sakellaridis and Akshay Venkatesh, *Periods and harmonic analysis on spherical varieties*, Astérisque (2017), no. 396, viii+360. MR 3764130
- [Wan26] X. Griffin Wang, *Multiplicative Hitchin fibrations and the fundamental lemma*, Annals of Mathematics Studies, Princeton University Press, Princeton, NJ, 2026.
- [WZ24] X. Griffin Wang and Zhiyu Zhang, *Relative fundamental lemmas for spherical Hecke algebras and multiplicative Hitchin fibrations: the Jacquet–Rallis case*, 2024, Preprint.
- [Yun11] Zhiwei Yun, *The fundamental lemma of Jacquet and Rallis*, Duke Math. J. **156** (2011), no. 2, 167–227, With an appendix by Julia Gordon. MR 2769216
- [Yun16] ———, *Lectures on springer theories and orbital integrals*, arXiv preprint arXiv:1602.01451 (2016).