



UNSW
A U S T R A L I A



UNIVERSITY OF NEW SOUTH WALES

SCHOOL OF MATHEMATICS AND STATISTICS

Assignment 3

Functional Analysis

Author:
Edward McDonald

Student Number:
3375335

Question 1

For this question, X is a Banach space and X_0 is a closed subspace.

Proposition 1. *There is an isometric embedding, $X/X_0 \hookrightarrow (X_0^\perp)^*$.*

Proof. Define $\rho : X/X_0 \rightarrow (X_0^\perp)^*$ as $\rho(x + X_0)(f) = f(x)$, for $x \in X$ and $f \in X_0^\perp$. This is a well defined linear map, since if $x' + X_0 = x + X_0$, then $f(x) = f(x')$ since $x - x' \in X_0$ and $f \in X_0^\perp$. ρ is linear, since $\rho(\alpha x + y + X_0) = f(\alpha x + y) = \alpha f(x) + f(y)$ for $x, y \in X$ and $\alpha \in \mathbb{C}$.

We need to show that ρ is an isometry. That is,

$$\|\rho(x + X_0)\|_{(X_0^\perp)^*} = \|x + X_0\|_{X/X_0}.$$

for any $x \in X$.

Expanding this out into definitions, we must prove that

$$\sup_{\|f\|_{X^*} \leq 1, f \in X_0^\perp} |f(x)| = \inf_{y \in X_0} \|x - y\|. \quad (1)$$

for any $x \in X$.

Suppose that $f \in X_0^\perp$ with $\|f\|_{X^*} \leq 1$, then for any $y \in X_0$ and $x \in X$.

$$|f(x)| = |f(x - y)| \leq \|x - y\|.$$

Hence,

$$|f(x)| \leq \inf_{y \in X_0} \|x - y\|.$$

So

$$\sup_{\|f\|_{X^*} \leq 1, f \in X_0^\perp} |f(x)| \leq \inf_{y \in X_0} \|x - y\|$$

follows. Now we must prove the opposite inequality.

Let $x \in X$. Then on the subspace $V := \mathbb{C}x$, define the functional

$$\omega(\lambda x) = \lambda \|x + X_0\|_{X/X_0}.$$

ω is linear, and for any $y \in V$, $|\omega(y)| = \|y + X_0\| \leq \|y\|$. So $\|\omega\| \leq 1$.

So ω is a linear functional on a subspace V of X bounded by the seminorm $\|\cdot\|_{X/X_0}$.

So by the Hahn-Banach theorem, there is a functional $f \in X^*$, with $f(y) = \omega(y)$ for $y \in V$ and $|f(z)| \leq \|z + X_0\|$ for any $z \in X$. Hence for $z \in X_0$, $f(z) = 0$, so $f \in X_0^\perp$ and $|f(z)| \leq \|z\|$, so $\|f\|_{X^*} \leq 1$.

Therefore, $|f(x)| = \|x + X_0\|_{X/X_0}$. Hence,

$$\sup_{\|f\|_{X^*} \leq 1, f \in X_0^\perp} |f(x)| \geq \|x + X_0\|_{X/X_0} = \inf_{y \in X_0} \|x - y\|.$$

So the equality 1 holds. Hence ρ is an isometric embedding. \square

Question 2

Theorem 1. Suppose $T : X \rightarrow Y$ is a linear mapping between normed spaces X and Y . Then T is bounded if and only if T has the property that if U is open in Y then $T^{-1}(U)$ is open in X .

Proof. Suppose that T is bounded, and let $U \subset Y$ be an open set with $T^{-1}(U) \neq \emptyset$.

Then let $x \in T^{-1}(U)$, and let ε be small enough such that $B_Y(Tx, \varepsilon) \subset U$.

Then choose $\varepsilon' = \varepsilon/\|T\|$. Then if $y \in B_X(x, \varepsilon')$,

$$\|Ty - Tx\| \leq \|T\|\|y - x\| \leq \varepsilon.$$

Hence $Ty \in B_Y(Tx, \varepsilon)$, so $Ty \in U$. Therefore $B_X(x, \varepsilon') \subset T^{-1}(U)$, and so $T^{-1}(U)$ is open.

Conversely, suppose that T has the property that $T^{-1}(U)$ is open in X whenever U is open in Y .

Let $U = B_Y(0, 1) \subset Y$. Then since $T^{-1}(U)$ is open, there is some $\varepsilon > 0$ such that $B_X(0, \varepsilon) \subset T^{-1}(U)$.

So $TB_X(0, \varepsilon) \subseteq B_Y(0, 1)$. Hence, by linearity, $TB_X(0, 1) \subseteq B_Y(0, \frac{1}{\varepsilon})$.

Hence, $\|T\| \leq 1/\varepsilon$. So T is bounded. □

Question 3

Let $X = \ell^1(\mathbb{N})$, and X_0 is the subspace defined by

$$X_0 = \{(\xi_k)_{k \geq 0} : \sum_{k \geq 0} \xi_k = 0\}.$$

Theorem 2. There is an isometric isomorphism,

$$X/X_0 \cong \mathbb{C}$$

Proof. Define function $S : X/X_0 \rightarrow \mathbb{C}$ by

$$S((\xi_k)_{k \geq 0} + X_0) = \sum_{k \geq 0} \xi_k.$$

This is well defined, since if $(\xi'_k)_{k \geq 0} \in X_0$, then

$$S((\xi_k)_{k \geq 0} + (\xi'_k)_{k \geq 0}) = \sum_{k \geq 0} \xi_k + \xi'_k = \sum_{k \geq 0} \xi_k + \sum_{k \geq 0} \xi'_k = S((\xi_k)_{k \geq 0}),$$

since we may rearrange the sum as it is absolutely convergent. S is linear, since if $(\eta_k)_{k \geq 0}, (\xi_k)_{k \geq 0} \in X$, then

$$S((\xi_k)_{k \geq 0} + (\eta_k)_{k \geq 0}) = \sum_{k \geq 0} \xi_k + \eta_k = \sum_{k \geq 0} \xi_k + \sum_{k \geq 0} \eta_k.$$

Furthermore, S is bijective. S is surjective since $S(\lambda, 0, 0, 0, \dots) = \lambda$ for any $\lambda \in \mathbb{C}$, and if $S(x + X_0) = S(y + X_0)$, then $x - y \in X_0$, so $x + X_0 = y + X_0$. Now we must prove that S is an isometry. Choose $(x_k)_{k \geq 0} \in X$ such that each $x_k \geq 0$ and $\sum_{k \geq 0} x_k = 1$. For example, $x_k = \frac{1}{2^{k+1}}$. Since we have shown X/X_0 is one dimensional, $X/X_0 = \mathbb{C}((x_k)_{k \geq 0} + X_0)$. Hence, any $y + X_0 \in X/X_0$ is of the form $\lambda((x_k)_{k \geq 0} + X_0)$ for some $\lambda \in \mathbb{C}$. Then,

$$|S(y)| = |S(\lambda((x_k)_{k \geq 0} + X_0))| = |\lambda| = \|y\|.$$

So S is an isometry. \square

Theorem 3. X_0^\perp is the one dimensional subspace of ℓ^∞ , $\mathbb{C}(1, 1, 1, \dots)$, where $(1, 1, 1, \dots)$ is a constant sequence with value 1.

Proof. We identify X^* with ℓ^∞ . For $(\eta_k)_{k \geq 0} \in X_0^\perp$, we require for all $(\xi_k)_{k \geq 0} \in X_0$,

$$\sum_{k \geq 0} \xi_k \overline{\eta_k} = 0.$$

Choose the sequence $(\xi_k)_{k \geq 0} \in X_0$ as $\xi_0 = 1$ and $\xi_p = -1$ for some $p > 0$, and $\xi_k = 0$ otherwise. Then,

$$\eta_0 - \eta_p = 0.$$

Hence $(\eta_k)_{k \geq 0}$ is a constant sequence. So X_0^\perp consists of constant sequences. Clearly any constant sequence is in X_0^\perp , so $X_0^\perp = \mathbb{C}(1, 1, 1, \dots)$. \square