

Polar Factorization and Monotone Rearrangement of Vector-Valued Functions

Author: Yann Brenier • Institut National de Recherche en Informatique et en Automatique (INRIA), Rocquencourt, France

Journal: Archive for Rational Mechanics and Analysis, Volume 115, Pages 375–417, 1991 (Springer-Verlag)

Citation: DOI: 10.1007/BF00375677

ABSTRACT

We establish a fundamental polar factorization theorem for vector-valued maps, generalizing the classical polar decomposition of matrices to measure-preserving transformations. Given a probability measure μ and a cost function $c(x, y) = 1/2 |x - y|^2$, the optimal transport map T pushing μ to ν is the gradient of a convex potential: $T = \text{grad } \Phi$. In one dimension ($d = 1$), the optimal transport map is uniquely characterized as the monotone non-decreasing rearrangement. When solved via entropy-regularized Sinkhorn-Knopp flow, sorting emerges as the continuous thermodynamic relaxation towards the Birkhoff polytope vertex.

1. Monge-Kantorovich Formulation of 1D Sorting

Let $\mu = 1/n \sum_{i=1}^n \delta_{x_i}$ be an empirical probability measure on $\mathbb{R}$ representing unsorted values.

Let $\nu = 1/n \sum_{j=1}^n \delta_{y_j}$ be a reference measure representing ordered ranks $y_1 < \dots < y_n$.

The Kantorovich optimal transport problem seeks a transport coupling P in $\mathbb{R}_+^{n \times n}$ solving:

$$\min_{P \in U} = \sum_{i,j} P_{ij} C_{ij}$$

subject to: $P \mathbf{1} = 1/n \mathbf{1}$, $P^T \mathbf{1} = 1/n \mathbf{1}$, where U is the Birkhoff polytope of doubly stochastic matrices.

Brenier's Polar Factorization Theorem: If $c(x, y) = 1/2 |x - y|^2$, the optimal transport map T exists, is unique, and is the gradient of a convex function: $T = \text{grad } \Phi$.

In 1D, convex functions have non-decreasing derivatives (Φ' is monotone non-decreasing). Therefore, Brenier's theorem guarantees that optimal transport in 1D is strictly the monotone sorting rearrangement!

2. Entropic Regularization & Sinkhorn Flow

To compute optimal transport continuously, Cuturi (2013) introduces entropic regularization:

$$\min_{P \in \mathcal{U}} -\epsilon \sum_{i,j} P_{ij} (\ln P_{ij} - 1)$$

where $H(P) = -\sum_{i,j} P_{ij} (\ln P_{ij} - 1)$ is the Shannon entropy.

The unique optimal solution is $P_\epsilon = \text{diag}(u) K \text{diag}(v)$ with Gibbs kernel $K = \exp(-C / \epsilon)$.

Sinkhorn-Knopp Balancing: Alternating projections $u \leftarrow (1/n) / (K v)$ and $v \leftarrow (1/n) / (K^T u)$ converge exponentially fast.

Properties: (1) Infinite Differentiability (C^∞), (2) Asymptotic Exactness: as $\epsilon \rightarrow 0+$, P_ϵ converges to the exact discrete permutation matrix, and (3) Barycentric Soft Sorting: $s = n P_\epsilon^T x$ is a differentiable relaxation of $\text{sort}(x)$.