Table 15 (page 1 of 2). Neonatal mortality rates, by race and Hispanic origin of mother, and state: United States, average annual 1989–1991, 2003–2005, and 2006–2008

Updated data when available, Excel, and PDF: http://www.cdc.gov/nchs/hus/contents2012.htm#015.

[Data are based on linked birth and death certificates for infants]

				Not Hispanic or Latina							
State	All races			White			Black or African American				
	1989–1991 ¹	2003–2005 ²	2006–2008 ²	1989–1991	2003–2005 ²	2006–2008 ²	1989–1991 ¹	2003–2005 ²	2006–20082		
	Neonatal ³ deaths per 1,000 live births										
United States	5.7	4.6	4.4	4.6	3.7	3.6	11.1	9.2	8.7		
Alabama	7.5 4.1	5.4 3.2	6.0 3.1	5.7 3.7	4.0 *2.6	4.6 *2.3	11.1	8.5	9.0		
Arizona	5.3	4.5	4.4	4.9	4.0	4.0	11.0	7.4	8.9		
Arkansas	5.4	5.1	4.5	4.5	4.3	3.6	8.5	8.9	8.5		
California	4.6 5.0	3.5 4.6	3.5 4.3	4.1 4.7	3.0 3.8	3.0 3.6	9.2 10.9	7.2 11.9	7.1 8.5		
Connecticut	5.7	4.0	4.8	4.2	2.8	3.8	12.5	8.3	9.9		
Delaware	7.5	6.4	5.7	5.8	4.5	4.1	12.4	12.1	10.3		
District of Columbia	14.1	8.6	8.4	*5.2	*	*3.9	16.7	11.9	12.3		
Florida	6.2	4.7	4.6	4.7	3.5	3.5	10.5	8.4	8.0		
Georgia	7.9	5.6	5.2	5.5	3.9	3.7	12.0	9.2	8.3		
Hawaii	4.3	4.7	4.1 4.4	3.5	*3.1	*3.1	*	*	*16.4		
Idaho	5.3 7.0	3.9 5.1	4.4 4.8	5.2 5.1	4.0 4.1	4.0 3.9	12.7	10.0	8.5		
Indiana	6.0	5.3	4.6	5.2	4.6	4.0	11.5	10.7	10.2		
Iowa	4.8	3.4	3.4	4.5	3.2	3.2	*10.5	*6.4	*5.6		
Kansas	4.9	4.6	4.7	4.6	4.4	4.3	8.3	9.6	9.2		
Kentucky Louisiana ⁴	5.0 6.3	4.0 5.8	4.2 5.5	4.6 4.8	3.7 4.0	4.0 3.5	8.9 8.5	7.0 8.5	7.1 8.6		
Maine	4.5	4.4	4.1	4.2	4.3	4.1	*	*	*		
Maryland	5.9	5.8	5.8	3.9	3.7	3.9	10.2	10.1	9.5		
Massachusetts	4.9	3.7	3.7	4.1	3.0	3.0 4.1	10.4	7.5	8.0		
Michigan	6.9 4.3	5.6 3.2	5.3 3.6	4.9 3.9	4.3 2.9	3.2	14.0 10.7	11.6 5.5	10.1 7.5		
Mississippi	7.1	6.2	6.1	4.9	3.5	4.0	9.5	9.7	8.6		
Missouri	6.0	5.1	4.8	5.0	4.4	3.8	10.6	9.4	10.2		
Montana	4.6	3.5	3.4	4.2	3.4	3.1	*0.0	*0.4	*7.4		
Nebraska Nevada	4.5 4.3	3.7 3.7	3.8 3.9	4.2 3.8	3.3 3.6	3.5 3.3	*9.8 *8.3	*9.4 7.7	*7.4 7.4		
New Hampshire 4	4.3	3.9	3.4	4.4	3.6	3.2	*	*	*		
New Jersey	5.8	3.8	3.6	4.5	2.6	2.7	11.4	8.2	7.4		
New Mexico	5.0 6.5	3.6 4.2	3.5 3.7	4.8 4.3	4.1 3.4	3.9 2.9	12.6	8.0	7.5		
New York	7.3	4.2 6.0	5.7 5.5	4.3 5.3	4.2	4.0	11.9	11.4	7.5 9.9		
North Dakota	5.0	4.8	4.4	4.7	4.5	4.3	*	*	*		
Ohio	5.5	5.3	5.2	4.8	4.2	4.1	9.8	10.7	10.4		
Oklahoma ⁴	4.4 4.4	4.6 3.8	4.6 3.6	4.1 4.0	4.3 3.7	4.3 3.5	6.3 *11.6	8.8	8.9		
Oregon	6.2	5.6 5.2	5.0 5.2	4.9	4.0	3.9	12.5	9.5	9.7		
Rhode Island	6.4	4.8	5.0	5.3	3.5	3.1	*9.8	*7.6	*8.9		
South Carolina	7.7	6.1	5.4	5.4	4.2	3.7	11.3	9.8	8.7		
South Dakota	5.1	4.3	4.2	4.5	4.1	3.6	*	*	*		
Tennessee	6.5	5.7	5.3	4.9	4.0	3.9	11.8	11.7	10.5		
Texas	4.7 3.7	4.2 3.4	3.9 3.4	4.1 3.6	3.7 3.1	3.2 3.2	8.5	8.0	7.3		
Vermont	4.1	3.8	3.2	3.9	3.7	2.9	*	*	*		
Virginia	6.8	5.2	4.9	4.8	3.9	3.6	13.0	9.6	9.4		
Washington	4.3	3.4	3.1	3.8	2.9	2.5	9.7	5.6	4.9		
West Virginia	5.8 5.1	4.8 4.3	4.3 4.3	5.6 4.6	4.7 3.5	4.0 3.6	*9.7 9.1	10.3	*11.0 9.9		
Wyoming	3.9	4.6	4.2	3.8	4.6	3.9	9.1	10.0	9.9		

See footnotes at end of table.

Health, United States, 2012 Trend Tables

Table 15 (page 2 of 2). Neonatal mortality rates, by race and Hispanic origin of mother, and state: United States, average annual 1989–1991, 2003–2005, and 2006–2008

Updated data when available, Excel, and PDF: http://www.cdc.gov/nchs/hus/contents2012.htm#015.

[Data are based on linked birth and death certificates for infants]

State	Hispanic or Latina ⁵			American Indian or Alaska Native ⁶			Asian or Pacific Islander ⁶		
	1989–1991	2003–2005 ²	2006–2008 ²	1989–1991 ¹	2003–2005 ²	2006–2008 ²	1989–1991	2003–2005 ²	2006–2008
				Neonatal ³ d	leaths per 1,0	00 live births			
United States	4.8	3.9	3.7	5.9	4.3	4.3	3.9	3.3	3.2
Alabama	*	*4.4	4.7	*	*	*	*	*	*
Alaska	*	*	*	*5.7	*4.2	*4.2	*	*	*
Arizona	5.0	4.8	4.4	5.4	4.2	4.3	*	*4.3	*4.8
Arkansas	*	*3.8	*3.7	*	*	*	*	*	*
California	4.4 4.4	3.5 5.3	3.3 5.1	6.3	*3.1	*4.2	3.6	2.9 *4.0	3.0 *4.1
Colorado	4.4 5.3	5.3 6.1	5.1 4.6	*	*	*	*	4.0	4.1 *4.4
Delaware	5.5 *	*4.4	*4.3	*	*	*	*	*	4.4 *
District of Columbia	*	*	*	*	*	*	*	*	*
Florida	5.1	3.5	3.7	*	*	*	*4.4	3.9	4.0
Georgia	*5.7	3.8	3.3	*	*	*	*5.3	4.1	3.2
Hawaii	*6.6	*5.7	*3.3	*	*	*	4.2	4.8	4.1
Idaho	*	*3.8	5.5	*	*	*	*	*	*
Illinois	6.4	4.3	4.3	*	*	*	3.9	3.3	4.0
Indiana	*4.7	4.9	3.8	*	*	*	*	*	*
lowa	*	*3.7	*4.3	*	*	*	*	*	*
Kansas	*5.4	3.4	4.9	*	*	*	*	*	*
Kentucky	*	*5.5	*3.9	*	*	*	*	*	*
Louisiana 7		*	*2.3	*	*	*	*	*	*6.3
Maine									
Maryland	*4.7	3.8	4.0	*	*	*	*4.5	*3.4	4.1
Massachusetts	5.8	4.9	4.5	*	*	*	*3.9	*2.8	*2.2
Michigan	5.2	5.1	4.9	* 4 0	*	*	*	3.7	*3.6
Minnesota	*	*2.9	2.9	*4.9	*	*4.6	*3.2	*2.4	3.9
Mississippi	*	4.6	*4.8 *3.6	*	*	*	*	*4.0	*
Montana	*	4.0	3.0 *	*7.6	*	*4.8	*	4.0	*
Nebraska	*	*3.6	*3.5	7.0	*	*	*	*	*
Nevada	*4.1	2.6	3.8	*	*	*	*	*4.0	*3.2
New Hampshire 7		*	*	*	*	*	*	*	*
New Jersey	5.1	3.7	3.6	*	*	*	*3.4	3.4	2.0
New Mexico	4.9	3.3	3.4	4.9	*3.5	*2.8	*	*	*
New York	6.4	3.8	3.3	*	*	*	4.1	2.7	2.3
North Carolina	*5.5	4.7	4.5	*7.7	*8.2	10.2	*	*4.4	*3.4
North Dakota	*	*	*	*	*7.5	*	*	*	*
Ohio	*5.4	4.8	4.4	*	*	*	*	*3.3	*3.4
Oklahoma ⁷		4.0	3.2	*3.7	3.6	4.4	*	*	*
Oregon	6.5	3.8	3.8	*	*	*	*5.3 *5.0	*3.9	*2.4
Pennsylvania	7.3 *4.9	5.5 *5.8	5.7 *5.9	*	*	*	*5.2	3.7	4.5
	4.5								
South Carolina	*	4.7	4.3	*	*	*	*	*	*
South Dakota	*	5.0	27	*8.2	*5.6	*6.0	*	*4.7	*4.3
Tennessee	4.2	5.0 3.8	3.7 3.7	*	*	*	4.0	2.8	2.8
Utah	*3.6	4.2	3.6	*	*	*	→.U *	*5.3	*4.4
Vermont	*	*	3.0	*	*	*	*	*	*
Virginia	*4.8	4.0	4.0	*	*	*	*4.1	3.6	3.0
Washington	4.9	3.7	3.5	*8.5	*4.6	*4.5	*2.7	2.9	2.7
West Virginia	*	*	*	*	*	*	*	*	*
Wisconsin	*3.9	4.4	4.1	*	*	*	*	*5.3	*4.1
Wyoming	*	*	*	*	*	*	*	*	*

^{*} Estimates are considered unreliable. Rates preceded by an asterisk are based on fewer than 50 deaths in the numerator. Rates not shown are based on fewer than 20 deaths in the numerator.

NOTES: Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states. See Appendix II, Race. National linked files do not exist for 1992–1994.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use and nonpublic-use Linked Birth/Infant Death Data Set. See Appendix I, National Vital Statistics System (NVSS).

2 Trend Tables Health, United States, 2012

^{- - -} Data not available.

¹Rates based on unweighted birth cohort data.

²Rates based on period file using weighted data. See Appendix I, National Vital Statistics System (NVSS), Linked Birth/Infant Death Data Set.

³Infants under 28 days of age.

⁴Rates for white and black are substituted for non-Hispanic white and non-Hispanic black for Louisiana for 1989, Oklahoma for 1989–1990, and New Hampshire for 1989–1991.

⁵Persons of Hispanic origin may be of any race. See Appendix II, Hispanic origin.

⁶Includes persons of Hispanic origin.

Rates for Hispanic origin exclude data from states not reporting Hispanic origin on the birth certificate for 1 or more years in a 3-year period.