Hypophthalmichthys nobilis (Bighead Carp)
Native
Exotic

Collection Info
Point Map
Specimen Profile
Animated Map
Impacts

U.S. Fish and Wildlife Service

Hypophthalmichthys nobilis (Richardson, 1845)

Common name: Bighead Carp

Synonyms and Other Names: *Aristichthys nobilis* (Richardson, 1845), *Leuciscus nobilis* (Richardson, 1845)

Taxonomy: available through ITIS

Nativity: This species is listed by the U.S. Fish and Wildlife Service as injurious wildlife.

Identification: The bighead carp is a large, narrow fish with eyes that project downward. Coloration of the body is dark gray, fading to white toward the underside, and with dark blotches on the sides. Its head has no scales, a large mouth with no teeth, and a protruding lower jaw. Its gill rakers are formed into sieve-like plates. Its dorsal fin has 3 to 4 very similar to the silver carp, and can be distinguished by the dark coloration on its sides. The bighead carp can be identified by a smooth keel between the anal and pelvic fins that does not extend anterior of the base of the pelvic fins.

Distinguishing characteristics were given in Berg (1949) and Jennings (1988). Distinguishing characteristics, along with keys that include this species and photographs or illustrations also were included in a few of the more recently published state field guides, such as Pflieger and Robertson 1988; Etnier and Starnes 1993; Cooper 1983; and Jenkins and Burkhead 1994. Maximum size: 60 kg and 1.8 m (Shireman 1984).

Size: 40 kg and 1.8 m

Native Range: Southern and central China (Li and Fang 1990; Robins et al. 1991).

Hydrologic Unit Codes (HUCs) Explained
Interactive maps: Point Distribution Maps

Nonindigenous Occurrence:

Table 1. States with nonindigenous occurrences, the earliest and latest observations in each state, and the number of HUCs with observations*. Names and dates are hyperlinked to their relevant specimen records. The list of references for all nonindigenous occurrences of *Hypophthalmichthys nobilis* are listed below.

State	First Observed	Last Observed	Total HUCs with observations*	HUCs with observations*
AL	1984	2021	11	Bayou Black Warrior Tombigbee Guntersville Lake Lower Alabama Lower Tallapoosa Middle Alabama Middle Tombigbee Chickasaw Middle Tombigbee Lubbub-Bucatunna Upper Black Warrior
AZ	2007	2007	1	Upper Santa Cruz
AR	1988	2018	21	Bayou Meto Beaver Lake Bayou Meto Boeuf Cadron Current Dardanelle Reservoir Frog Multberry Lake Conway Point Remove Lower Arkansas Lower Arkansas Maumelle Lower Black Lower Little Arkansas Oklahoma Lower Mississippi Greenville Lower Mississippi Memphis Lower Ouachita Bayou De Loutre Lower St Francis Lower White Bayou Bartholomew Lake Village Middle Black Upper Little Missouri Upper White Village
CA	1992	2013	3	Rio Chico Creek Sacramento River Russian Thomes Creek Sacramento River
CO	1996	2009	3	Cache La Poudre Lower Republican Middle Arkansas Cherry Creek
FL	1989	2019	7	Lake Okeechobee Lower Ocklawaha/Silver St. Andrew St. Joseph Bays
IL	1986	2015	36	Kaskaskia Big Muddy Cache Cahokia-Joachim Chicagoland Cypress Ditch Hay Pittance Embarrass Illinois Headwaters Kankakee Little Wabash Lower Kaskaskia Lower Little Wabash Lower Mississippi Hydrologic Accounting Unit Lower Wabash Lower Fox Lower Illinois Lower Kankakee Lower Rock Lower Sangamon Lower Wabash Mazonnie Middle Kaskaskia Middle Wabash Busseron Peruque-Piasa Pike-Keye Skillet Spoon Du Sac Upper Illinois Upper Kaskaskia Upper Mississippi Cape Girardeau Upper Sangamon
IN	1984	2015	14	Blue Sinkhole Hall Hoffman Prairie Little Calumet-Galien Lower East Fork White Lower Ohio Little Pigeon Lower Wabash Lower White Middle Ohio-Laughery Middle Wabash Busseron Middle Wabash-Little Vermillion Missinewa Muscatatuck Ohio Ravine Silver Little Kentucky Tippecanoe Upper East Fork White Upper Wabash
LA	1986	2014	19	Amite River Bayou De Pandilla Bayou Sara Bodcau-Bayou Middle Bayou Sara Cross Lake Cooper Creek Flint-Henderson Keys Wooten Water Lake Red Rock Little Sunnie Lower Bayou Des Cannes Lower Bayou Lacombe Middle Little Middle Levee Nonlevee Plain Sabine Upper Bayou Des Cannes Upper Bayou Teche Upper Ouachita Bayou De Loutre Upper Sabine Upper White Village
KS	1982	2014	10	Big Nemaha Independence-Sugar Kansas Lower Kansas Lower Republican Middle Neosho Turkey Wolf Upper Salt Fork Arkansas Upper Walnut River
KY	1986	2021	19	Barren Elk Clarks River Bel Rio Sinkhole Highland-Pigeon Kentucky Lake Lalcksie Little Scioto-Tygarts Lower Cumberland Lower Green Lower Kentucky Lower Missouri-Hungerly Lower Ohio-Little Pigeon Lower Ohio-Little Powell Lower Ohio-Little Sandy Middle Ohio-Laurberry Ohio Brush Whitewoi Rolling Fork Rough Silver Little Kentucky
LA	1983	2014	13	Atchafalaya Bayou Teche Boeuf East Central Louisiana Coastal Lake Maurepas Lake Pontchartrain Lower Mississippi Baton Rouge Lower Mississippi New Orleans Lower Pearl Lower Vermilion Bayou Des Cannes Lower Bayou Lacombe Middle Red Lowlands
MN	1996	2020	9	Buffalo Whitewater Cross Yellow Black Yellow Medicine La Crosse-Pine Little Sioux Lower St. Croix Middle Minnesota Rush Vermillion Twin Cities
MS	1986	2012	14	Big Sunflower Bogue Chitto Bogue Falaya Lalbabuchia Lower Mississippi Hydrologic Accounting Unit Lower Mississippi-Natchez Lower Mississippi-Vicksburg Lower Yazoo Pascagoula Pascagoula Tallahatchie Upper Yazoo Yalobusha
MO	1967	2022	29	Bear Wyaconda Big Muddy Black Lamine Cahokia-Joachim Independence Sugar Lake of the Ozarks Lamine Lower Chariton Lower Grand Lower Mississippi-Memphis Lower Missouri Lower Missouri-Crooked Lower Missouri-Moreau Lower Osage One Hundred and Two Perrapue Piasa Tarkio-Wolf The Sny Town of Marshall Saint Johns Bayou Upper Mississippi-Kaskaskia Wakenda-Big Lower Grand
NE	1990	2012	15	Big Papillion-Mosquito Blackbird-Soldier Keg Weeping Water Lewis and Clark Lake Little Nemaha Lower Elkhorn Lower North Loup Lower North Platte Lower Platte Middle Platte-Buffalo Middle Platte-Prairie Salt Turkey Wolf Upper Elkhorn
NJ	2010	2010	1	Middle Delaware-Musconetcong
NC	2011	2011	1	Chowan
ND	2019	2019	1	Upper James
OH	1995	2022	10	Lake Erie Little Muskingum Middle Island Lower Scioto Mahoning Middle Ohio-Laurberry Ohio Brush Whitewoi Scioto Upper Ohio Upper Ohio-Wheeling
OK	1992	2021	5	Bois D'arc Island Elk Kiamichi Lake O' The Cherokees Lower Cimarron-Skeleton Lower Neosho Twin Marshfields
PA	2014	2014	1	Upper Ohio
SD	1998	2022	5	Levya and Clark Lake Lower James Middle James Vermillion
TN	1951	2017	9	Guntersville Lake Hatchie-Loosahatchie Lower Cumberland Lower Tennessee Lower Hatchie Lower Mississippi-Memphis Middle Tennessee-Chickamauga Obion Rutts Bar Lake
TX	1989	2015	11	Bois D'arc Island Buffalo-San Jacinto Cadde Lake Lake O'the Pines Lower Sabine Rita Blanca Upper Cross Cross Timbers Upper Neches Upper Sabine Wert Fork San Jacinto
VA	1996	2007	1	South Fork Shenandoah
WV	1997	2024	4	Lower Kanawha Raccoon-Symmes Upper Ohio-Shade Upper Ohio-Wheeling
WI	1996	2022	5	Buffalo-Whitewater Cross Yellow Great Lakes Marquette La Crosse-Pine Lower Chippewa Lower St. Croix Lower Wisconsin Rush Vermillion St. Louis

Table last updated 5/28/2025

* Populations may not be currently present.

Ecology: Bighead carp is a powerful filter-feeder with a wide food spectrum that grows fast and reproduces quickly (Xie and Chen 2001), which makes this species a strong competitor. The diet of this species overlaps with that of the native paddlefish, gizzard shad, and threadfin shad (Kolar et al. 2005). Bighead carp are thought to displace paddlefish and buffalo fishes for native larval fishes and mussels (Cech and Papp 1986; Kolar et al. 2005; Lohmeyer and Garvey 2009; Sampson et al. 2009).

Fireside bighead carp reach sexual maturity at three years of age, while males can reach sexual maturity in two years; however, this varies significantly with changing environmental conditions (Hart 1970; Kolar et al. 2005). Bighead carp spawn in open water in the presence of flowing water and high turbidity (Kolar et al. 2005). Fecundity increases with age and body weight and is directly related to growth rate (Hartig et al. 1990). In its native range, Bighead Carp has a fecundity ranging from 280,000-1.1 million eggs. In North America, fecundity ranged from 4,792-1.6 million eggs (Kaja et al. 2011). Bighead carp produce semi-buoyant eggs that are 5-6 mm in diameter and hatch in 24-36 hours (Kolar et al. 2005). The eggs float for 40-60 hours before hatching.

Means of Introduction: Bighead carp were first imported into the United States in 1973 by a private fish farmer in Arkansas who wanted to use them in combination with other phytoplankton feders to improve water quality (Freeze and Henderson 1982; Kolar et al. 2005). In 1974 the Arkansas and Fish Farming Experimental Station received a shipment of bighead carp. Bighead carp escaped from ponds that they were held in during flooding events. By 1980 the species began to appear in open waters, the Ohio and Mississippi rivers, in the early 1980s, likely as a result of escapes from aquaculture facilities (Jennings 1988). In April 1994, several thousand bighead carp, along with a few black carp, escaped from a private fish farm in Missouri (Kolar et al. 2005). Bighead carp were also released into the Mississippi River in 1994 during the Great Flood of 1993 (Kolar et al. 2005). Bighead carp were also released into Lake Erie in 1995