

boson no spin

$$T_0^2 = 1$$

$$T_0 \psi(t, \vec{r}) = \psi^*(-t, \vec{r})$$

fermion spin  $\frac{1}{2}$

$$T^2 = -1$$

$$T \begin{pmatrix} \psi_1(t, \vec{r}) \\ \psi_2(t, \vec{r}) \end{pmatrix} = e^{i\varphi} \sigma_y \begin{pmatrix} \psi_1^*(-t) \\ \psi_2^*(-t) \end{pmatrix}$$

zero spin

$$\hat{H} \psi_{nl} = E_n \psi_{nl} \quad l=1 \dots d(n)$$

general sol of Schrödinger eqn  $\psi(\vec{r}, t) = \sum_n a_n \psi_{nl}(\vec{r}) e^{-\frac{i}{\hbar} E_n t}$   
( $\hat{H}$  is real, not explicitly dep. on t)

$$\hat{T}_0 \psi(\vec{r}, t) = \sum_n a_n^* \psi_{nl}^*(\vec{r}) e^{-\frac{i}{\hbar} E_n t}$$

effectively  $\hat{T}_0 : \psi_{nl}(\vec{r}) \rightarrow \psi_{nl}^*(\vec{r})$

if  $\hat{H}$  is real,  $\hat{H} \psi_{nl} = E_n \psi_{nl}$   $\psi_{nl}$  is eigenstate

$\Rightarrow \hat{H} \psi_{nl}^* = E_n \psi_{nl}^*$   $\psi_{nl}^*$  is eigenstate

$\psi_{nl}$  basis in  $\mathcal{H}^{(n)}$ ,  $\psi_{nl}^*$  is basis in  $\mathcal{H}^{(n)*}$

whether  $\mathcal{H}^{(n)}$  and  $\mathcal{H}^{(n)*}$  the same?

(1)  $\mathcal{H}^{(n)} \cong \mathcal{H}^{(n)*}$  equivalent rep of symm. group

(2)  $\mathcal{H}^{(n)} \not\cong \mathcal{H}^{(n)*}$  not equivalent ....

$$\text{eigenspace} = \mathcal{H}^{(n)} \oplus \mathcal{H}^{(n)*}$$

real rep, pseudo real rep, and complex rep

Given a symm. group  $G$ , unitary irrep  $D: g \mapsto D(g) \in \mathcal{H}^{(n)}$

Def  $D(g)$  is unitary on  $\mathcal{H}^{(n)}$ ,  $\psi_{hi}$  orthonormal basis in  $\mathcal{H}^{(n)}$

(complex  
conjugate  
irrep)

$$D(g) \psi_{hi}(\vec{r}) = \sum_j \psi_{hj}(\vec{r}) D_{ji}(g)$$

$D_{ji}(g)$  unitary matrix

complex conjugate :  $D(g)^* \psi_{hi}^*(\vec{r}) = \sum_j \psi_{hj}^*(\vec{r}) D_{ji}^*(g)^*$

$$\psi_{hi}^*(\vec{r}) = T_0 \psi_{hi}(\vec{r})$$

$D_{ij}(g)$  unitary irrep matrix  
(irrep  $D$ )

$\xrightarrow{T_0} D_{ij}^*(g)$  unitary irrep matrix  
irrep  $D^*$ :

Complex conjugate irrep

relation between  $D$  and  $D^*$

(1) if  $\exists$  unitary transf.  $U: \mathcal{H}^{(n)} \rightarrow \mathcal{H}^{(n)}$  s.t

$$U D(g) U^{-1} = D_0(g) \quad \forall g \in G$$

↑  
real matrix

$$\Rightarrow \underline{D \text{ is equivalent to } D^*}, \text{ since } U^* D(g)^* U^{*-1} = D_0(g)$$

$$\Rightarrow \underline{D^*(g) = U^{*-1} D_0(g) U^*} = \underline{U^{*-1} U D(g) U^{-1} U^*}$$

$U^{*-1} U$  is unitary

we say  $D \simeq D^*$  is a real rep.

(2)  $D$  is <sup>unitarily</sup> equivalent to  $D^*$ , but they are not equivalent to real rep.

$$\nexists U \text{ s.t. } D(g) = U D_0(g) U^{-1} \quad \forall g \in G$$

$\uparrow$   
real matrix

We say,  $D \simeq D^*$  is pseudo-real

(3)  $D$  is inequivalent to  $D^*$ .

We say,  $D$  and  $D^*$  are complex reps

distinguish (3) from (1) and (2), check character

$$\text{if } D \simeq D^*, \text{ then } \chi(g) = \chi^*(g) \quad \forall g \in G$$

if  $\chi(g) \neq \chi^*(g)$ , then  $D$  &  $D^*$  are complex reps.

distinguish between (1) & (2)

$$D^* \simeq D \rightarrow D^*(g) = Z D(g) Z^{-1} \quad \forall g \in G$$

$\uparrow$   
unitary matrix

Complex conjugate  $D(g) = Z^* D^*(g) Z^{*-1}$

$$\Rightarrow D^*(g) = Z Z^* D^*(g) (Z Z^*)^{-1}$$

$$\therefore [Z Z^*, D^*(g)] = 0 \quad \forall g \in G$$

by Schur's lemma, when  $D^*$  is irrep,  $Z Z^* = c \mathbb{1}$   
 $c \in \mathbb{C}$

$$Z \text{ is unitary, } Z^* = (Z^T)^{-1}$$

$$\Rightarrow Z (Z^T)^{-1} = c \mathbb{1} \Rightarrow \underline{Z = c Z^T}$$

$$\xRightarrow{\text{transpose}} Z^T = c Z$$

$$\Rightarrow Z = c^2 Z \Rightarrow c^2 = 1, c = \pm 1$$

$$D^* \simeq D \Rightarrow Z Z^* = \pm \mathbb{1}$$

Thm:  $Z Z^* = \mathbb{1}$  iff  $D$  is real

$Z Z^* = -\mathbb{1}$  iff  $D$  is pseudo-real

pf if  $D$  is real  $\rightarrow D^*(g) = (U^*)^{-1} D_0(g) U^*$   
 $= (U^*)^{-1} U D(g) U^{-1} U^*$

on the other hand  $D^*(g) = Z D(g) Z^{-1}$

$$\Rightarrow (U^*)^{-1} U = b Z \quad b \in \mathbb{C}$$

Complex conjugate  $U^{-1} U^* = b^* Z^*$

$$U^{-1} U^* (U^*)^{-1} U = b^* b Z^* Z = |b|^2 Z^* Z$$

$$\parallel$$
$$1$$

$$\underline{Z^* Z = |b|^{-2} \mathbb{1}}$$

$$\underline{Z Z^* = c \mathbb{1}}$$

$$\Rightarrow |b|^2 c = 1$$

$$c = 1 \text{ or } -1$$

$$c = 1 \quad |b|^2 > 0$$

conversely if  $c = 1$  ( $Z Z^* = \mathbb{1}$ )

$Z$  unitary,  $Z = e^{iA}$   $A$  hermitian

$$Z^* = Z^{-1} \Leftrightarrow Z^T = Z, \quad Z \text{ symmetric}$$

$$Z^* = (Z^T)^{-1}$$

$$\text{def. } U = Z^{\frac{1}{2}} = e^{iA/2} \quad \text{unitary}$$

$$U^2 = Z$$

$$\underline{U^* Z = e^{-iA/2} e^{iA} = e^{\frac{1}{2}iA} = U}$$

$$\begin{aligned} \forall g \in G, [U D(g) U^{-1}]^* &= U^* D(g)^* U^{*-1} \\ &= U^* Z D(g) Z^{-1} U^{*-1} \\ &= U D(g) U^{-1} = D_o(g) \text{ real} \end{aligned}$$

$\Rightarrow D$  is real rep.

$$Z Z^* = \mathbb{1} \Leftrightarrow D \text{ is real rep}$$

$$\Rightarrow Z Z^* = -\mathbb{1} \Leftrightarrow D \text{ is pseudo-real rep.} \quad \square$$

Distinguish real & pseudo-real reps by characters

$$\underline{D^*(g) = Z D(g) Z^{-1}}$$

orthogonality  $\sum_g D_{\alpha\delta}^*(g) D_{\beta\gamma}(g) = \delta_{\alpha\beta} \delta_{\gamma\delta} \frac{h}{\dim(D)=d}$

$$\sum_{\alpha, \delta} \underline{Z_{\tau\alpha}^{-1}} \sum_g \sum_{\sigma, \rho} \underline{Z_{\alpha\sigma}} D_{\sigma\rho}(g) \underline{Z_{\rho\delta}^{-1}} D_{\beta\gamma}(g) \times \underline{Z_{\delta\chi}}$$

$$\Rightarrow \sum_g D_{\tau\chi}(g) D_{\beta\gamma}(g) = Z_{\gamma\chi} Z_{\tau\beta}^{-1} \frac{h}{d}$$

let  $\chi = \rho$ .  $\sum_{\beta}$ ,  $\sum_g \sum_{\beta} D_{\tau\beta}(g) D_{\beta\gamma}(g)$

$$\sum_g D_{\tau\gamma}(g^2) = (Z Z^*)_{\tau\gamma} \frac{h}{d}$$

$$= \pm \delta_{\tau\gamma} \frac{h}{d}$$

let  $\tau = \gamma$ ,  $\sum_{\tau}$ ,

$$\boxed{\sum_g \chi(g^2) = \pm h}$$

for complex rep.  $D \neq D^* \Rightarrow \sum_g [D_{\alpha\beta}^*(g)]^* D_{\gamma\delta}(g) = 0$

$$\sum_j \text{Dop}(j) \text{Dss}(j)$$

$$\Rightarrow \sum_j \chi(j^2) = 0$$

Summary :

$$\frac{1}{h} \sum_{j \in G} \chi(j^2) = \begin{cases} 1 & \text{real rep} \\ -1 & \text{pseudo-real rep} \\ 0 & \text{complex.} \end{cases}$$

Extra-degeneracy of  $\hat{H}$  due to time-reversal inv.

$$\hat{H} \psi_{nl} = E_n \psi_{nl} \quad l = 1 \dots d$$

$$\psi_{nl} \in \mathcal{H}^{(n)} \quad \text{unitary irrep. of symmetry group } G$$

$$l = 1 \dots d \quad \dim \mathcal{H}^{(n)} = d$$

find degeneracy at  $E_n$ , deg. at  $E_n$  is either  $d$  or  $2d$  (spin-zero)

if  $\hat{H}$  is real and not explicitly dep on  $t$

$$\hat{H} \psi_{nl} = E_n \psi_{nl}$$

$$\hat{H} \psi_{nl}^* = E_n \psi_{nl}^*$$

$$\psi_{nl}^*(\vec{r}) = \hat{T}_0 \psi_{nl}(\vec{r})$$

• if  $\forall \psi_{nl}^*$ ,  $\psi_{nl}^* \in \mathcal{H}^{(n)}$  spanned by  $\psi_{nl}$

then  $\mathcal{H}^{(n)}$  is the final eigenspace i.e.  $\mathcal{H}^{(n)} = \mathcal{H}^{(n)*}$

degeneracy at  $E_n$  is  $d$

- otherwise, eigenspace  $= \mathcal{H}^{(n)} \oplus \mathcal{H}^{(n)*}$

degeneracy at  $E_n = 2d$  (extra degeneracy)