


- (1) In following Fig., the four particles form a square of edge length $a = 5.00$ cm and have charges $q_1 = +10.0$ nC, $q_2 = -20.0$ nC, $q_3 = +20.0$ nC, and $q_4 = -10.0$ nC. In unit vector notation, what net electric field do the particles produce at the square's center.



Soln:

EXPRESS Applying the superposition principle, the net electric field at the center of the square is

$$\vec{E} = \sum_{i=1}^4 \vec{E}_i = \sum_{i=1}^4 \frac{1}{4\pi\epsilon_0} \frac{q_i}{r_i^2} \hat{r}_i.$$

With $q_1 = +10$ nC, $q_2 = -20$ nC, $q_3 = +20$ nC, and $q_4 = -10$ nC, the x component of the electric field at the center of the square is given by, taking the signs of the charges into consideration,

$$\begin{aligned} E_x &= \frac{1}{4\pi\epsilon_0} \left[\frac{|q_1|}{(a/\sqrt{2})^2} + \frac{|q_2|}{(a/\sqrt{2})^2} - \frac{|q_3|}{(a/\sqrt{2})^2} - \frac{|q_4|}{(a/\sqrt{2})^2} \right] \cos 45^\circ \\ &= \frac{1}{4\pi\epsilon_0} \frac{1}{a^2/2} (|q_1| + |q_2| - |q_3| - |q_4|) \frac{1}{\sqrt{2}}. \end{aligned}$$

Similarly, the y component of the electric field is

$$\begin{aligned} E_y &= \frac{1}{4\pi\epsilon_0} \left[-\frac{|q_1|}{(a/\sqrt{2})^2} + \frac{|q_2|}{(a/\sqrt{2})^2} + \frac{|q_3|}{(a/\sqrt{2})^2} - \frac{|q_4|}{(a/\sqrt{2})^2} \right] \cos 45^\circ \\ &= \frac{1}{4\pi\epsilon_0} \frac{1}{a^2/2} (-|q_1| + |q_2| + |q_3| - |q_4|) \frac{1}{\sqrt{2}}. \end{aligned}$$

The magnitude of the net electric field is $E = \sqrt{E_x^2 + E_y^2}$.

$$E_x = \frac{1}{4\pi\epsilon_0} \frac{\sqrt{2}}{a^2} (|q_1| + |q_2| - |q_3| - |q_4|) = \frac{1}{4\pi\epsilon_0} \frac{\sqrt{2}}{a^2} (10 \text{ nC} + 20 \text{ nC} - 20 \text{ nC} - 10 \text{ nC}) = 0$$


and

$$E_y = \frac{1}{4\pi\epsilon_0} \frac{\sqrt{2}}{a^2} (-|q_1| + |q_2| + |q_3| - |q_4|) = \frac{1}{4\pi\epsilon_0} \frac{\sqrt{2}}{a^2} (-10 \text{ nC} + 20 \text{ nC} + 20 \text{ nC} - 10 \text{ nC})$$


$$= \frac{(8.99 \times 10^9 \text{ N} \cdot \text{m}^2 / \text{C}^2)(2.0 \times 10^{-8} \text{ C})\sqrt{2}}{(0.050 \text{ m})^2}$$

$$= 1.02 \times 10^5 \text{ N/C}.$$

Thus, the electric field at the center of the square is $\vec{E} = E_y \hat{j} = (1.02 \times 10^5 \text{ N/C}) \hat{j}$.



- (2) In Fig. below three particles are fixed in place and have charges $q_1 = q_2 = +e$ and $q_3 = +2e$. Distance $a = 6.00 \text{ mm}$. What are the (a) magnitude and (b) direction of the net electric field at point P due to the particles?



Soln:

By symmetry we see that the contributions from the two charges $q_1 = q_2 = +e$ cancel each other, and we simply compute magnitude of the field due to $q_3 = +2e$.


(a) The magnitude of the net electric field is

$$|\vec{E}_{\text{net}}| = \frac{1}{4\pi\epsilon_0} \frac{2e}{r^2} = \frac{1}{4\pi\epsilon_0} \frac{2e}{(a/\sqrt{2})^2} = \frac{1}{4\pi\epsilon_0} \frac{4e}{a^2}$$

$$= (8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2) \frac{4(1.60 \times 10^{-19} \text{ C})}{(6.00 \times 10^{-6} \text{ m})^2} = 160 \text{ N/C}.$$

(b) This field points at 45.0° , counterclockwise from the x axis.

- (3) Figure, below shows an electric dipole. What are the (a) magnitude and (b) direction (relative to the positive direction of the x axis) of the dipole's electric field at point P, located at distance $r \gg d$?



Soln:

(a) Consider the figure below. The magnitude of the net electric field at point P is

$$|\vec{E}_{\text{net}}| = 2E_1 \sin \theta = 2 \left[\frac{1}{4\pi\epsilon_0} \frac{q}{(d/2)^2 + r^2} \right] \frac{d/2}{\sqrt{(d/2)^2 + r^2}} = \frac{1}{4\pi\epsilon_0} \frac{qd}{[(d/2)^2 + r^2]^{3/2}}$$

For $r \gg d$, we write $[(d/2)^2 + r^2]^{3/2} \approx r^3$ so the expression above reduces to

$$|\vec{E}_{\text{net}}| \approx \frac{1}{4\pi\epsilon_0} \frac{qd}{r^3}.$$

(b) From the figure, it is clear that the net electric field at point P points in the $-\hat{j}$ direction, or -90° from the $+x$ axis.

