



GRAPE-Dilepton (Version 1.1) A generator for dilepton production in ep collisions

Tetsuo Abe

Department of Physics, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8654, Japan

Received 24 April 2000; received in revised form 4 October 2000

Abstract

GRAPE-Dilepton is a Monte Carlo event generator for dilepton production in ep collisions. The cross-section calculation is based on the exact matrix elements in the electroweak theory at tree level. The dilepton productions via $\gamma\gamma$, γZ^0 , $Z^0 Z^0$ collisions and via photon internal conversion are taken into account. In addition, the effects of the Z^0 on/off-shell production are also included. The relevant Feynman amplitudes are generated by the automatic calculation system GRACE. The calculation of the proton vertex covers the whole kinematical region. This generator has an interface to PYTHIA and SOPHIA to obtain complete hadronic final states. This program can be downloaded from the CPC Program Library under catalogue identifies <http://www-zeus.desy.de/~abe/grape/>. © 2001 Elsevier Science B.V. All rights reserved.

PROGRAM SUMMARY

Number of bytes in distributed program, including test data, etc.:
999 438

Title of program: GRAPE-Dilepton (v1.1)

Distribution format: tar gzip file

Catalogue identifier: ADNR

Keywords: Dilepton, lepton-pair, ep collision, Bethe–Heitler, Z boson, dipole form factor, hadron tensor, lepton tensor, structure function, parton density, GRACE

Program Summary URL: <http://cpc.cs.qub.ac.uk/summaries/ADNR>

Program obtainable from: CPC Program Library, Queen's University of Belfast, N. Ireland and from <http://www-zeus.desy.de/~abe/grape/>

Nature of physical problem

A precise estimation of the cross-section of the electroweak dilepton production in ep collisions is required in various physics analyses, where 8~48 Feynman diagrams can contribute.

Operating system under which the program has been tested: UNIX

Method of solution

The automatic calculation system GRACE is used to obtain all of the relevant helicity amplitudes. The phase space is divided into the

Programming language used: Fortran77

Memory required to execute with typical data: 7 Mwords for integrations, 9 Mwords for event generations

E-mail address: abe@helios.kek.jp (T. Abe).

3 regions according to the kinematics at the proton vertex, and the 3 different calculation methods are applied. The radiative corrections are included using the structure function and the parton shower methods.

Restrictions on the complexity of the problem

Higgs, the proton- Z^0 coupling and lepton-pair production through

photon radiation from the proton are not included. The contribution from the resolved photon, i.e. Drell–Yan process in ep collisions is not included.

Typical running time

1 hour for a cross-section integration and 1 msec per 1 event for an event generation.

LONG WRITE-UP

1. Introduction

In the study of electron/positron–proton (ep) collisions, a precise estimation of the dilepton¹ production cross-sections in the electroweak (EW) interaction is important since it could become a significant background for various physics analyses such as, for example, exclusive J/ψ or Υ production and new physics searches. So far only the generator LPAIR [1] has been used in experimental analyses to estimate the dilepton background [2]. The calculation of LPAIR is based on the diagrams of the photon–photon collision process [3], so-called *two-photon Bethe–Heitler* (2γ BH), corresponding to the diagrams of Fig. 1(a) or Fig. 2(a) with the photon contribution only. This process is dominant in most of the phase space. It is, however, expected that in the region of low invariant masses of the dilepton system, QED-Compton type (CO) diagrams (i.e. photon internal conversion process) as seen in Figs. 1(b) and 2(b) become dominant. In the high mass region, there is an additional interesting process, i.e. Z^0 production, which is implemented in the MC event generator EPVEC [4]. EPVEC, however, does not include 2γ BH nor CO diagrams. In the di- e channel, interference effects of the final state e^-e^- or e^+e^+ should also be taken into account, which are included neither in LPAIR nor in EPVEC.

In this paper, a new MC event generator GRAPE-Dilepton for the dilepton production in ep collisions is presented. The FORTRAN code to calculate the Feynman amplitudes is generated by GRACE [5] which is an automatic calculation system. GRACE has been used mainly for e^+e^- interactions so far. This is the first time for GRACE to be applied to the case where there is a composite particle (i.e. proton) in the initial state. GRAPE stands for a *GRACE-based generator for Proton–Electron collisions*.

This generator has the following features.

- The cross-section calculation is based on the exact matrix elements in the electroweak theory at tree level. Not only 2γ BH but also the dilepton productions via γZ^0 and $Z^0 Z^0$ collisions are taken into account. CO and Z^0 on/off-shell production are also included. Interference effects of the final state $e^\pm e^\pm$ are taken into account in the di- e channel. It is possible to select any sub-set of diagrams in the calculation.
- All fermion masses are kept non-zero both in the matrix elements and in the kinematics, which makes it possible to use this program with arbitrary small scattering angles of e^\pm and/or small invariant masses of dilepton down to the kinematical limits.
- The calculation of the proton vertex covers the whole kinematical region by dividing it into 3 categories of elastic, quasi-elastic and DIS (Deep Inelastic eq Scattering) processes, as described in the next section in detail.
- Both of Initial State Radiation (ISR) and Final State Radiation (FSR) can be included.

¹ The word *dilepton* represents di-electron (di- e), di-muon (di- μ) and di-tau (di- τ) in this paper.

2. Physics aspects

This generator simulates the ep interaction: $e_{(\text{in})}^{\pm} p_{(\text{in})} \rightarrow e^{\pm} l^{+} l^{-} X$ where $e_{(\text{in})}^{\pm}$ and $p_{(\text{in})}$ indicate the electron/positron and the proton in the initial state, respectively, e^{\pm} and $l^{+} l^{-}$ are the scattered electron/positron and the produced dilepton, respectively. The relevant processes are classified into 3 categories using the negative momentum transfer squared at the proton vertex (Q_p^2) and the invariant mass of the hadronic system (M_{had});

$$Q_p^2 \stackrel{\text{def}}{=} -\{p_{e^{\pm}(\text{in})} - (p_{e^{\pm}} + p_{l^{+}} + p_{l^{-}})\}^2, \quad (1)$$

$$M_{\text{had}}^2 \stackrel{\text{def}}{=} \{(p_{e^{\pm}(\text{in})} + p_{p(\text{in})}) - (p_{e^{\pm}} + p_{l^{+}} + p_{l^{-}})\}^2, \quad (2)$$

where $p_{e^{\pm}(\text{in})}$ and $p_{p(\text{in})}$ are the 4-momenta of the incoming lepton and the proton after ISR, respectively. $p_{e^{\pm}}$ and $p_{l^{\pm}}$ are those of the scattered lepton and the produced leptons before FSR, respectively. The 3 categories are

- $M_{\text{had}} = M_p$ (elastic),
- $Q_p^2 < Q_{\text{min}}^2$ or $M_p + M_{\pi^0} < M_{\text{had}} < M_{\text{cut}}$ (quasi-elastic),
- $Q_p^2 > Q_{\text{min}}^2$ and $M_{\text{had}} > M_{\text{cut}}$ (DIS),

where M_p and M_{π^0} are the masses of the proton and the neutral pion, respectively. Q_{min}^2 is set to around 1 GeV depending on the Parton Density Function (PDF) used in the DIS process. The recommended value for M_{cut} is 5 GeV.

For the elastic process, the diagrams in Fig. 1 are calculated with the following dipole form factor for the proton–proton–photon vertex ($\Gamma_{pp\gamma}^{\mu}$) with the on-shell proton. The general form of the elastic proton vertex can be written as

$$\Gamma_{pp\gamma}^{\mu} = e_p \left(F_1(Q_p^2) \gamma^{\mu} + \frac{\kappa_p}{2M_p} F_2(Q_p^2) i\sigma^{\mu\nu} q_{\nu} \right), \quad (3)$$

where e_p indicates the electric charge of the proton, q is the 4-momentum transfer at the proton vertex ($q^2 = -Q_p^2$), $F_1(Q_p^2)$ and $F_2(Q_p^2)$ are the 2 independent form factors, and κ_p is the anomalous magnetic moment of the proton (see, for example, [6]). The electric and magnetic form factors $G_E^p(Q_p^2)$ and $G_M^p(Q_p^2)$, respectively are defined as follows,

$$\begin{pmatrix} G_E^p(Q_p^2) \\ G_M^p(Q_p^2) \end{pmatrix} = \begin{pmatrix} F_1(Q_p^2) - \frac{\kappa_p Q_p^2}{4M_p^2} F_2(Q_p^2) \\ F_1(Q_p^2) + \kappa_p F_2(Q_p^2) \end{pmatrix}. \quad (4)$$

Using the Gordon decomposition and the scaling law of the form factor,

$$G_E^p(Q_p^2) = G_M^p(Q_p^2) / |\mu_p|, \quad (5)$$

the following formula which is used in this program is obtained,

$$\Gamma_{pp\gamma}^{\mu} = e_p \left(\mu_p G_E^p(Q_p^2) \gamma^{\mu} - \frac{(p_{p(\text{in})}^{\mu} + p_{p(\text{out})}^{\mu})}{2M_p} \frac{\kappa_p}{1 + \frac{Q_p^2}{4M_p^2}} G_E^p(Q_p^2) \right), \quad (6)$$

where $\mu_p = (1 + \kappa_p) \mu_B$, μ_B is the Bohr magneton, and $p_{p(\text{out})}$ indicates the 4-momentum of the scattered proton. $G_E^p(Q_p^2)$ is calculated according to the formula of the dipole fit,

$$G_E^p(Q_p^2) = \left(1 + \frac{Q_p^2}{0.71 \text{ GeV}^2} \right)^{-2}. \quad (7)$$

The only difference between the elastic and the quasi-elastic processes is the treatment of the proton vertex and the simulation of the hadronic final state. The quasi-elastic proton vertex can be described using the hadron tensor in the following form assuming parity and current conservation (for example, see [6]),




Fig. 1. Feynman diagrams included in the (quasi)-elastic process. $e = \{e^+, e^-\}$, $l^\pm = \{e^\pm, \mu^\pm, \tau^\pm\}$. N means a (dissociated) proton or a nucleon resonance.

$$W^{\mu\nu} = W_1 \left(-g^{\mu\nu} + \frac{q^\mu q^\nu}{q^2} \right) + W_2 \frac{1}{M_p^2} \left(p_{p(\text{in})}^\mu - \frac{p_{p(\text{in})} \cdot q}{q^2} q^\mu \right) \left(p_{p(\text{in})}^\nu - \frac{p_{p(\text{in})} \cdot q}{q^2} q^\nu \right). \quad (8)$$

$W_1(Q_p^2, M_{\text{had}})$ and $W_2(Q_p^2, M_{\text{had}})$ are the electromagnetic proton structure functions. The hadron tensor is contracted with the lepton tensor $L^{\mu\nu}$ numerically to obtain the cross-section,

$$d\sigma \sim L_{\mu\nu} W^{\mu\nu}. \quad (9)$$

In this version, W_1 and W_2 are parameterized with Brasse et al. [7] for $M_{\text{had}} < 2$ GeV (the proton resonance region), and with ALLM97 [8] for $M_{\text{had}} > 2$ GeV. These two parameterizations are based on fits to the experimental data on the measurement of the total γ^*p cross-sections. The exclusive hadronic final state is generated using the MC event generator SOPHIA [9] in the event generation step.

For the DIS process with the Quark Parton Model, the diagrams in Fig. 2 are calculated. PDFLIB [10] is linked to obtain parton densities with Q_p^2 as a QCD scale. The simulation of the proton remnant and the hadronization are performed by PYTHIA [11]. It should be noted that the lowest order calculation in this process is valid only for the region of

$$u \stackrel{\text{def}}{=} |\{p_{q(\text{in})} - (p_{l^+} + p_{l^-})\}^2| \gtrsim 25 \text{ GeV}^2, \quad (10)$$

where $p_{q(\text{in})}$ is the 4-momentum of the incoming quark. The value of u corresponds to the virtuality of the u -channel quark in the diagrams in Fig. 2(b), (c). When it is nearly or smaller than 25 GeV^2 , the lowest order calculation is not correct as explained in [4] since QCD corrections become large. In this case, the dilepton




Fig. 2. Feynman diagrams included in the DIS process. $e = \{e^+, e^-\}$, $l^\pm = \{e^\pm, \mu^\pm, \tau^\pm\}$ and $q = \{u^{(-)}, d^{(-)}, s^{(-)}, c^{(-)}, b^{(-)}, t^{(-)}\}$.

production should be treated as Drell–Yan process between the proton and the resolved photon from the beam lepton, which is not implemented in this program. The cut: $u > 25 \text{ GeV}^2$ is explicitly applied in this program if the diagrams other than BH are included.

The effect of ISR is included in the cross-section calculation using the structure function method described in [12], where the momentum transfer squared on the beam lepton, i.e. $\{p_{e^\pm(\text{in})} - p_{e^\pm}\}^2$ is used as a QED scale. When ISR turns on, the correction for the photon self energy, i.e. the vacuum polarization, is included according to the parameterization in [13] by modifying photon propagators. FSR is performed by PYTHIA using the parton shower method when the event is generated.

3. Program structure

Physics events are generated with the 2 steps; the MC integration step by the executable: `integ` and the event generation step by the executable: `spring`, as illustrated in Fig. 3. In both steps, the program is controlled by an ASCII file: `grape.cards`. The file is read by the executables with help of `FFREAD` [14]. The contents of `grape.cards` are explained in the next section.

In the integration step by the executable: `integ`, an effective total cross-section (in unit of pb) and probability distributions are calculated by BASES [15]. The results are stored in a file: `bases.rz` which has the Ntuple format provided by the HBOOK package [16]. At the same time, the information related to the convergency status of the integration is output into an ASCII file: `bases.result`.

In the event generation step by the executable: `spring`, unweighted events are generated. This is done by an routine: `SPRING` [15] according to the probability distributions in `bases.rz`. The results of the event generation are stored in the PYTHIA common block `/PYJETS/`. After filling `/PYJETS/`, `spring` calls a routine: `USRSTR` in which user specific procedures are put. Its template is found in the appendix. The event information in `/PYJETS/` is also available in a Ntuple file: `grp.rz`.




Fig. 3. Flowchart for the program structure.

The calculated cross-section is found in `bases.result` or at the end of the standard output from `spring`. The status of the event generation is output into an ASCII file: `spring.result`. Looking at the file, users should find a reasonable agreement between generated distributions by `spring` and calculated ones by `integ`. The procedure to make the executables is described in the README file.

4. Input data cards

The input data in `grape.cards` are explained in this section. All of the items are optional and are set to default values if not specified. Default values are written in the brackets starting with D=. The items are not explicitly displayed in case that they are the only one for their cards.

- **KFLBEAM**

KF code of the lepton beam (INTEGER, D=-11); 11: electron, -11: positron.

- **EPOL** P θ ϕ

Polarization of the lepton beam (REAL);

P = degree of the polarization in the range $[-1, 1]$ (D=0.),

θ = polar angle of the polarization vector in degree (D=0.),

ϕ = azimuthal angle of the polarization vector in degree (D=0.).

The positive direction of the z-axis on the polarization vector is in the direction of the lepton beam.

- **EBEAM**

Lepton beam momentum in MeV/c (REAL, D=27520.).

- **PBEAM**

Proton beam momentum in MeV/c (REAL, D=820000.).

- **PROCESS**

Process type of the proton vertex (INTEGER, D=1);

1: elastic, 2: quasi-elastic, 3: DIS.

- **LPAIR**

Dilepton channel (INTEGER, D=2); 1: di- e , 2: di- μ , 3: di- τ .

Table 1
Possible combinations of **QFLV** and **MERGE**

QFLV	MERGE	Quarks	BH	QED/EW/CO/Z ⁰
1	1234	$u + \bar{u} + d + \bar{d}$	Yes	No
1	123456	$u + \bar{u} + d + \bar{d} + s + \bar{s}$	Yes	No
1	12345678	$u + \bar{u} + d + \bar{d} + s + \bar{s} + c + \bar{c}$	Yes	No
1	1234567890	$u + \bar{u} + d + \bar{d} + s + \bar{s} + c + \bar{c} + b + \bar{b}$	Yes	No
1	17	$u + c$	Yes	Yes
2	28	$\bar{u} + \bar{c}$	Yes	Yes
3	35	$d + s$	Yes	Yes
4	46	$\bar{d} + \bar{s}$	Yes	Yes
3	359	$d + s + b$	Yes	Yes
4	460	$\bar{d} + \bar{s} + \bar{b}$	Yes	Yes

- **ISR**

Initial state radiation flag for the beam lepton (INTEGER, D=1); 0: off, 1: on.

- **QFLV**

Scattered quark in the DIS process (INTEGER, D=1);

1: u , 2: w , 3: d , 4: \bar{d} , 5: s , 6: \bar{s} , 7: c , 8: \bar{c} , 9: b , 10: \bar{b} , 11: t , 12: \bar{t} .

- **MERGE**

Merging mode in the DIS process (INTEGER, D=0); 0: off.

In some cases, contributions from different quarks can be included in the cross-section calculation adding the parton densities if the mass difference is negligible. The possible combinations of **QFLV** and **MERGE** are written in Table 1. The mass of the quark specified with **QFLV** is used in the amplitude and the kinematics calculations.

- **NGROUP**

Author group described in the PDFLIB manual (INTEGER, D=5).

- **NSET**

PDF set described in the PDFLIB manual (INTEGER, D=5).

The default is GRV94(LO).

- **GRASEL**

Feynman diagram selection (INTEGER, D=3);

- 1: 2- γ Bethe–Heitler (without $e^\pm e^\pm$ interference in case of di- e),
- 2: 2- γ Bethe–Heitler (including $e^\pm e^\pm$ interference in case of di- e),
- 3: QED diagrams (i.e. all the diagrams except for the Z^0 contribution),
- 4: EW diagrams (i.e. all the diagrams),
- 13: QED-Compton type diagrams only,
- 14: Z^0 production diagrams only.

In case of di- μ , τ , the first and the second selections give the same result.

- **ITMX1**

Number of iterations in the grid optimization step of BASES (INTEGER, D=4).

This should be larger than 2.

- **ITMX2**

Number of iterations in the integration step of BASES (INTEGER, D=10).

This should be larger than 5.

- **NCALL**
Number of sampling points in each iteration of BASES (INTEGER, D=1000000).
This should be large so that any *accuracy* of each iteration in the integration step of BASES is better than 0.5%.
- **NGEN**
Number of events to be generated by spring (INTEGER, D=100).
- **NMOD** N_{mod}
Printing a message per N_{mod} events in the event generation (INTEGER, D=1000).
- **PSISR**
Switch for the initial state parton shower by PYTHIA (INTEGER, D=1); 0: off, 1: on.
This has an effect only on event generations of the DIS process. No effect on elastic and quasi-elastic events.
This item is copied to MSTP (61) in the PYTHIA common block /PYPARS/.
- **PSFSR**
Switch for the final state parton shower by PYTHIA (INTEGER, D=1); 0: off, 1: on.
This item is copied to MSTP (71) in the PYTHIA common block /PYPARS/.
- **PSBRA**
Parton shower branchings in PYTHIA (INTEGER, D=2);
1: QCD, 2: QCD + QED.
This item is copied to MSTJ (41) in the PYTHIA common block /PYDAT1/.
- **PSSUP**
Suppression of the PYTHIA parton shower (INTEGER, D=0); 0: off, ≥ 1 : on.
This item is copied to MSTJ (40) in the PYTHIA common block /PYDAT1/.
- **PYDECAY**
Switch for fragmentation and decay in PYTHIA (INTEGER, D=1); 0: off, 1: on.
No effect on elastic and quasi-elastic events.
This item is copied to MSTP (111) in the PYTHIA common block /PYPARS/.
- **PRIPT**
Primordial k_t distribution in the proton (INTEGER, D=1);
0: off, 1: gaussian, 2: exponential.
No effect on elastic and quasi-elastic events. This item is copied to MSTP (91) in the PYTHIA common block /PYPARS/.
- **PYLIST**
Printing the contents of /PYJETS/ (LOGICAL, D=TRUE).
- **NLIST**
Number of events whose /PYJETS/ is printed out (INTEGER, D=10).
- **NTPYT**
Output of generated events into a Ntuple file: `grp.rz`
from the PYTHIA common block /PYJETS/ (LOGICAL, D=FALSE).
The meanings of the Ntuple variables are in the following.
 - `npv`: Number of particles (integer),
 - `px(1:npv)`, `py(1:npv)`, `pz(1:npv)`: x-, y-, z-component of momentum in GeV/c (real*4),
 - `pe(1:npv)`: Energy in GeV (real*4),
 - `pm(1:npv)`: Mass in GeV (real*4)
 - `kf(1:npv)`: KF code (integer),
 - `sta(1:npv)`: Status code (integer),
 - `mot(1:npv)`: Line number of the mother particle (integer).
- **Q2RNGME** Min Max
Range for the negative momentum transfer squared at the electron vertex Q_e^2 without ISR (REAL), i.e.
 $Q_e^2 = -\{p_{e^\pm(\text{in})} - p_{e^\pm}\}^2$ where $p_{e^\pm(\text{in})}$ is a 4-momentum of the incoming lepton after ISR.

Min = the minimum in GeV^2 ($D=0$.),

Max = the maximum in GeV^2 ($D=1$. E20).

In case of di- e with $e^\pm e^\pm$ interference, smaller one of the two Q_E^2 values is used.

- **Q2RNGOB** Min Max

Range for the negative momentum transfer squared at the electron vertex Q_e^2 including ISR (REAL), i.e.

$Q_e^2 = -\{p_{e^\pm(\text{in})} - p_{e^\pm}\}^2$ where $p_{e^\pm(\text{in})}$ is a 4-momentum of the incoming lepton before ISR.

Min = the minimum in GeV^2 ($D=0$.),

Max = the maximum in GeV^2 ($D=1$. E20).

In case of di- e with $e^\pm e^\pm$ interference, smaller one of the two Q_e^2 values is used.

- **MHAD** Min Max

Range for the mass of the hadronic system M_{had} (REAL);

Min = the minimum in GeV ($D=1$. 08),

Max = the maximum in GeV ($D=1$. E20).

No effect on elastic events.

- **Q2P** Min Max

Range for the negative momentum transfer squared at the proton vertex Q_p^2 (REAL);

Min = the minimum in GeV^2 ($D=0$.),

Max = the maximum in GeV^2 ($D=1$. E20).

In case of the DIS process, Q_p^2 is used as a QCD scale for PDF.

- **THMIN** $\theta_{\min}^{(1)}$ $\theta_{\min}^{(2)}$ $\theta_{\min}^{(3)}$ $\theta_{\min}^{(4)}$
- **THMAX** $\theta_{\max}^{(1)}$ $\theta_{\max}^{(2)}$ $\theta_{\max}^{(3)}$ $\theta_{\max}^{(4)}$
- **EMIN** $E_{\min}^{(1)}$ $E_{\min}^{(2)}$ $E_{\min}^{(3)}$ $E_{\min}^{(4)}$
- **EMAX** $E_{\max}^{(1)}$ $E_{\max}^{(2)}$ $E_{\max}^{(3)}$ $E_{\max}^{(4)}$
- **PMIN** $P_{\min}^{(1)}$ $P_{\min}^{(2)}$ $P_{\min}^{(3)}$ $P_{\min}^{(4)}$
- **PMAX** $P_{\max}^{(1)}$ $P_{\max}^{(2)}$ $P_{\max}^{(3)}$ $P_{\max}^{(4)}$
- **PTMIN** $P_{t\min}^{(1)}$ $P_{t\min}^{(2)}$ $P_{t\min}^{(3)}$ $P_{t\min}^{(4)}$
- **PTMAX** $P_{t\max}^{(1)}$ $P_{t\max}^{(2)}$ $P_{t\max}^{(3)}$ $P_{t\max}^{(4)}$

(1): for scattered proton or quark,

(2): for scattered e^\pm ,

(3): for produced l^\mp ,

(4): for produced l^\pm .

The above 8 data cards are used for describing the detector cut in the laboratory frame (REAL). Each final state particle is required to satisfy the following,

$$\theta_{\min}^{(i)} < \theta < \theta_{\max}^{(i)} \quad \text{AND} \quad E_{\min}^{(i)} < E < E_{\max}^{(i)} \quad \text{AND}$$

$$P_{\min}^{(i)} < P < P_{\max}^{(i)} \quad \text{AND} \quad P_{t\min}^{(i)} < P_t < P_{t\max}^{(i)}$$

where θ (degree), E (GeV), P (GeV/c) and P_t (GeV/c) indicate polar angle, energy, momentum and transverse momentum, respectively. The default values correspond to not applying this cut.

- **THPTMCT** θ_{\min} θ_{\max}

- **PTMXCT** $P_{t\min}$ $P_{t\max}$

Using the above 2 data cards, final state leptons are required to satisfy the following (REAL),

$$\theta_{\min} < \theta^M < \theta_{\max} \quad \text{AND} \quad P_{t\min} < P_t^M < P_{t\max}$$

where P_t^M (GeV/c) indicates the maximum transverse momentum among the 3 final state leptons (e^\pm, l^\mp, l^\pm), and θ (degree) is the polar angle of the lepton with P_t^M . The default values correspond to not applying this cut.

- **MASSLL** Min1 Max1 Min2 Max2

Range for the mass of the produced dilepton system (REAL);

Min1 = the minimum in GeV ($D=0$),

Max1 = the maximum in GeV ($D=1$. E20).

In case of di- μ , τ , Min2 and Max2 are not used.

In case of di- e with $e^\pm e^\pm$ interference, there are two masses; $M_{e^+e^-}^{(1)}$, $M_{e^+e^-}^{(2)}$ ($M_{e^+e^-}^{(1)} < M_{e^+e^-}^{(2)}$), and they are required to satisfy the following,

$$\text{Min1} < M_{ee}^{(1)} < \text{Max1} \quad \text{AND} \quad \text{Min2} < M_{ee}^{(2)} < \text{Max2}.$$

- **MASSELL** Min Max

Range for the mass of the final state lepton system of $e^\pm l^\mp l^\pm$ (REAL);

Min = the minimum in GeV ($D=1$.),

Max = the maximum in GeV ($D=1$. E20).

- **MASSQLL** Min Max

Range for the mass of the scattered quark and produced dilepton system of ql^+l^- (REAL);

Min = the minimum in GeV ($D=5$.),

Max = the maximum in GeV ($D=1$. E20).

This cut has an effect only on the DIS process.

In case of di- e with $e^\pm e^\pm$ interference, smaller one of the 2 values is used.

- **IVISI** N_{visi}

- **THEVMIN** $\theta_{\min}^{(1)}$ $\theta_{\min}^{(2)}$ $\theta_{\min}^{(3)}$ $\theta_{\min}^{(4)}$

- **THEVMAX** $\theta_{\max}^{(1)}$ $\theta_{\max}^{(2)}$ $\theta_{\max}^{(3)}$ $\theta_{\max}^{(4)}$

- **EVMIN** $E_{\min}^{(1)}$ $E_{\min}^{(2)}$ $E_{\min}^{(3)}$ $E_{\min}^{(4)}$

- **EVMAX** $E_{\max}^{(1)}$ $E_{\max}^{(2)}$ $E_{\max}^{(3)}$ $E_{\max}^{(4)}$

- **PTVMIN** $Pt_{\min}^{(1)}$ $Pt_{\min}^{(2)}$ $Pt_{\min}^{(3)}$ $Pt_{\min}^{(4)}$

- **PTVMAX** $Pt_{\max}^{(1)}$ $Pt_{\max}^{(2)}$ $Pt_{\max}^{(3)}$ $Pt_{\max}^{(4)}$

(1): for scattered proton or quark,

(2): for scattered e^\pm ,

(3): for produced l^\mp ,

(4): for produced l^\pm .

The above 6 data cards are used for describing the detector cut in the laboratory frame (REAL except for N_{visi} : INTEGER). N_{visi} particle(s) are(is) required to satisfy the following,

$$\theta_{\min}^{(i)} < \theta < \theta_{\max}^{(i)} \quad \text{AND} \quad E_{\min}^{(i)} < E < E_{\max}^{(i)} \quad \text{AND} \quad Pt_{\min}^{(i)} < Pt < Pt_{\max}^{(i)},$$

where θ (degree), E (GeV) and Pt (GeV/c) indicate polar angle, energy and transverse momentum, respectively. As for N_{visi} , $D=-1$, which corresponds to not applying this cut. The test run at the end of this paper is instructive for understanding this cut.

5. Summary

A new Monte Carlo generator for dilepton production in the framework of the electroweak theory was presented. The whole kinematical region on the proton vertex is covered. This generator can be used for quantitative and precise estimations of processes which come in addition to the two-photon Bethe–Heitler contributions.

Acknowledgements

I am very thankful to Prof. Y. Shimizu, Prof. S. Yamada and Prof. K. Tokushuku for their useful comments and various supports. I shall never forget the collaboration with the members of the Minami-Tateya group at KEK;

J. Fujimoto, T. Ishikawa, T. Kaneko, K. Kato, S. Kawabata, T. Kon, Y. Kurihara, H. Tanaka and T. Watanabe. I also wish to thank the conveners of the working group on QED radiative effects in HERA Monte Carlo Workshop at DESY; L. Favart, S. Schlenstedt and H. Spiesberger for the helpful discussions. This work was supported in part by the Japanese Ministry of Education, Science and Culture (the Monbusho) and its grant for Scientific Research (No. 11206203).

Appendix A. Event store in /PYJETS/

Each line in the PYTHIA event store has the following meaning according to the PYTHIA convention.

Line number	Meaning
1, 2	Beam particles (1: p , 2: e^\pm)
3, 4	Partons from the beam particles <i>before</i> ISR (3: from p , 4: from e^\pm)
5, 6	Partons from the beam particles <i>after</i> ISR; the initial state in the matrix element calculation (5: from p , 6: from e^\pm)
7, 8, 9, 10	Final state particles <i>before</i> FSR; the final state in the matrix element calculation (7: p , 8: e^\pm , 9: l^\mp , 10: l^\pm)
11~	Final state particles <i>after</i> FSR, fragmentation and decay

In case of the (quasi-)elastic process,

- a parton from the beam proton is the proton itself, i.e. the 1st and the 3rd lines are the same,
- the beam proton always makes no ISR, so that the 1st, 3rd and 5th lines are the same,
- the 2nd and the 4th lines are also the same.

In case of the DIS process with both ISR and FSR, all of the lines have different contents in general.

As for di- e events, there are 2 identical particles in the final state. In GRAPE, those 2 particles are distinguished in the following way;

- in case of $e^\pm e^\pm$ interference *off*, a particle stored in the 8th line is a scattered lepton, and one in the 10th line comes from the $2\text{-}\gamma$ collision (i.e. a produced lepton),
- in case of $e^\pm e^\pm$ interference *on*, a lepton stored in the 8th line has smaller transverse momentum than that of a lepton in the 10th line.

Appendix B. Routines/function related to users

- **Function** DRN(ISEED)
provides uniform random numbers. All other random number routines are linked to this one. This routine is stored in the BASES library (libbases.a).
- **Subroutine** DRNSET(ISEED)
performs an initialization for DRN(ISEED). This is also stored in libbases.a.
- **Subroutine** READ_CARDS(LUN, filename)
reads input data cards from grape.cards.
- **Subroutine** SETMAS
provides masses/widths of particles and the QED coupling constant.

- **Subroutine** USRSTR(Ievt, Ngen)

can be modified by users to access the information on generated events. User initialization/termination procedures are also put in this routine.

Appendix C. User event storing routine: USRSTR

This routine is called ($N_{\text{gen}} + 2$) times where N_{gen} is the number of generated events. The 1st call is for the user initialization and the last one for the user termination phase. The following is the template file prepared in the GRAPE package (usrstr.f).

```

      subroutine USRSTR(Ievt, Ngen)
      implicit NONE
*----- Arguments -----
      integer Ievt, Ngen
* Ngen : # of events to be generated
* Ievt : Counter --- < 1      ==> Initialization   phase
*              1 - Ngen ==> Event generation phase
*              > Ngen      ==> Termination       phase
*-----
*----- PYTHIA common -----
      integer          N,NPAD, K(4000,5)
      double precision P(4000,5), V(4000,5)
      common /PYJETS/ N,NPAD,K,P,V          !!! Event Record !!!

      integer          MINT(400)
      double precision VINT(400)
      common /PYINT1/ MINT,VINT
* (See PYTHIA manual for details.)
*-----
*----- Local variables -----
      integer          LUN1,    LUN2,    LUN3
      parameter (LUN1=41, LUN2=42, LUN3=43)
* (You can use the above logical unit numbers.)
*-----

***** Initialization of USER Event Storing *****
      if (Ievt .LT. 1) then

          endif
*****

***** <<< USER Event Storing >>> *****
      if ((Ievt .GE. 1).and.(Ievt .LE. Ngen)) then

          endif
*****
***** Termination of USER Event Storing *****
      if (Ievt .GT. Ngen) then

          endif

```

```
return  
end
```

References

- [1] S.P. Baranov, O. Dünger, H. Shoohtari, J.A.M. Vermaseren, LPAIR — A Generator for Lepton Pair Production, in: W. Buchmüller, G. Ingelman (Eds.), Proc. of the Workshop on Physics at HERA, Vol. 3, DESY, 1992, p. 1478.
- [2] D. Hoffmann, L. Favart, Lepton Pair Monte Carlo Generators for HERA Physics, in: A.T. Doyle, G. Grindhammer, G. Ingelman, H. Jung (Eds.), Proc. of the Workshop on Monte Carlo Generators for HERA Physics, DESY, 1999.
- [3] J.A.M. Vermaseren, Nucl. Phys. B 229 (1983) 347.
- [4] U. Baur, J.A.M. Vermaseren, D. Zeppenfeld, Nucl. Phys. B 375 (1992) 3.
- [5] T. Ishikawa, T. Kaneko, K. Kato, S. Kawabata, Y. Shimizu, H. Tanaka, KEK Report 92-19 (1993).
- [6] F. Halzen, A.D. Martin, Quarks and Leptons: An Introductory Course in Modern Particle Physics, John Wiley & Sons, 1984.
- [7] F.W. Brasse et al., Nucl. Phys. B 110 (1976) 413.
- [8] H. Abramowicz, A. Levy, DESY-97-251, hep-ph/9712415.
- [9] A. Mücke et al., Comput. Phys. Commun. 124 (2000) 290.
- [10] H. Plochow-Besch, Comput. Phys. Commun. 75 (1993) 396.
- [11] T. Sjöstrand, Comput. Phys. Commun. 82 (1994) 74.
- [12] J. Fujimoto, M. Igarashi, N. Nakazawa, Y. Shimizu, K. Tobimatsu, Prog. Theor. Phys. Suppl. 100 (1990) 285.
- [13] H. Burkhardt, B. Pietrzyk, Phys. Lett. B 356 (1995) 398.
- [14] R. Brun et al., FFREAD User Guide and Reference Manual, CERN DD/US/71, CERN Program Library I302, February 1987.
- [15] S. Kawabata, Comput. Phys. Commun. 88 (1995) 309.
- [16] R. Brun, D. Lienart, HBOOK User Guide — Version 4, CERN Program Library Y250, 1988.
- [17] M. Glück, E. Reya, A. Vogt, Z. Phys. C 67 (1995) 433.

TEST RUN INPUT

An example of the input data cards

One example is presented with the following condition;

- process: $e^+q \rightarrow e^+q\mu^+\mu^-$ ($q = \{ \overset{(-)}{u}, \overset{(-)}{d}, \overset{(-)}{s} \}$ with GRV94(LO) [17]),
- BH diagrams only,
- 70% polarization of the e^+ beam in the direction of the proton beam,
- cuts: (1) & (2) & (3) & (4),
 - (1) $Q_p^2 > 1 \text{ GeV}^2$ & $M_{\text{had}} > 5 \text{ GeV}$,
 - (2) for scattered q , $\theta > 10^\circ$ & $P_t > 15 \text{ GeV}/c$,
 - (3) (invariant mass of $\mu^+\mu^-$) $> 4 \text{ GeV}$,
 - (4) 2 of the following 3 requirements for the final state leptons are satisfied,
 - for e^+ : $5^\circ < \theta < 175^\circ$ & $P_t > 5 \text{ GeV}$,
 - for μ^+ : $20^\circ < \theta < 160^\circ$ & $P_t > 3 \text{ GeV}$,
 - for μ^- : $20^\circ < \theta < 160^\circ$ & $P_t > 3 \text{ GeV}$.

Following is the corresponding grape.cards. The same file is put in the directory sample.

```

LIST
NCALL 1400000
C =====
C << Polarization of the Lepton Beam >>
C      (1)      (2)      (3)
EPOL      -0.7      0.      0.
C =====
C << Process in the Proton Vertex >> (1:elastic, 2:quasi-elastic, 3:DIS)
PROCESS      3
C =====
C << Produced Lepton-pair >> (1:di-e, 2:di-mu, 3:di-tau)
LPAIR      2
C =====
C << Scattered Quark in DIS >>
C (1:u, 2:u-bar, 3:d, 4:d-bar, 5:s, 6:s-bar, 7:c, ... , 12:t-bar)
QFLV      1
MERGE 123456
C =====
C << PDF set in DIS >> (See PDFLIB manual.)
NGROUP      5
NSET      5
C =====
C #####
C =====
C << Electroweak Dilepton Production >>
GRASEL      2
C =====
C #####
C =====
C << Mass Range for the Hadronic System >> (only for quasi-elastic and DIS)
MHAD      5.      300.

```

```

C -----
C << Q2 Range for the Proton Vertex >>
Q2P      1.      1.E20
C -----
C << Cuts for each Final-state Particle >>
C      <p/q>  <e+->  <l+->  <l+->
THMIN      10.      0.      0.      0.
THMAX     180.     180.     180.     180.
EMIN        0.      0.      0.      0.
EMAX       1.E20    1.E20    1.E20    1.E20
PMIN        0.      0.      0.      0.
PMAX       1.E20    1.E20    1.E20    1.E20
PTMIN      15.      0.      0.      0.
PTMAX      1.E20    1.E20    1.E20    1.E20
C -----
C << Mass cuts >>
MASSL      4.      1.E20
           0.      1.E20
C -----
C << Cuts for One or Some of the Final-state Particles >>
IVISI      2
C      <p/q>  <e+->  <l+->  <l+->
THEVMIN     0.      5.     20.     20.
THEVMAX     0.    175.    160.    160.
EVMIN       0.      0.      0.      0.
EVMAX       0.     1.E20    1.E20    1.E20
PTVMIN      0.      5.      3.      3.
PTVMAX      0.     1.E20    1.E20    1.E20
C =====
STOP

```

TEST RUN OUTPUT

A part of the standard output from `integ`

```
#####
#####
#####
##### GGGGGG RRRRRR A PPPPPP EEEEE +++++#
##### G G R R A A P P E +++++#
##### G R R A A P P E +++++#
##### G GGGGG R RRRR A A PPPPPP EEEEE +++++#
##### G G G R R AAAAAA P E +++++#
##### G G G R R A A P E +++++#
##### GGGG G R RR A A P EEEEEEE +++++#
#####
##### GRACE-based generator for Proton-Electron collisions +++++#
##### GRAPE-Dilepton_version1.1 +++++#
##### +++++#
##### Mar. 14 2000 +++++#
##### Comments/bug-report to Tetsuo ABE(tabe@post.kek.jp) +++++#
##### (abe@mail.desy.de) +++++#
#####
#####
<<<<<<<<< This is an INTEGRATION step. >>>>>>>>>

---> DIS process
      (Scattering of e and u quark)
---> Muon-pair production

      grace 2.1(5)

      (c)Copyright 1990-1998 Minami-Tateya Group (Japan)

>>> Graph selection
jselg =
1 : 0
2 : 0
3 : 0
4 : 0
5 : 0
6 : 0
7 : 0
8 : 0
9 : 1
10 : 0
11 : 0
12 : 0
13 : 1
14 : 0
15 : 0
16 : 0
17 : 0
18 : 0
19 : 0
20 : 0
21 : 0
22 : 0
23 : 0
24 : 0
25 : 0
```


=====> Start of Kinematics Initialization

***** Information (in Lab. frame) *****
(in unit of GeV)

```
P of electrons   = 27.5200
P of protons     = 820.000
Mass of electron = 5.10999E-04
Mass of proton   = 0.938272
sqrt(S)          = 300.444
P of CMS         = 792.480
E of CMS         = 847.521
gamma of CMS     = 2.82089
beta*gamma of CMS = 2.63770
```

<< Mass range for the hadronic system >>

```
Min. = 5.00000 GeV
Max. = 300.000 GeV
```

-----> PDFLIB Initialization started

```
***** PDFLIB Version: 7.09 Released on 970702 at 16.05
in the CERN Computer Program Library W5051 *****
***** Library compiled on 970702 at 16.05 *****
```

```
Parm = Nptype Ngroup Nset
Val = 1.0000 5.0000 5.0000
```

```
Nptype = 1 Ngroup = 5 Nset = 5 Name = "GRV94-L0" CrMode = -1
Nfl = -5, L0 = 1, Tmas = 180.00 GeV/c**2
QCDL4 = 0.2000 GeV, QCDL5 = 0.1530 GeV
Xmin = 0.10E-05, Xmax = 0.99999E+00,
Q2min = 0.400 (GeV/c)**2, Q2max = 0.10E+07 (GeV/c)**2
```

-----> PDFLIB Initialization finished

-----> ISR for incoming lepton using Structure Function method

=====> End of Kinematics Initialization

>>> e+ beam

Date: 0/ 4/12 21:29

```
*****
*
*      BBBBBBB AAAA SSSSSS EEEEE EEEEE SSSSSS
*      BB BB AA AA SS SS EE SS SS
*      BB BB AA AA SS EE SS
*      BBBBBBB AAAAAAA SSSSSS EEEEE SSSSSS
*      BB BB AA AA SS EE SS
*      BB BB AA AA SS SS EE SS SS
*      BBBB BB AA AA SSSSSS EEEEE SSSSSS
*
*
*      BASES Version 5.1
*      coded by S.Kawabata KEK, March 1994
*****
```

<< Parameters for BASES >>

- (1) Dimensions of integration etc.
 # of dimensions : Ndim = 9 (50 at max.)
 # of Wilds : Nwild = 7 (15 at max.)
 # of sample points : Ncall = 1399680(real) 1400000(given)
 # of subregions : Ng = 48 / variable
 # of regions : Nregion = 6 / variable
 # of Hypercubes : Ncube = 279936

- (2) About the integration variables

i	XL(i)	XU(i)	IG(i)	Wild
1	0.000000E+00	1.000000E+00	1	yes
2	0.000000E+00	1.000000E+00	1	yes
3	0.000000E+00	1.000000E+00	1	yes
4	0.000000E+00	1.000000E+00	1	yes
5	0.000000E+00	1.000000E+00	1	yes
6	0.000000E+00	1.000000E+00	1	yes
7	0.000000E+00	1.000000E+00	1	yes
8	0.000000E+00	1.000000E+00	0	no
9	0.000000E+00	1.000000E+00	0	no

- (3) Parameters for the grid optimization step

Max.# of iterations: ITMX1 = 4
 Expected accuracy : Acc1 = 0.2000 %

- (4) Parameters for the integration step

Max.# of iterations: ITMX2 = 10
 Expected accuracy : Acc2 = 0.0100 %

Date: 0/ 4/12 21:29

Convergency Behavior for the Grid Optimization Step

<- Result of each iteration ->				<- Cumulative Result		>- < CPU time >	
IT	Eff	R_Neg	Estimate	Acc %	Estimate(+/- Error)order	Acc % (H: M: Sec)	
1	2	0.00	4.586E-02	6.435	4.586406(+/-0.295122)E-02	6.435	0: 1:37.57
2	29	0.00	5.205E-02	0.295	5.203249(+/-0.015359)E-02	0.295	0: 7:23.86
3	37	0.00	5.196E-02	0.201	5.198557(+/-0.008625)E-02	0.166	0:13:52.62

Date: 0/ 4/12 21:29

Convergency Behavior for the Integration Step

<- Result of each iteration ->				<- Cumulative Result		>- < CPU time >	
IT	Eff	R_Neg	Estimate	Acc %	Estimate(+/- Error)order	Acc % (H: M: Sec)	
1	38	0.00	5.196E-02	0.194	5.196162(+/-0.010077)E-02	0.194	0:20:27.03
2	39	0.00	5.201E-02	0.198	5.198416(+/-0.007197)E-02	0.138	0:27: 0.89
3	38	0.00	5.201E-02	0.395	5.198705(+/-0.006793)E-02	0.131	0:33:34.96
4	39	0.00	5.199E-02	0.191	5.198759(+/-0.005605)E-02	0.108	0:40: 9.71
5	38	0.00	5.200E-02	0.195	5.198989(+/-0.004905)E-02	0.094	0:46:43.94
6	39	0.00	5.200E-02	0.193	5.199218(+/-0.004405)E-02	0.085	0:53:18.60
7	38	0.00	5.187E-02	0.193	5.197301(+/-0.004032)E-02	0.078	0:59:53.16
8	38	0.00	5.185E-02	0.193	5.195602(+/-0.003739)E-02	0.072	1: 6:27.22
9	38	0.00	5.205E-02	0.193	5.196769(+/-0.003505)E-02	0.067	1:13: 1.21
10	38	0.00	5.191E-02	0.196	5.196186(+/-0.003313)E-02	0.064	1:19:34.81

***** END OF BASES *****

<< Computing Time Information >>

```
(1) For BASES      H: M: Sec
    Overhead       : 0: 0: 0.12
    Grid Optim. Step : 0:13:52.62
    Integration Step : 1: 5:42.20
    Go time for all  : 1:19:34.94

(2) Expected event generation time
    Expected time for 1000 events : 0.40 Sec
```

Making bases.rz...

```
----> BASE1 : finished
----> BASE3 : finished
----> BASE4 : finished
----> BASE5 : finished
----> RANDM : finished
----> PLOTH : finished
----> PLOTB : finished
----> BSRSLT: finished
```

====> Directory : //bn

```
3 (N) BASES data(real*8)
1 (N) BASES data(integer*4)
2 (N) BASES data(real*4)
```

Suppressing decay of the following particles in PYTHIA,						
K_S0	eta	Lambda0	Sigma+	Sigma0	Sigma-	Xi0
Xi-	Omega-	D+	D0	D_s+	Lambda_c+	mu-
tau-	pi+	K+	K_L0	pi0		

```

Loading bases.rz...
---> BASE1 : finished
---> BASE3 : finished
---> BASE4 : finished
---> BASE5 : finished
---> RANDM : finished
---> PLOTH : finished
---> PLOTB : finished
---> BSRSLT : finished

```

```

*****
*****
**                                     **
**                                     **
**          *.....*                  Welcome to the Lund Monte Carlo! **
**      *:~::~!:~::~!:~::~!:~::~!*    PPP Y Y TTTT H H III A **
**      *:~::~!:~::~!:~::~!:~::~!*    P P Y Y T H H I AA **
**      *:~::~!:~::~!:~::~!:~::~!*    PPP Y T HHHH I AAAAA **
**      *:~::~!:~::~!:~::~!:~::~!*    P Y T H H I AA **
**      *:~::~!:~::~!:~::~!:~::~!*    P Y T H H III A A **
**      *:~::~!:~::~!:~::~!:~::~!* !! **
**      !! *:~::~!:~::~!:~::~!:~::~:* !! This is PYTHIA version 6.136 **
**      !! !* -><- * !! Last date of change: 30 Nov 1999 **
**      !! !! !! **
**      !! !! !! **
**      !! !! !! **
**      !! ep !! Disclaimer: this program comes **
**      !! !! !! without any guarantees. Beware **
**      !! pp !! of errors and use common sense **
**      !! e+e- !! when interpreting results. **
**      !! !! **
**      Copyright T. Sjostrand (1999) **
** **
** An archive of program versions and documentation is found on the web: **
** http://www.thep.lu.se/~torbjorn/Pythia.html **
** **
** When you cite this program, currently the official reference is **
** T. Sjostrand, Computer Physics Commun. 82 (1994) 74. **
** The supersymmetry extensions are described in **
** S. Mrenna, Computer Physics Commun. 101 (1997) 232 **
** Also remember that the program, to a large extent, represents original **
** physics research. Other publications of special relevance to your **
** studies may therefore deserve separate mention. **
** **
** Main author: Torbjorn Sjostrand; Department of Theoretical Physics 2, **
** Lund University, Solvegatan 14A, S-223 62 Lund, Sweden; **
** phone: + 46 - 46 - 222 48 16; e-mail: torbjorn@thep.lu.se **
** SUSY author: Stephen Mrenna, Physics Department, UC Davis, **
** One Shields Avenue, Davis, CA 95616, USA; **
** phone: + 1 - 530 - 752 - 2661; e-mail: mrenna@physics.ucdavis.edu **
** **

```


 1***** PYINIT: initialization of PYTHIA routines *****

```
=====
I                                     I
I      PYTHIA will be initialized for p+ on e+ user configuration      I
I                                     I
I                                     I
I      px (GeV/c)   py (GeV/c)   pz (GeV/c)   E (GeV)                   I
I      p+          0.000        0.000        820.000   820.001         I
I      e+          0.000        0.000       -27.520   27.520         I
I                                     I
I      corresponding to      300.444 GeV center-of-mass energy         I
I                                     I
=====
```

***** PYMAXI: summary of differential cross-section maximum search *****

```
=====
I                                     I   I
I      ISUB   Subprocess name           I   Maximum value   I
I                                     I   I
I      308    e+ uu^dd^ss^ -> e+ q m+ m-   I       5.1962D-02   I
I                                     I   I
=====
```

***** PYINIT: initialization completed *****

=====> START of SPRING at 1:21(13/ 4/ 0)

Number of generated events = 100

Event listing (summary)

I	particle/jet	KS	KF	orig	p_x	p_y	p_z	E	m
1	!p+!	21	2212	0	0.000	0.000	820.000	820.001	0.938
2	!e+!	21	-11	0	0.000	0.000	-27.520	27.520	0.001
3	!d!	21	1	1	-0.343	-0.058	168.409	168.409	0.000
4	!e+!	21	-11	2	0.000	0.000	-27.520	27.520	0.000
5	!d!	21	1	3	1.167	-0.401	138.481	138.487	0.000
6	!e+!	21	-11	4	0.000	0.000	-27.497	27.497	0.000
7	!d!	21	1	0	-10.149	16.228	96.119	98.007	0.004
8	!e+!	21	-11	0	0.009	0.020	-6.958	6.958	0.001
9	!mu-!	21	13	0	19.804	-7.523	-12.433	24.564	0.106
10	!mu+!	21	-13	0	-8.498	-9.126	34.256	36.455	0.106
11	e+	1	-11	8	0.009	0.020	-6.958	6.958	0.001
12	mu-	1	13	9	19.804	-7.523	-12.433	24.564	0.106
13	gamma	1	22	4	0.000	0.000	0.000	0.000	0.000
14	mu+	1	-13	10	-7.882	-8.464	31.770	33.810	0.106
15	gamma	1	22	2	0.000	0.000	0.000	0.000	0.000
16	(d)	A 12	1	7	-9.379	15.009	77.561	79.555	0.004
17	(g)	I 12	21	7	-1.386	0.556	21.044	21.097	0.000
18	(g)	I 12	21	3	-1.510	0.343	29.905	29.945	0.000
19	(uu_1)	V 11	2203	1	0.343	0.058	651.591	651.591	0.771

20 (string)	11	92	16	-11.932	15.967	780.101	782.188	53.513
21 (rho-)	11	-213	20	-2.614	3.803	19.032	19.599	0.763
22 eta	1	221	20	-5.726	8.838	49.702	50.808	0.547
23 (K*+)	11	323	20	-0.313	0.722	5.628	5.751	0.885
24 (K*-)	11	-323	20	-1.601	1.458	13.870	14.064	0.862
25 (rho+)	11	213	20	-0.642	0.548	18.730	18.778	1.033
26 pi0	1	111	20	-0.205	0.308	3.644	3.665	0.135
27 (rho-)	11	-213	20	-0.597	-0.035	42.069	42.081	0.796
28 (rho0)	11	113	20	-0.171	0.378	46.359	46.369	0.877
29 pi0	1	111	20	0.093	-0.222	22.348	22.349	0.135
30 K+	1	321	20	-0.371	0.069	95.962	95.964	0.494
31 Sigma0	1	3212	20	0.358	0.006	317.588	317.590	1.193
32 pi+	1	211	20	-0.143	0.094	145.170	145.170	0.140
33 pi-	1	-211	21	-1.237	1.578	6.910	7.196	0.140
34 pi0	1	111	21	-1.376	2.225	12.122	12.402	0.135
35 K+	1	321	23	-0.236	0.131	2.313	2.380	0.494
36 pi0	1	111	23	-0.078	0.591	3.315	3.370	0.135
37 (Kbar0)	11	-311	24	-0.822	0.679	8.481	8.562	0.498
38 pi-	1	-211	24	-0.779	0.779	5.389	5.502	0.140
39 pi+	1	211	25	-0.837	0.227	15.119	15.144	0.140
40 pi0	1	111	25	0.195	0.321	3.611	3.633	0.135
41 pi-	1	-211	27	-0.260	0.270	29.720	29.722	0.140
42 pi0	1	111	27	-0.337	-0.305	12.350	12.359	0.135
43 pi+	1	211	28	-0.149	-0.213	8.213	8.218	0.140
44 pi-	1	-211	28	-0.022	0.591	38.146	38.151	0.140
45 K_S0	1	310	37	-0.822	0.679	8.481	8.562	0.498
=====								
sum:		2.00		0.000	0.000	792.480	847.521	300.444

====> Directory : //grp
 =====> END of SPRING at 1:21(13/ 4/ 0)

1***** PYSTAT: Statistics on Number of Events and Cross-sections *****

I	I	I	I	I
I	Subprocess	I	Number of points	I
I		I		I
I		I		I
I		I		I
I		I		I
I	N:o Type	I	Generated	I
I		I	Tried	I
I		I		I
=====				
I		I		I
I	0 All included subprocesses	I	100	I
I	308 e+ uu^dd^ss^-> e+ q m+ m-	I	100	I
I		I		I
=====				

***** Fraction of events that fail fragmentation cuts = 0.00000 *****

```

***** << Total Cross-section >> *****
*
*      ( 5.196186 +- 0.003313 )E-02 pb
*
*****

```