1. The probability of a leap year selected at random contain 53					
Sunday is:					
53/ 366	(b) 1/7	(c) 2/7	(d) 53/365		
			marble is drawn at		
random. The prob	pability of draw	ing a black bal	lis:		
(a) 3/5	(b) 2/5	6 0/5	(d) 1/5		
3. The probability	y that it will rai	n tomorrow is (0.85. What is the		
probability that it	will not rain to	morrow			
(a) 0.25	(b) 0.145	4 3/20	(d) none of these		
			ed from the numbers		
(1, 2, 3,,15)) is a multiple o	of 4?			
1/5	(b) 4/5	(c) 2/15	(d) 1/3		
5. What are the t	otal outcomes	when we throw	v three coins?		
(a) 4					
			ed at random from the		
numbers (1,2,3,	35) is :				
(a) 12/35	(b) 11/35	(c) 13/35	none of these		
7. The sum of the					
(a) 2	(c)	0 (d) non	e of these.		
8. The following	probabilities a	re given; choos	se the correct answer		
for that which is r	not possible.				
(a) 0.15	(b) 2/7	7/5	(d) none of these.		
			an the probability of		
getting at least tw	vo heads, is:				
(a) 1/4	4) 3/8	(c) ½	(d) 1/8		
10. A letter is ch	osen at randor	m from the lette	ers of the word		
ASSASSINATION	N�. The prob	ability that the	letter chosen has:		
(a) 6/13	(b) 7/13	(c) 1	letter chosen has: none of these.		
11. A dice is throw	wn. Find the pr	obability of get	ting an even number.		
(A) 2/3	(B) 1	(C) 5/6	(-) 1/2		
12. Two coins are thrown at the same time. Find the probability of					
getting both head		, -			
(A) 3/4 (S) 1/4	(C) 1/2	(D) 0			
13. Two dice are thrown simultaneously. The probability of getting a					

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1	/9 (D) 4/9	
14. 100 cards are numbered from 1 to 100. Find the probability of					
(A) 3/4	me number. (B) 27/50	(6,1)	' 4	(D) 29/100	
15. A bag contains 5 red balls and some blue balls .If the probability of drawing a blue ball is double that of a red ball, then the number of blue balls in a bag is:					
(A) 5	10	(C) 15	(D) 20		
	random from ve bulb is:	this box.		olbs. One bulb is obability that it is	
mixed thoro the probabil		rd is draw mber on c	n from this l ard is a perf	-	1
18. What is (A) 1/7	the probability 53/366	•	•	ys in a leap year? (D) 7/366	
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13					
equally likel 1,2,312 ;	y to come to r	est pointinability that	ng to one of it will point	arrow which is the number to an odd number) 5/12	is:
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4					

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is: (A) 364/365 (B) 31/365 (D) 1/133225						
(A) 304/303	(b) 31/303	(1/303	(D) 1/133223			
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (D) $4/5$						
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (D) 7						
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: 7/50 (B) 4/25 (C) 1/25 (D) 2/25						
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) $\frac{1}{7}$	b) $\frac{5}{42}$	$\int \frac{1}{21}$	d) $\frac{4}{21}$			
	27. Runs scored by batsman in 5 one day matches are 50, 70, 82, 93, and 20. The standard deviation is					
	b) 25.49		d) 25.69			
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
a) 13, 15	13, 18		d) 13, 16			
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is						
a) $^{1}/_{2}$	b) $^1\!/_3$	d 3. The value of I	d) $\frac{1}{6}$			
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?						

a) 3 c) 5 d) 7 32. Out of the following values, which one is not possible in probability? b) $\sum x P(x) = 3$ c) P(x) = -0.5a) P(x) = 1c) P(x) = 0.533.If E(x) = 2 and E(z) = 4, then E(z - x) = ?a) 2 **9**) 6 d) Insufficient data 34. The covariance of two independent random variable is \bigcirc - 1 d) Undefined b) 0 a) 1 35.If $\Sigma P(x) = k^2 - 8$ then, the value of k is? d) Insufficient data **(**) 3 a) 0 b) 1 36.If P(x) = 0.5 and x = 4, then E(x) = ?d) 2 a) 1 **9**) 0.5 c) 4 37.In a discrete probability distribution, the sum of all probabilities is always? d) Undefined b) Infinite a) 0 38.If the probability of hitting the target is 0.4, find mean and variance. a 0.4, 0.24 b) 0.6, 0.24 c) 0.4. 0.16 d) 0.6. 0.16 39. If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 **0.4, 0.16** d) 4, 1.6

a) 2 b) 4 c) 8

40. Find the mean of tossing 8 coins.

41. What is the mean and variance for standard normal distribution?

•				•		d variance and variand	
	/ariance X)				s given by 2) – (E(X)	•	d) (E(X))2
a) E(X)	₽) E(X	2)	c) E(X2	ven by) - (E(X))		d) (E(X))2
44.M a) 0	lean of	a constai b) a	nt 'a' is _	c) a/2	_ •	d) 1	
45.V	ariance	of a cons b) a		c) a/		d) 1	
46.F i	46.Find the mean and variance of X?						
	Х	0	1	2	3	4	
	f(x)	1/9	2/9	3/9	2/9	1/9	
a) 2,	4/3	b)	3, 4/3		c) 2, 2/3		d) 3, 2/3
47. Find the expectation of a random variable X?							
				_			
	х	1	2 3				
_	f(x) 1/		2/6 1/6				
a) 0.	5	9) 1.5		c) 2.5		d) 3.5	

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

b) npq

c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- \triangle) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- (a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}