

**APPENDIX A: FIR CATALOGUE OF SNRS
WITH *HERSCHEL***

APPENDIX B: PPMAP RESULTS




Figure A1. *Herschel* three colour images of the locations of all supernova remnants in our sample. Colours are red = 250 μm , green = 160 μm , and blue = 70 μm . Purple crosses indicate the source radio centre from Green (2014) and cyan crosses indicate the X-ray centre. Magenta diamonds indicate the location of 1720 MHz OH masers. The border colour indicates the associated detection level in this study, as given in Table 1, where green = level 1, orange = level 2, magenta = level 3, grey = level 4, and black = unstudied. In some panels dashed magenta circles and white arrows indicate particular FIR dust emission features associated with the SNR. X-ray contours are overlaid onto the images of G11.2–0.3 and G15.9+0.2, MIPS contours are overlaid onto G14.1–0.1, and radio contours are overlaid onto the image of G14.3+0.1.




Figure A1 – continued Radio 20 cm contours are overlaid onto the image of G20.4+0.1, G31.9+0.0, and G33.2–0.6. X-ray contours are overlaid onto the images of G21.5–0.9, G27.4+0.0, G29.7–0.3, and G33.3+0.1.




Figure A1 – continued Radio contours are overlaid onto the images of G39.2–0.3, G43.3–0.2, G336.7+0.5. X-ray contours are overlaid onto the image of G332.4–0.4




Figure A1 – continued MIPS contours are overlaid onto the image of G344.7–0.1 and X-ray contours are overlaid onto the image of G349.7+0.2.




Figure A2. First column: *Herschel* three colour images where colours are red = $250\,\mu\text{m}$, green = $160\,\mu\text{m}$, and blue = $70\,\mu\text{m}$. Overlaid contours are X-ray unless indicated. Second column: *Spitzer* IRAC four colour images. Third column: *Spitzer* MIPS $24\,\mu\text{m}$ images. Fourth column: Radio images: VLA 20 cm where $l < 48.5^\circ$, CGPS 1420 MHz in the range $52 < l < 192^\circ$, and MOST 0.843 GHz where $l > 255^\circ$. The border colour indicates the associated detection level as given in Table 1, where green = level 1, orange = level 2, magenta = level 3, grey = level 4, and black = unstudied. This is given for this study in the first column, R06 in the second column, and PG11 in the third column. In some panels dashed magenta circles and white arrows indicate particular FIR dust emission features associated with the SNR. X-ray contours are overlaid onto the *Herschel* image of G11.2–0.3.




Figure A2 – continued MIPS 24 μm contours are overlaid onto the *Herschel* image of G14.1–0.1.




Figure A2 – continued Radio contours are overlaid onto the *Herschel* image of G14.3+0.1 and X-ray contours onto the *Herschel* image of G15.9+0.2.




Figure A2 – *continued*




Figure A2 – continued Radio contours are overlaid onto the *Herschel* image of G20.4+0.1 and X-ray contours onto the *Herschel* image of G21.5–0.9.




Figure A2 – continued X-ray contours are overlaid onto the *Herschel* image of G27.4+0.0.




Figure A2 – continued X-ray contours are overlaid onto the *Herschel* image of G29.7–0.3 and radio contours onto the *Herschel* image of G31.9+0.0.




Figure A2 – continued Radio contours are overlaid onto the *Herschel* image of G33.2–0.6, X-ray contours onto the *Herschel* image of G33.6+0.1.




Figure A2 – continued Radio contours are overlaid onto the *Herschel* images of G39.2–0.3 and G43.3–0.2.




Figure A2 – continued Radio contours are overlaid onto the *Herschel* images of G296.8–0.3, G304.6+0.1, and G310.6–0.3.




Figure A2 – continued Radio contours are overlaid onto the *Herschel* image of G310.8–0.4 and X-ray contours are overlaid onto the *Herschel* image of G332.4–0.4.




Figure A2 – continued Radio contours are overlaid onto the *Herschel* images of G336.7+0.5, G337.2–0.7, G340.4+0.4, and G340.6+0.3.




Figure A2 – continued Radio contours are overlaid onto the images of G342.0–0.2, G345.7–0.2, and G346.6–0.2. MIPS 24 μm contours are overlaid onto the *Herschel* image of G344.7–0.1.




Figure A2 – continued Radio contours are overlaid onto the images of G348.5–0.0, G348.5+0.1, and G349.2–0.1.




Figure A2 – continued X-ray contours are overlaid onto the *Herschel* image of G349.7+0.2.




Figure B1. Estimated column density with β using a Gaussian β prior with mean 1.9 and $\sigma = 0.25$. (a) G11.2-0.3: for the PWN, the shell and the ISM dust. (b) G21.5-0.9: PWN and ISM and (c) G29.7-0.3. The column density in each region is normalised by dividing by the peak column density for comparison purposes. As we have low signal to noise for these SNRs, we cannot constrain β . PPMAP is effectively returning the Gaussian posterior for these three PWNe.