

Definition

The distribution function of the random variable X is given by

$$F_X(t) = \mathbb{P}[X \leq t] = \int_{-\infty}^t f_X(x) dx.$$

1.

$$\mathbb{P}[X \geq t] = 1 - F_X(t).$$

2.

$$\mathbb{P}[t_1 \leq X \leq t_2] = F_X(t_2) - F_X(t_1).$$

12

Relation between the probability density function and the distribution

Theorem

The relation between the probability density function and the distribution function is given by

1. $F_X(t)$ can be obtained from the density function by integration:

$$F_X(t) = \int_{-\infty}^t f_X(x) dx.$$

2. $f_X(t)$ can be obtained from $F_X(t)$ by differentiation:

$$f_X(x) = F'_X(x).$$

13

The uniform distribution

The simplest continuous distribution is the uniform distribution.

Definition

A random variable X has uniform distribution over the interval $[a, b]$, if its probability density function is given by

$$f_X(t) = \begin{cases} \frac{1}{b-a} & \text{if } a \leq t \leq b \\ 0 & \text{otherwise} \end{cases}$$

When we say, choose a random number between a and b , we always talk about uniform distribution.

14

The special case of the uniform random variable

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$$

$$F_X(t) = \begin{cases} 0 & \text{if } t \leq a \\ \frac{t-a}{b-a} & \text{if } a \leq t \leq b \\ 1 & t \geq b. \end{cases}$$

15

Example

Suppose X has uniform distribution over the interval $[1, 3]$, Find the probability of the following events: (a) $1 \leq X \leq 2$. (b) $X \geq 2$. (c) $1 \leq X \leq 4$.

The distribution function is given by

$$F_X(t) = \begin{cases} 0 & \text{if } t \leq 1 \\ \frac{t-1}{2} & \text{if } 1 \leq t \leq 3 \\ 1 & t \geq 3. \end{cases}$$

Hence we have

$$\mathbb{P}[1 \leq X \leq 2] = \frac{1}{2}.$$

$$\mathbb{P}[X \geq 2] = 1 - F_X(2) = 1 - \frac{1}{2} = \frac{1}{2}.$$

$$\mathbb{P}[1 \leq X \leq 4] = F_X(4) - F_X(1) = 1 - 0 = 1.$$

16

Example

Suppose that the distribution of a random variable X is given by

$$F_X(t) = \begin{cases} 1 - e^{-t^2} & \text{if } t \geq 0 \\ 0 & \text{if } t < 0 \end{cases}$$

Find the probability density function of X .

$$f_X(t) = F'_X(t) = \begin{cases} 2te^{-t^2} & \text{if } t \geq 0 \\ 0 & \text{if } t < 0 \end{cases}$$

17

Transformation of random variables

Example

Suppose X has uniform distribution over the interval $[0, 2]$ and set $Y = X^3$. What is the distribution function and the probability density function of Y ?

First we compute $F_Y(t)$.

$$F_Y(t) = \mathbb{P}[Y \leq t] = \mathbb{P}[X^3 \leq t] = \mathbb{P}[X \leq t^{1/3}].$$

$$F_Y(t) = \begin{cases} 1 & \text{if } t \leq 0 \\ \frac{1}{2}t^{1/3} & \text{if } 0 \leq t \leq 8 \\ 1 & \text{if } t \geq 8. \end{cases}$$

$$f_Y(t) = \begin{cases} \frac{1}{6}t^{-2/3} & \text{if } 0 < t < 8 \\ 0 & \text{otherwise} \end{cases}$$

18

Transformation of random variables

Suppose X has uniform distribution over the interval $[-1, 1]$ and set $Y = X^2$. What is the distribution function and the probability density function of Y ?

Since X takes values in $[-1, 1]$, we know that $Y = X^2$ takes values in $[0, 1]$.

For $0 < t < 1$ we have

$$F_Y(t) = \mathbb{P}[Y \leq t] = \mathbb{P}[X^2 \leq t] = \mathbb{P}[-\sqrt{t} \leq X \leq \sqrt{t}] = \int_{-\sqrt{t}}^{\sqrt{t}} \frac{1}{2} dx = \sqrt{t}.$$

Hence we have

$$f_Y(t) = \begin{cases} \frac{1}{2}t^{-1/2} & \text{if } 0 < t < 1 \\ 0 & \text{if } t < 0 \text{ or } t > 1. \end{cases}$$

19

Other prominent continuous random variables

Definition

A random variable X is said to have exponential distribution with parameter λ when its density function is given by

$$f_X(t) = \begin{cases} \lambda e^{-\lambda x} & \text{if } x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

The distribution function is thus given by for values of $t > 0$:

$$F_X(t) = \int_0^t \lambda e^{-\lambda x} dx = 1 - e^{-\lambda t}.$$

In particular, we have

$$\mathbb{P}[X > t] = 1 - F_X(t) = e^{-\lambda t}.$$

20

Exponential random variables do not capture memory

$$\mathbb{P}[X > s + t | X > s] = \frac{e^{-\lambda(s+t)}}{e^{-\lambda s}} = e^{-\lambda t} = \mathbb{P}[X > t].$$

Situations where exponential distributions may arise:

- Time between two car accidents.
- The number of words between two typos in a book.
- Lifetime of items that do not age.

21

Normal random variables

Normal or Gaussian random variables are some of the most important examples of continuous random variables. They arise naturally in *the central limit theorem*.

Definition


A continuous random variable X is said to have Gaussian or normal distribution with parameters (μ, σ^2) , written $X \sim N(\mu, \sigma^2)$, if the probability density function of X is given by

$$f_X(t) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}.$$

A random variable with normal distribution with parameters $\mu = 0$ and $\sigma = 1$ is called a *standard normal distribution* or a standard Gaussian.

22

Probability density functions of normal random variables



23