

# Understanding Analysis, 2nd Edition: Stephen Abbott

## Chapter 3 - Section 3.2 - Open and Closed Sets

1. (a) Where in the proof of Theorem 3.2.3 part (ii) does the assumption that the collection of open sets be *finite* get used?
- (b) Give an example of a countable collection of open sets  $\{O_1, O_2, O_3, \dots\}$  whose intersection  $\cap_{n=1}^{\infty} O_n$  is closed, not empty, and not all of  $R$ .

### Solution:

- (a) The part where we want to take the minimum  $\epsilon$ -neighborhood  $\epsilon = \min\{\epsilon_1, \epsilon_2, \dots, \epsilon_n\} > 0$ . A minimum does not necessarily exist in an infinite set.

To be even more precise though, what we're lacking is a positive lower bound. Had there been a positive infimum (but not minimum), we could use that as  $\epsilon$  (note that the proof is still valid even if we replace minimum with infimum. It's just that taking minimum is simpler for a finite set). However, the infimum may be zero, as can be seen in the next example.

- (b) Let  $O_n = (-\frac{1}{n}, \frac{1}{n})$ . Both endpoints converge to 0, so  $\cap_{n=1}^{\infty} O_n = \{0\}$  which is closed, not empty, and not all of  $R$ , as required.

2. Let  $A = \{(-1)^n + \frac{2}{n} : n = 1, 2, 3, \dots\}$  and  $B = \{x \in Q : 0 < x < 1\}$ . Answer the following questions for each set:
  - (a) What are the limit points?
  - (b) Is the set open? Closed?
  - (c) Does the set contain any isolated points?
  - (d) Find the closure of the set.