

# Understanding Analysis, 2nd Edition

## Chapter 5 - Section 5.2 - Derivatives and the Intermediate Value Property

1. Supply proofs for part (i) and (ii) of Theorem 5.2.4.

**Solution:** Part (i):  $(f + g)'(c) = f'(c) + g'(c)$

Proof:

$$\begin{aligned}(f + g)'(c) &= \lim_{x \rightarrow c} \frac{f(x) + g(x) - f(c) - g(c)}{x - c} = \lim_{x \rightarrow c} \left( \frac{f(x) - f(c)}{x - c} + \frac{g(x) - g(c)}{x - c} \right) \\ &= f'(c) + g'(c)\end{aligned}$$

where the last step is justified because we know both limits exist, so we can apply the Algebraic Limit Theorem (Theorem 4.2.4).

Part (ii):  $(kf)'(c) = kf'(c)$  for all  $k \in \mathbf{R}$ .

Proof:

$$(kf)'(c) = \lim_{x \rightarrow c} k \left( \frac{f(x) - f(c)}{x - c} \right) = k \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = kf'(c)$$

2. Exactly one of the following requests is impossible. Decide which it is, and provide examples for the other three. In each case, assume the functions are defined on all of  $\mathbf{R}$ .

- (a) Functions  $f$  and  $g$  not differentiable at zero but where  $fg$  is differentiable at zero.
- (b) A function  $f$  not differentiable at zero and a function  $g$  differentiable at zero where  $fg$  is differentiable at zero.
- (c) A function  $f$  not differentiable at zero and a function  $g$  differentiable at zero where  $f + g$  is differentiable at zero.
- (d) A function  $f$  differentiable at zero but not differentiable at any other point.

**Solution:**

- (a)  $f(x) = 0$  for  $x \geq 0$  and  $f(x) = 1$  otherwise.  $g(x) = 1$  for  $x \geq 0$  and  $g(x) = 0$  otherwise. Both functions are discontinuous at zero hence not differentiable there, but  $fg$  is the zero function.

- (b)  $f(x) = 0$  for  $x \geq 0$  and  $f(x) = 1$  otherwise.  $g(x) = 0$  everywhere.
- (c) Impossible. If  $f + g$  were differentiable at zero, then  $(f + g) - g = f$  must be differentiable at zero as well by Theorem 5.2.4.
- (d)  $f(x) = x^2$  for  $x \in \mathbf{Q}$  and  $f(x) = 0$  otherwise.

It's differentiable at zero because for  $c = 0$ ,  $\frac{f(x)-f(c)}{x-c} = x$  and  $\lim_{x \rightarrow 0} x = 0$ .

It's not differentiable at  $c \neq 0$ . Consider a sequence  $(x_n) \rightarrow c$ . We can split it into two subsequences  $(a_n) \in \mathbf{Q}$  and  $(b_n) \notin \mathbf{Q}$ . We have  $(a_n) \rightarrow c^2$  and  $(b_n) \rightarrow 0$ . Since  $c^2 \neq 0$  for  $c \neq 0$ , the two subsequences converge to different limits, hence  $(x_n)$  doesn't converge.

3. (a) Use Definition 5.2.1 to produce the proper formula for the derivation of  $f(x) = 1/x$ .
- (b) Combine the result in part (a) with the Chain Rule to supply a proof for part (iv) of Theorem 5.2.4.
- (c) Supply a direct proof of Theorem 5.2.4 (iv) by algebraically manipulating the difference quotient for  $(f/g)$  in a style similar to the proof of Theorem 5.2.4 (iii).

**Solution:**

(a)

$$\frac{f(x) - f(c)}{x - c} = \frac{\frac{1}{x} - \frac{1}{c}}{x - c} = \frac{-(x - c)}{xc} \cdot \frac{1}{x - c} = -\frac{1}{xc}$$

So we have  $f'(c) = \lim_{x \rightarrow c} (-\frac{1}{xc}) = -\frac{1}{c^2}$  for  $c \neq 0$ .

(b) We first derive a formula for the derivative of  $h(x) = 1/g(x)$ .

$$\frac{h(x) - h(c)}{x - c} = \frac{-(g(x) - g(c))}{g(x)g(c)} \cdot \frac{1}{x - c} = -\frac{1}{g(x)g(c)} \cdot \frac{g(x) - g(c)}{x - c}$$

Taking the limit to  $c$  and noting that the second term of the product is the definition of  $g'(c)$

$$h'(c) = \lim_{x \rightarrow c} -\frac{1}{g(x)g(c)} \cdot \frac{g(x) - g(c)}{x - c} = -\frac{g'(c)}{g(c)^2}$$

Now apply the product rule to  $(fh)'(c)$  to get the formula for  $(f/g)'(c)$ .

— Simpler solution by chatgpt —

Derive the formula of  $h'(x)$  by defining  $h(x) = 1/x$  then apply the chain rule to  $h(g(x))$ .

(c)

$$\begin{aligned}\frac{\frac{f}{g}(x) - \frac{f}{g}(c)}{x - c} &= \frac{\frac{f}{g}(x) - \frac{f(x)}{g(c)} + \frac{f(x)}{g(c)} - \frac{f}{g}(c)}{x - c} \\&= \frac{f(x)}{x - c} \left( \frac{g(c) - g(x)}{g(x)g(c)} \right) + \frac{1}{g(c)(x - c)} (f(x) - f(c)) \\&= -\frac{f(x)}{g(x)g(c)} \left( \frac{g(x) - g(c)}{x - c} \right) + \frac{1}{g(c)} \left( \frac{f(x) - f(c)}{x - c} \right)\end{aligned}$$

Take the limit to  $c$  to get

$$-\frac{f(c)g'(c)}{g(c)^2} + \frac{f'(c)}{g(c)} = \frac{f'(c)g(c) - f(c)g'(c)}{g(c)^2}$$

4. Follow these steps to provide a slightly modified proof of the Chain Rule.

- (a) Show that a function  $h : A \rightarrow \mathbf{R}$  is differentiable at  $a \in A$  if and only if there exists a function  $l : A \rightarrow \mathbf{R}$  which is continuous at  $a$  and satisfies

$$h(x) - h(a) = l(x)(x - a) \quad \text{for all } x \in A.$$

- (b) Use this criterion for differentiability (in both directions) to prove Theorem 5.2.5.

**Solution:**

- (a) First we prove that if such a function  $l$  exists then  $h$  is differentiable at  $a$ . We have

$$h'(a) = \lim_{x \rightarrow a} \frac{h(x) - h(a)}{x - a} = \lim_{x \rightarrow a} l(x) = l(a)$$

Next, we prove that if  $h$  is differentiable at  $a$  then such a function  $l$  exists. Define

$$l(x) = \begin{cases} \frac{h(x) - h(a)}{x - a} & x \neq a \\ h'(a) & (x = a) \end{cases}$$

$l$  is continuous at  $a$  since  $\lim_{x \rightarrow a} l(x) = \lim_{x \rightarrow a} \frac{h(x) - h(a)}{x - a} = h'(a) = l(a)$ .

$h(x) - h(a) = l(x)(x - a)$  for all  $x \in A$  by definition.

- (b) TODO.

5. Let  $l(x) = \begin{cases} x^a & x > 0 \\ 0 & x \leq 0 \end{cases}$ .

- (a) For which values of  $a$  is  $f$  continuous at zero?
- (b) For which values of  $a$  is  $f$  differentiable at zero? In this case, is the derivative function continuous?
- (c) For which values of  $a$  is  $f$  twice-differentiable?

**Solution:**

(a)  $a > 0$ . The limit at 0 from the left is 0. If  $a < 0$  then the right limit diverges. If  $a = 0$  then the right limit is 1. But if  $a > 0$  then the right limit is 0.

(b) From the direction of positive  $x$ , we have  $\lim_{x \rightarrow 0^+} \frac{x^a - 0^a}{x - 0} = \lim_{x \rightarrow 0^+} x^{a-1} = 0^{a-1}$ . This is only defined for  $a > 1$ , in which case it's 0 and agrees with the left limit, so the derivative is defined.

The derivative is continuous, but seems to require us to use the fact that  $l'(x) = ax^{a-1}$  for *non-integer*  $a$ , which we have not proved yet.

(c) The next right-side derivative at zero is  $\lim_{x \rightarrow 0^+} x^{a-2} = 0^{a-2}$ , so we need  $a > 2$ .

6. Let  $g$  be defined on an interval  $A$  and let  $c \in A$ .

- (a) Explain why  $g'(c)$  in Definition 5.2.1 could have been given by

$$g'(c) = \lim_{h \rightarrow 0} \frac{g(c+h) - g(c)}{h}$$

- (b) Assume  $A$  is open. If  $g$  is differentiable at  $c \in A$ , show

$$g'(c) = \lim_{h \rightarrow 0} \frac{g(c+h) - g(c-h)}{2h}$$

**Solution:**

- (a) Let  $h = x - c$ . Then we have

$$g'(c) = \lim_{(x-c) \rightarrow 0} \frac{g(x) - g(c)}{x - c}$$

And use the fact that  $\lim_{(x-c) \rightarrow 0} = \lim_{x \rightarrow c}$ .

(b)

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{g(c+h) - g(c-h)}{2h} &= \lim_{h \rightarrow 0} \frac{g(c+h) - g(c) + g(c) - g(c-h)}{2h} \\ &= \frac{1}{2} \lim_{h \rightarrow 0} \left( \frac{g(c+h) - g(c)}{h} + \frac{g(c) - g(c-h)}{h} \right) \\ &= \frac{1}{2} (g'(c) + g'(c)) = g'(c)\end{aligned}$$

We used the fact that  $g'(c) = \lim_{h \rightarrow 0} \frac{g(c) - g(c-h)}{h}$ , which we'll now prove. Let  $h = c - x$ . Then we have

$$\lim_{(c-x) \rightarrow 0} \frac{g(c) - g(x)}{c - x} = \lim_{(c-x) \rightarrow 0} \frac{g(x) - g(c)}{x - c}$$

And use the fact that  $\lim_{(c-x) \rightarrow 0} = \lim_{(x-c) \rightarrow 0} = \lim_{x \rightarrow c}$  (this is easy to prove with the usual  $\epsilon - \delta$ ).

One more thing to note is why the problem assumes  $A$  is open. This is so that both  $g(c-h)$  and  $g(c+h)$  are defined.

— **Slight improvement by chatgpt** —

To prove  $g'(c) = \lim_{h \rightarrow 0} \frac{g(c) - g(c-h)}{h}$ , set  $k = -h$ . Then as  $h \rightarrow 0$ , also  $k \rightarrow 0$ , and

$$\frac{g(c) - g(c-h)}{h} = \frac{g(c+k) - g(c)}{k}$$

Then reuse part (a).

7. TODO

8. TODO

9. True / false with proof / counterexample.

- (a) If  $f'$  exists on an interval and is not constant, then  $f'$  must take on some irrational values.
- (b) If  $f'$  exists on an open interval and there is some point  $c$  where  $f'(c) > 0$ , then there exists a  $\delta$ -neighborhood  $V_\delta(c)$  around  $c$  in which  $f'(x) > 0$  for all  $x \in V_\delta(c)$ .
- (c) If  $f$  is differentiable on an interval containing zero and if  $\lim_{x \rightarrow 0} f'(x) = L$ , then it must be that  $L = f'(0)$ .

**Solution:**

(a) True, by Darboux Theorem and the fact that the irrationals are dense in  $\mathbf{R}$ .

(b) — **Incorrect** —

True. Otherwise  $f'(c)$  is an isolated point and that violates Darboux Theorem.

— **Corrected** —

The above is wrong because Darboux Theorem only guarantees Intermediate Value Property and not continuity (see Section 4.5).

I would not have been able to come up with the counterexample myself. It's

$$f(x) = \begin{cases} x/2 + x^2 \sin(1/x) & x \neq 0 \\ 0 & x = 0 \end{cases}$$

The  $x/2$  term makes  $f'(0) = \frac{1}{2}$  (compare this with  $f(x) = x^2 + \sin(1/x)$  where  $f'(0)$  would be 0).

For  $x \neq 0$ ,  $f'(x) = \frac{1}{2} - \cos(1/x) + 2x \sin(1/x)$ . For any  $V_\delta(0)$  there's always an  $x$  such that  $\cos(1/x) = 1$ , which would make  $f'(x)$  negative.

(c) — **Incorrect** —

We need the following lemma: if  $\lim_{a \rightarrow b} f(a)$  is defined for some  $b$  near  $c$ , and  $\lim_{a \rightarrow c} f(a)$  is also defined, then  $\lim_{b \rightarrow c} (\lim_{a \rightarrow b} f(a)) = \lim_{a \rightarrow c} f(a)$ .

The proof is straightforward with  $\epsilon - \delta$ . There's a  $\delta$  such that  $|a - c| < \delta \implies |f(a) - L| < \epsilon$ . Make  $|b - c| < \delta/2$  and  $|a - b| < \delta/2$ .

It's very important to note that  $\lim_{a \rightarrow b} f(a)$  \*must\* be defined for  $b$  near  $c$  for this lemma to work. Otherwise, one counterexample is the Dirichlet function.  $\lim_{a \rightarrow 0} f(a) = 0$ . But the limit is undefined for all  $b \neq 0$  so the expression  $\lim_{b \rightarrow 0} (\lim_{a \rightarrow b} f(a))$  is undefined.

Now back to the original problem.  $f$  is differentiable on an interval containing zero, so  $f'(0)$  exists and also

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$$

exists for  $c$  near 0.

We also have

$$\lim_{c \rightarrow 0} \left( \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \right) = L$$

Now apply the lemma to conclude that

$$\lim_{c \rightarrow 0} \left( \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \right) = \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = f'(0) = L$$

— **Corrected** —

The nested limit lemma above only applies to functions with a single parameter. For a counterexample, consider  $\lim_{c \rightarrow 0} (\lim_{x \rightarrow c} \frac{x}{c}) = \lim_{c \rightarrow 0} 1 = 1$ . But  $\lim_{x \rightarrow 0} \frac{x}{c}$  is undefined.

Instead we can use Darboux Theorem. Suppose for contradiction that  $f'(0) \neq L$  and let  $d = |f'(0) - L|$ . By convergence to  $L$ , there exists  $\delta$  such that  $x \in V_\delta(0) \implies |f'(x) - L| < d/2$ .

But by Darboux, for such  $x \in V_\delta(0)$  then the derivative must take every value between  $f'(x)$  and  $f'(0)$ . In particular, there exists  $y \in V_\delta(0)$  such that  $|f'(y) - f'(0)| < d/2$ , contradicting convergence to  $L$  ( $f'(y)$  cannot be within  $d/2$  of both  $L$  and  $f'(0)$ ).