

Understanding Analysis, 2nd Edition

Chapter 6 - Section 6.2 - Uniform Convergence of a Sequence of Functions

1. Let

$$f_n(x) = \frac{nx}{1 + nx^2}$$

- (a) Find the pointwise limit of (f_n) for all $x \in (0, \infty)$.
- (b) Is the convergence uniform on $(0, \infty)$?
- (c) Is the convergence uniform on $(0, 1)$?
- (d) Is the convergence uniform on $(1, \infty)$?

Solution:

- (a) Intuitively, as n grows the $+1$ term starts to not matter, so it should converge to $\frac{nx}{nx^2} = \frac{1}{x}$. Formally

$$\lim_{n \rightarrow \infty} \frac{nx}{1 + nx^2} = \lim_{n \rightarrow \infty} \frac{x}{1/n + x^2} = 1/x$$

- (b) No. Given $\epsilon > 0$, let N such that for all $n \geq N$, $|f_n(x) - \frac{1}{x}| < \epsilon$. Or in other words

$$\left| \frac{nx}{1 + nx^2} - \frac{1}{x} \right| = \frac{1}{x(1 + nx^2)} < \epsilon$$

For this to be uniformly convergent, the N needs to work for all x . But the lhs is unbounded as $x \rightarrow 0$; there's always a small enough x such that the lhs is eventually greater than ϵ .

- (c) No, for the same reason as part (b). We can always choose small enough x to break the uniform convergence.
- (d) Now the answer is yes. Reusing the result from part (b), for $x > 1$, the value is upper-bounded by $x = 1$

$$\frac{1}{x(1 + nx^2)} < \frac{1}{1 + n}$$

This doesn't depend on x , so we can simply choose $N \geq \frac{1}{\epsilon} - 1$.

2. (a) Define a sequence of functions on $\mathbf{R}$ by

$$f_n(x) = \begin{cases} 1 & \text{if } x = 1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$$

and let f be the pointwise limit of f_n .

Is each f_n continuous at zero? Does $f_n \rightarrow f$ uniformly on $\mathbf{R}$? Is f continuous at zero?

- (b) Repeat this exercise using the sequence of functions

$$g_n(x) = \begin{cases} x & \text{if } x = 1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$$

- (c) Repeat once more with

$$h_n(x) = \begin{cases} 1 & \text{if } x = \frac{1}{n} \\ x & \text{if } x = 1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n-1} \\ 0 & \text{otherwise} \end{cases}$$

In each case, explain how the results are consistent with the Continuous Limit Theorem (Theorem 6.2.6).

Solution:

- (a) Yes, each f_n is continuous at zero. Simply choose $\delta < \frac{1}{n}$.

To answer uniform convergence, let's first define f . For x not of the form $\frac{1}{n}$, $f(x)$ is clearly 0. For x of the form $\frac{1}{n}$, then for all $m \geq n$, $f_m(x) = 1$ i.e. eventually all $f_n(x)$ becomes 1. So f is

$$f(x) = \begin{cases} 1 & \text{if } x = \frac{1}{n} \text{ for } n \in \mathbf{N} \\ 0 & \text{otherwise} \end{cases}$$

It's not uniformly convergent. Let $\epsilon < 1$. Now say we choose $N = 5$, then for $x = \frac{1}{6}$, we have $f_5(x) = 0$ but $f(x) = 1$. In general, no single N can work for all x because we can choose $x = \frac{1}{N+1}$, then $|f_N(x) - f(x)| = 1 > \epsilon$.

Finally, f is not continuous at zero. We have $f(0) = 0$. But since $\frac{1}{n} \rightarrow 0$, there's always x of the form $\frac{1}{n}$ close enough to zero such that $f(x) = 1$.