

CS 577 - Discrete Primer

Marc Renault

Department of Computer Sciences
University of Wisconsin – Madison

Summer 2023

TopHat Section 001 Join Code: 275653



DISCRETE MATHEMATICS

Definition

Rigorous mathematical study of discrete structures.

DISCRETE MATHEMATICS

Definition

Rigorous mathematical study of discrete structures.

Key Discrete Concepts for CS 577

Core

- Logic
- Sets
- Recurrences
- Relations and Function
- Graphs and Trees
- Counting

DISCRETE MATHEMATICS

Definition

Rigorous mathematical study of discrete structures.

Key Discrete Concepts for CS 577

Core

- Logic
- Sets
- Recurrences
- Relations and Function
- Graphs and Trees
- Counting

Applied in CS 577

- Proofs esp. **Induction**
- Invariants
- Program Correctness

LOGIC

PROPOSITIONS

Definition

A statement that is either true or false.

PROPOSITIONS

Definition

A statement that is either true or false.

Example

True proposition:

- Empire is the best Star Wars movie

PROPOSITIONS

Definition

A statement that is either true or false.

Example

True proposition:

- ~~Empire is the best Star Wars movie~~
- Ottawa is the capital of Canada.

PROPOSITIONS

Definition

A statement that is either true or false.

Example

True proposition:

- ~~Empire is the best Star Wars movie~~
- Ottawa is the capital of Canada.

False proposition:

- Toronto is the capital of Canada.

PROPOSITIONS

Definition

A statement that is either true or false.

Example

True proposition:

- ~~Empire is the best Star Wars movie~~
- Ottawa is the capital of Canada.

False proposition:

- Toronto is the capital of Canada.

Operations

- And: \wedge , $\&$, $\&\&$
- Or: \vee , $|$, $||$
- Negation: \neg , $!$

- Implies: \implies

- If and only if (iff): \iff

$$P \iff Q \equiv P \implies Q \wedge Q \implies P$$

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F				
F	T				
T	F				
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F			
F	T				
T	F				
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F		
F	T				
T	F				
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	
F	T				
T	F				
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T				
T	F				
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F			
T	F				
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T		
T	F				
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	
T	F				
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F				
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F			
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F	T		
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F	T	F	
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F	T	F	F
T	T				

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F	T	F	F
T	T	T			

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F	T	F	F
T	T	T	T		

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F	T	F	F
T	T	T	T	T	F

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F	T	F	F
T	T	T	T	T	F

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F	T	F	F
T	T	T	T	T	F

Logical Equivalence

TopHat 1: Is $P \implies Q$ equivalent to $\neg P \implies \neg Q$?

TRUTH TABLES

a	b	$a \wedge b$	$a \vee b$	$a \implies b$	$\neg a$
F	F	F	F	T	T
F	T	F	T	T	T
T	F	F	T	F	F
T	T	T	T	T	F

Logical Equivalence

TopHat 1: Is $P \implies Q$ equivalent to $\neg P \implies \neg Q$?

Exercise: Prove it!

PREDICATES

Definition

For an underlying domain D . A predicate is a mapping of D to propositions.

PREDICATES

Definition

For an underlying domain D . A predicate is a mapping of D to propositions.

Quantifiers

- For all: $\forall. \forall x \in \mathbb{Z}, \text{Even}(x) \iff \text{Odd}(x + 1)$
- There exists: $\exists. \exists \text{person} \in \text{This Room}, \text{LovesStarWars}(x)$
- Order matters when combining quantifiers!

PREDICATES

Definition

For an underlying domain D . A predicate is a mapping of D to propositions.

Quantifiers

- For all: $\forall. \forall x \in \mathbb{Z}, \text{Even}(x) \iff \text{Odd}(x + 1)$
- There exists: $\exists. \exists \text{person} \in \text{This Room}, \text{LovesStarWars}(x)$
- Order matters when combining quantifiers!

Logical Equivalence

TopHat 2: What is the logical equivalence of $\neg(\forall x S(x))$?

SETS

SETS

Definition

- A well-defined collection of elements from some domain. Each element in a set is unique.
- A *multiset* may contain duplicates.

SETS

Definition

- A well-defined collection of elements from some domain. Each element in a set is unique.
- A *multiset* may contain duplicates.

Ex: $A = \{11, 12, 13\}$

$B = \{x \in \mathbb{Z} \mid 10 < x \leq 13\}$

SETS

Definition

- A well-defined collection of elements from some domain. Each element in a set is unique.
- A *multiset* may contain duplicates.

Ex: $A = \{11, 12, 13\}$

$B = \{x \in \mathbb{Z} \mid 10 < x \leq 13\}$

Basic Notations


$A \subset B$ A is a proper subset of B , meaning that A contains some (or none) of the elements of B but not all.

$A \subseteq B$ A is subset of B and A may contain all of the elements of B .


$|A|$ The cardinality of A is the number of elements in the set.

SET OPERATIONS


Union: $A \cup B$



Intersection: $A \cap B$




Set difference: $A \setminus B$ or $A - B$




SET OPERATIONS


Union: $A \cup B$



Intersection: $A \cap B$



Set difference: $A \setminus B$ or $A - B$



Other Notions

\emptyset or $\{\}$ The null or empty set.

$\mathcal{P}(A)$ Power set of A . A set of all possible subsets of A (including \emptyset).

TOPHATS

TopHat 3

What is $\{a, b, c\} \setminus \{c, d, e\}$?

TOPHATS

TopHat 3

What is $\{a, b, c\} \setminus \{c, d, e\}$?

TopHat 4

What is the size of $\mathcal{P}(A)$ for some set A ?

RELATIONS AND FUNCTIONS

RELATIONS

Cartesian Product

For two set A and B , $A \times B = \{(a, b) \mid a \in A \wedge b \in B\}$.

RELATIONS

Cartesian Product

For two set A and B , $A \times B = \{(a, b) \mid a \in A \wedge b \in B\}$.

Definition

A relation between sets A and B is a defined subset of $A \times B$.

Ex: $R = \{(x, y) \mid x \neq y\}$

RELATIONS

Definition

A relation between sets A and B is a defined subset of $A \times B$.

Ex: $R = \{(x, y) \mid x \neq y\}$

Properties of Relations

Reflexive If $\forall a \in A, R(a, a)$. (*antireflexive*: $\forall a \in A, \neg R(a, a)$)

Symmetric If $\forall a, b \in A, R(a, b) \iff R(b, a)$. (*antisymmetric*:
 $\forall a, b \in A, R(a, b) \cap R(b, a) \implies a = b$)

Transitive If $\forall a, b, c \in A, R(a, b) \cap R(b, c) \implies R(a, c)$.

RELATIONS

Definition

A relation between sets A and B is a defined subset of $A \times B$.

Ex: $R = \{(x, y) \mid x \neq y\}$

Properties of Relations

Reflexive If $\forall a \in A, R(a, a)$. (*antireflexive*: $\forall a \in A, \neg R(a, a)$)

Symmetric If $\forall a, b \in A, R(a, b) \iff R(b, a)$. (*antisymmetric*:
 $\forall a, b \in A, R(a, b) \cap R(b, a) \implies a = b$)

Transitive If $\forall a, b, c \in A, R(a, b) \cap R(b, c) \implies R(a, c)$.

Types of Relations

- Equivalence Relations: reflexive, symmetric, and transitive.
- Order Relations: antisymmetric and transitive.
- Functions

FUNCTIONS

Definition

$f : A \rightarrow B$ is a function from A to B . That is for every $a \in A$ there is at most one $b \in B$.

Ex. $f(x) = y + 1$ for $x, y \in \mathbb{R}$.

FUNCTIONS

Definition


$f : A \rightarrow B$ is a function from A to B . That is for every $a \in A$ there is at most one $b \in B$.

Ex. $f(x) = y + 1$ for $x, y \in \mathbb{R}$.


Terminology

- **Domain:** The values of A .
- **Range / Codomain:** The values of B


FUNCTIONS




An injective non-surjective function
(injection, not a **bijection**)



An injective surjective function
(**bijection**)



A non-injective surjective function
(**surjection**, not a **bijection**)



A non-injective non-surjective
function (also not a **bijection**)

Types of Functions

- one-to-one / injective
- onto / surjective
- bijection (both onto and one-to-one)

INDUCTION

PROOF BY INDUCTION

What is induction?

- The most important proof technique in discrete math and CS.
- It proves that $P(n)$ holds for every natural number n , i.e., $n = 0, 1, 2, 3, \dots$

PROOF BY INDUCTION

What is induction?

- The most important proof technique in discrete math and CS.
- It proves that $P(n)$ holds for every natural number n , i.e., $n = 0, 1, 2, 3, \dots$

Induction Formula

Step 1 State the induction hypothesis.

Step 2 Show that the induction hypothesis holds for the base case(s).

Step 3 Assume hypothesis is true for k , show that it holds for $k + 1$.

PROOF BY INDUCTION

Induction Formula

- Step 1** State the induction hypothesis.
- Step 2** Show that the induction hypothesis holds for the base case(s).
- Step 3** Assume hypothesis is true for k , show that it holds for $k + 1$.

Special Types of Induction

- Strong induction: we assume true for 1 to k instead of just k .
- Structural induction: we are reasoning about a structure that we map to the natural numbers.

PROOF BY INDUCTION

Induction Formula

Step 1 State the induction hypothesis.

Step 2 Show that the induction hypothesis holds for the base case(s).

Step 3 Assume hypothesis is true for k , show that it holds for $k + 1$.

Induction Exercises

- Show $\sum_1^n 2^n = 2^{n+1} - 2$.

PROOF BY INDUCTION

Induction Formula

Step 1 State the induction hypothesis.

Step 2 Show that the induction hypothesis holds for the base case(s).

Step 3 Assume hypothesis is true for k , show that it holds for $k + 1$.

Induction Exercises

- Show $\sum_1^n 2^n = 2^{n+1} - 2$.
- Show, for $n \geq 5$, $4n < 2^n$.

PROOFS

PROOFS

Definition

A proof of a proposition P is a chain of logical deductions ending in P and starting from some set of axioms.

PROOFS

Definition

A proof of a proposition P is a chain of logical deductions ending in P and starting from some set of axioms.

Types (other than induction)

Proof by Picture Actually not valid!

PROOFS

Definition

A proof of a proposition P is a chain of logical deductions ending in P and starting from some set of axioms.

Types (other than induction)

Direct Proof Series of implications.

PROOFS

Definition

A proof of a proposition P is a chain of logical deductions ending in P and starting from some set of axioms.

Types (other than induction)

Direct Proof Series of implications.

Indirect Proof (Contrapositive) To show $P \implies Q$, we show that $\neg Q \implies \neg P$.

PROOFS

Definition

A proof of a proposition P is a chain of logical deductions ending in P and starting from some set of axioms.

Types (other than induction)

Direct Proof Series of implications.

Indirect Proof (Contrapositive) To show $P \implies Q$, we show that $\neg Q \implies \neg P$.

If and only if $P \iff Q$ means proving $P \implies Q$ and $Q \implies P$.

PROOFS

Definition

A proof of a proposition P is a chain of logical deductions ending in P and starting from some set of axioms.

Types (other than induction)

Direct Proof Series of implications.

Indirect Proof (Contrapositive) To show $P \implies Q$, we show that $\neg Q \implies \neg P$.

If and only if $P \iff Q$ means proving $P \implies Q$ and $Q \implies P$.

Proof by Contradiction Assume claim is not true, and derive a contradiction.

PROOFS

Definition

A proof of a proposition P is a chain of logical deductions ending in P and starting from some set of axioms.

Types (other than induction)

Direct Proof Series of implications.

Indirect Proof (Contrapositive) To show $P \implies Q$, we show that $\neg Q \implies \neg P$.

If and only if $P \iff Q$ means proving $P \implies Q$ and $Q \implies P$.

Proof by Contradiction Assume claim is not true, and derive a contradiction.

Proof by Cases (Brute Force / Exhaustion) Split into cases and prove separately for each case.

COUNTING

COUNTING

Basic Techniques

- *k*-to-1 Rule: Is there a *k* to 1 ratio between 2 sets?
- Sum Rule: Combine disjoint sets; add cardinality.
- Product Rule: Cartesian product of sets; multiply cardinality.

COUNTING

Basic Techniques

- **k -to-1 Rule:** Is there a k to 1 ratio between 2 sets?
- **Sum Rule:** Combine disjoint sets; add cardinality.
- **Product Rule:** Cartesian product of sets; multiply cardinality.

Perms and Comb

- **k -Permutation:** $k!$
- **r -Permutation of n items:** ${}_nP_r = P(n, r) = \frac{n!}{(n-r)!}$
- **r -Combination of n items:** ${}_nC_r = C(n, r) = \frac{n!}{r!(n-r)!}$

COUNTING

Basic Techniques

- **k -to-1 Rule:** Is there a k to 1 ratio between 2 sets?
- **Sum Rule:** Combine disjoint sets; add cardinality.
- **Product Rule:** Cartesian product of sets; multiply cardinality.

Perms and Comb

- **k -Permutation:** $k!$
- **r -Permutation of n items:** ${}_nP_r = P(n, r) = \frac{n!}{(n-r)!}$
- **r -Combination of n items:** ${}_nC_r = C(n, r) = \frac{n!}{r!(n-r)!}$

Pigeonhole Principal

If n pigeons are placed into m holes, and $n > m$, then at least one hole has more than one pigeon.

INVARIANTS

INVARIANTS

Definition

- A property of a system that holds after every step.

INVARIANTS

Definition

- A property of a system that holds after every step.
- Often critical to showing program correctness.

INVARIANTS

Definition

- A property of a system that holds after every step.
- Often critical to showing program correctness.

Robot Exercise

Suppose we have a robot which walks on a 2-dimensional grid. The rows and columns of the grid are labelled by integers. Our robot starts at position $(0, 0)$, and can only move diagonally, one square at a time. Can we get to $(8, 9)$? Why or why not?

PROGRAM CORRECTNESS

PROGRAM CORRECTNESS

Definition

A program/algorithm is correct if it is:

- **Partial correctness/correct/sound** (any returned value is true), and
- **Termination/complete** (returns a value for all valid inputs).

PROGRAM CORRECTNESS

Definition

A program/algorithm is correct if it is:

- **Partial correctness/correct/sound** (any returned value is true), and
- **Termination/complete** (returns a value for all valid inputs).

Proving correctness

- Requires 2 proofs (one for soundness and one for completeness).
- Often requires identifying invariants and induction.

RECURRENCES

RECURRENCES

Definition

An inductive (or recursive) definition of a sequence.

RECURRENCES

Definition

An inductive (or recursive) definition of a sequence.

Methods for Solving Recurrences

- Guess Method / Recurrence Tree
- Unwind
- Master Theorem

RECURRENCES

Definition

An inductive (or recursive) definition of a sequence.

Methods for Solving Recurrences

- Guess Method / Recurrence Tree
- Unwind
- Master Theorem

Exercises

Assume $T(1) = 1$ for all.

- $T(n) = T(n/2) + 1$
- $T(n) = T(n/2) + n$
- $T(n) = 3T(n/3) + n$

GRAPHS AND TREES

GRAPHS

Definition

A graph G is a pair $G = (V, E)$, where V is a set of vertices/nodes and E is a set of edges/arcs connecting a pair of vertices. That is, $E \in V \times V$.

GRAPHS

Definition

A graph G is a pair $G = (V, E)$, where V is a set of vertices/nodes and E is a set of edges/arcs connecting a pair of vertices. That is, $E \in V \times V$.

Some Special Graphs

- Complete graph (K_4)

GRAPHS

Definition

A graph G is a pair $G = (V, E)$, where V is a set of vertices/nodes and E is a set of edges/arcs connecting a pair of vertices. That is, $E \in V \times V$.

Some Special Graphs

- Complete graph (K_4)
- Cycle (C_4)

GRAPHS

Definition

A graph G is a pair $G = (V, E)$, where V is a set of vertices/nodes and E is a set of edges/arcs connecting a pair of vertices. That is, $E \in V \times V$.

Some Special Graphs

- Complete graph (K_4)
- Cycle (C_4)
- Path (P_4)

GRAPHS

Definition

A graph G is a pair $G = (V, E)$, where V is a set of vertices/nodes and E is a set of edges/arcs connecting a pair of vertices. That is, $E \in V \times V$.

Some Special Graphs

- Complete graph (K_4)
- Cycle (C_4)
- Path (P_4)
- Trees

GRAPHS

Definition

A graph G is a pair $G = (V, E)$, where V is a set of vertices/nodes and E is a set of edges/arcs connecting a pair of vertices. That is, $E \in V \times V$.

Some Special Graphs

- Complete graph (K_4)
- Cycle (C_4)
- Path (P_4)
- Trees
- Digraph

GRAPHS

Definition

A graph G is a pair $G = (V, E)$, where V is a set of vertices/nodes and E is a set of edges/arcs connecting a pair of vertices. That is, $E \in V \times V$.

Some Special Graphs

- Complete graph (K_4)
- Cycle (C_4)
- Path (P_4)
- Trees
- Digraph
- Directed Acyclic Graph (DAG)

GRAPHS

Definition

A graph G is a pair $G = (V, E)$, where V is a set of vertices/nodes and E is a set of edges/arcs connecting a pair of vertices. That is, $E \in V \times V$.

Some Special Graphs

- Complete graph (K_4)
- Cycle (C_4)
- Path (P_4)
- Trees
- Digraph
- Directed Acyclic Graph (DAG)
- Bipartite

GRAPHS


Definition

A graph G is a pair $G = (V, E)$, where V is a set of vertices/nodes and E is a set of edges/arcs connecting a pair of vertices. That is, $E \in V \times V$.

Some Special Graphs

- Complete graph (K_4)
- Cycle (C_4)
- Path (P_4)
- Trees
- Digraph
- Directed Acyclic Graph (DAG)
- Bipartite
- Forests


GRAPH ENCODINGS



Representations

- **Adjacency matrix:** $|V|$ by $|V|$ matrix with a 1 if nodes are adjacent.


GRAPH ENCODINGS



Representations

- **Adjacency matrix:** $|V|$ by $|V|$ matrix with a 1 if nodes are adjacent.
- **Adjacency list:** For each node, list adjacent nodes.


GRAPH ENCODINGS



Representations

- **Adjacency matrix:** $|V|$ by $|V|$ matrix with a 1 if nodes are adjacent.
- **Adjacency list:** For each node, list adjacent nodes.
- **Edge list:** List of all node pairs representing the edges.

GRAPH ENCODINGS



Representations

- **Adjacency matrix:** $|V|$ by $|V|$ matrix with a 1 if nodes are adjacent.
- **Adjacency list:** For each node, list adjacent nodes.
- **Edge list:** List of all node pairs representing the edges.
- **Incidence matrix:** $|V|$ by $|E|$ matrix with a 1 if node is incident to the edge.

TREES

Definition

- A connected graph without cycles.
- A single node may be designated as the *root* of the tree.
- Any node with degree 1 that is not the root is a *leaf*.

TREES

Definition

- A connected graph without cycles.
- A single node may be designated as the *root* of the tree.
- Any node with degree 1 that is not the root is a *leaf*.

Properties of a tree T

- 1 If $|V| \geq 2$, (unrooted) T has at least 2 leaves.
- 2 For all nodes u and v , there exists one path between them in T .
- 3 $|V| = |E| + 1$ for $|V| \geq 1$.

TREES

Definition

- A connected graph without cycles.
- A single node may be designated as the *root* of the tree.
- Any node with degree 1 that is not the root is a *leaf*.

Properties of a tree T

- 1 If $|V| \geq 2$, (unrooted) T has at least 2 leaves.
- 2 For all nodes u and v , there exists one path between them in T .
- 3 $|V| = |E| + 1$ for $|V| \geq 1$.

TopHat 6

Is P_{10} a tree?

APPENDIX


REFERENCES

IMAGE SOURCES I



WISCONSIN
UNIVERSITY OF WISCONSIN-MADISON

<https://brand.wisc.edu/web/logos/>



A non-injective non-surjective function (neither one-to-one nor onto)

A bijective function (one-to-one and onto)

A non-injective surjective function (neither one-to-one nor onto)

A non-surjective injective function (neither one-to-one nor onto)

<https://en.wikipedia.org/wiki/Bijection>