

Python Global Model Documentation

This document describes the PyGMol global model and explicitly the equations which are being solved for inside in the model, with the intention, to give the most complete and transparent description.

Contents

Symbols	2
Model Description	3
Input parameters for the model:	4
Model outputs:	4
Particle Density Balance	4
Volumetric Reactions Contribution	4
Flow Contribution	5
Inflow	5
Pressure Regulation	6
Diffusion Contribution	6
Diffusion losses	7
Boundary sources	9
Minimal Allowed Species Density	9
Electron Energy Density Balance	10
Contribution of Elastic and Inelastic Collisions	10
Electron Generation and Loss Contribution	11
Energy Loss by Electron Transport	11
Energy Loss by Ion Transport	11
Minimal Allowed Electron Energy Density	12
References	13

Symbols

Table 1: Overview of symbols used in the description of the PyGMol global modelling framework.

Symbol	Unit	Description
i, k		Index, running over species in a kinetic scheme
i_0, i_+, i_-		Indices i running over neutral, positive and negative species respectively
j		Index, running over reactions in a kinetic scheme
N_S		number of species in a kinetic scheme
n_i	$[m^{-3}]$	Number density of the i -th species in a kinetic scheme
n_e	$[m^{-3}]$	Electron number density
ρ_e	$[eV \cdot m^{-3}]$	Electron energy density
P	$[W]$	Absorbed power
p_0	$[Pa]$	Desired pressure
p	$[Pa]$	Instantaneous pressure
Q_i	$[scm]$	Feed flow for i -th species in a kinetic scheme
R_p, Z_p	$[m]$	Plasma dimensions: radius and length
V	$[m^3]$	Plasma volume
T_g	$[K]$	Neutral temperature
T_i	$[K]$	Positive ion temperature
T_e	$[K]$	Electron temperature
T_g	$[eV]$	Neutral temperature
T_i	$[eV]$	Positive ion temperature
T_e	$[eV]$	Electron temperature
k_j	$[m^{-3+3m}s^{-1}]$	Reaction rate coefficient of the j -th reaction of an order m in a kinetic scheme
R_j	$[m^{-3}s^{-1}]$	Reaction rate of the j -th reaction in a kinetic scheme
$M_{cp,j}$	$[kg]$	Mass of the collision partner in j -th electron reaction
M_i	$[kg]$	Mass of the i -th species
m_e	$[kg]$	Electron mass
q_i	$[e]$	Charge of the i -th species
e	$[C]$	Electron charge
σ_i^{LJ}	$[m]$	σ parameter of the Lennard-Jones potential for i -th species
$\Delta E_{e,j}^{inel}$	$[eV]$	Electron energy loss due to an inelastic collision j
a_{ij}^L		stoichiometric coefficient of i -th distinct species on left hand side in j -th reaction
a_{ij}^R		stoichiometric coefficient of i -th distinct species on right hand side in j -th reaction
a_{ij}		Net stoichiometric coefficient of i -th distinct species in j -th reaction

A_j	$[m^{-3+3m}s^{-1}]$	Arrhenius parameter – pre-exponential factor
n_j		Arrhenius parameter – exponent
$E_{a,j}$	$[eV]$	Arrhenius parameter – activation energy
$E_{a,j}$	$[K]$	Arrhenius parameter – activation energy
k_B	$[JK^{-1}]$	Boltzmann constant
s_i		Sticking coefficient – probability of i -th species sticking to a plasma boundary ($s_i \in [0, 1]$)
r_{ik}		Return coefficient – number of i -th species returned for each <i>one</i> of <i>stuck</i> k -th species ($r_{ik} \in \mathbf{R}_0^+$)
D_i	$[m^2s^{-1}]$	Diffusion coefficient of i -th species
D_a	$[m^2s^{-1}]$	Ambipolar diffusion coefficient
Λ	$[m]$	Characteristic diffusion length
λ_i	$[m]$	Mean free path of i -th species
\bar{v}_i	$[ms^{-1}]$	Mean speed of i -th species
σ_{ik}^m	$[m^2]$	Momentum transfer cross section for i -th species scattering on k -th species
\bar{V}_s	$[V]$	Mean sheath voltage
n_{\min}	$[m^{-3}]$	Minimal allowed particle density
$\varrho_{e,\min}$	$[eV \cdot m^{-3}]$	Minimal allowed electron energy density

Model Description

The model solves the set of following ordinary differential equations (ODE):

- **Particle density balance equation for heavy species**, including contributions from volumetric reactions, flow and from diffusion sinks and surface sources of n_i .
- **Electron energy density balance equation**, including contributions of power absorbed by the plasma, elastic and inelastic collisions between electrons and heavy species, generation and loss of electrons in volumetric reactions and power lost to walls by electrons and ions.

The electron density n_e is not solved for explicitly but rather implicitly by enforcing the charge neutrality. Also, the heavy species temperature is not resolved in the model, but rather treated as a constant input parameter. The collisional kinetics is not described by cross sections, but rather parametrized for each reaction with the Arrhenius formula (5), (6). The model was developed mainly for the purpose of plasma chemistry reduction, which justifies it's simplicity and the degree of ap-

proximation. For those reasons, the model should only be used with a great care to obtain any sort of quantitative results.

Input parameters for the model:

- Plasma parameters: $P, p_0, Q_i, R_p, Z_p, T_g$
- Reaction set parameters: $M_i, q_i, \sigma_i^{\text{LJ}}, k_j, \Delta E_{e,j}^{\text{inel}}$

Model outputs:

- Densities of heavy species and electrons: n_i, n_e
- Electron temperature T_e

The rest of this chapter describes all the solved equations in detail.

Particle Density Balance

The time derivative of all heavy species densities is expressed as a sum of contributions from volumetric processes, flow sources and sinks and surface (diffusion) processes:

$$\frac{dn_i}{dt} = \left(\frac{\delta n_i}{\delta t} \right)_{\text{vol}} + \left(\frac{\delta n_i}{\delta t} \right)_{\text{flow}} + \left(\frac{\delta n_i}{\delta t} \right)_{\text{diff}}. \quad (1)$$

Volumetric Reactions Contribution

The contribution of volumetric reactions is

$$\left(\frac{\delta n_i}{\delta t} \right)_{\text{vol}} = \sum_j G_{ij} - \sum_j L_{ij}, \quad (2)$$

where G_{ij} and L_{ij} are contributions of generation and loss of n_i due to inelastic reaction j . In greater detail, it can be written as

$$\left(\frac{\delta n_i}{\delta t} \right)_{\text{vol}} = \sum_j (a_{ij}^{\text{R}} - a_{ij}^{\text{L}}) R_j, \quad (3)$$

$$R_j = k_j \prod_l n_{lj}^L, \quad (4)$$

where n_{lj}^L is the density of l^{th} species on the left hand side of reaction j . The reaction rate coefficient k_j in this model takes form of the Arrhenius equation

$$k_j = A_j \left(\frac{T_g}{300\text{K}} \right)^{n_j} \exp \left(-\frac{E_{a,j}}{T_g} \right) \quad (5)$$

for heavy species reactions j (reactions where all the reactants are heavy species) and

$$k_j = A_j \left(\frac{T_e}{1\text{eV}} \right)^{n_j} \exp \left(-\frac{E_{a,j}}{T_e} \right) \quad (6)$$

for electron processes j (reactions where at least one reactant is an electron). The Arrhenius parameters A_j , n_j and $E_{a,j}$ (or $E_{a,j}$) describe the collisional kinetics of the model.

Flow Contribution

The contribution of flow to the time evolution of heavy species densities will consist of inflow and outflow terms as well as a term regulating the pressure.

$$\left(\frac{\delta n_i}{\delta t} \right)_{\text{flow}} = \left(\frac{\delta n_i}{\delta t} \right)_{\text{flow}}^{\text{in}} + \left(\frac{\delta n_i}{\delta t} \right)_{\text{flow}}^{\text{out}} + \left(\frac{\delta n_i}{\delta t} \right)_{\text{flow}}^{\text{reg}} \quad (7)$$

Inflow

$$\left(\frac{\delta n_i}{\delta t} \right)_{\text{flow}}^{\text{in}} = \frac{Q'_i}{V}, \quad (8)$$

where

$$Q'_i = \frac{N_A}{V_m \cdot 60} Q_i = 4.478 \times 10^{17} \cdot Q_i$$

is the inflow expressed in [particles/sec] rather than in [sccm]. $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ is Avogadro constant and $V_m = 2.241 \times 10^4 \text{ cm}^3 \text{mol}^{-1}$ is the molar volume for ideal gas at standard temperature and pressure.

Outflow

The outflow term is set in such a way that only neutrals are leaving the plasma region due to the flow, the neutral species flow rate is proportional to the species density and the total flow rate out of the plasma region is the same as total inflow rate:

$$\left(\frac{\delta n_i}{\delta t}\right)_{\text{flow}}^{\text{out}} = \begin{cases} -\frac{\sum Q'_i}{\sum n_{i_0}} \cdot \frac{n_i}{V} & \text{neutrals,} \\ 0 & \text{ions,} \end{cases} \quad (9)$$

where the index i_0 runs only over neutral species.

Pressure Regulation

A term regulating the plasma pressure is added to the particle balance equation, accounting for changes in p due to dissociation/association processes and to diffusion losses and surface sources. This term, similarly to the outflow term, only acts upon the neutral species and can be viewed as an addition to the outflow term, or physically as adjusting a pressure-regulation valve between a plasma chamber and a pump, based on the instantaneous pressure.

$$\left(\frac{\delta n_i}{\delta t}\right)_{\text{flow}}^{\text{reg}} = \begin{cases} -\frac{p - p_0}{p_0} \frac{n_i}{\tau_p} & \text{neutrals,} \\ 0 & \text{ions.} \end{cases} \quad (10)$$

Here, p is the instantaneous pressure from the state equation for an ideal gas

$$p = k_B T_g \cdot \sum_i n_i, \quad (11)$$

and τ_p is a pressure recovery time scale. A value of $\tau_p = 10^{-3}$ s was found to yield satisfactory results for a wide range of process parameters.

Diffusion Contribution

The diffusion contribution towards the particle balance equation is ultimately controlled by vector of sticking coefficients s_i and matrix of return coefficients r_{ik}

and the diffusion model:

$$\left(\frac{\delta n_i}{\delta t}\right)_{\text{diff}} = \left(\frac{\delta n_i}{\delta t}\right)_{\text{diff}}^{\text{out}} + \left(\frac{\delta n_i}{\delta t}\right)_{\text{diff}}^{\text{in}}. \quad (12)$$

Diffusion losses

The rate of species loss to the plasma boundaries due to diffusion is expressed as

$$\left(\frac{\delta n_i}{\delta t}\right)_{\text{diff}}^{\text{out}} = \frac{A}{V} \Gamma_{\text{wall},i}, \quad (13)$$

where V and A are plasma volume and plasma boundaries area respectively, while the wall fluxes $\Gamma_{\text{wall},i}$, as used (among others) by Kushner in GlobalKin [1], is expressed as

$$\Gamma_{\text{wall},i} = -\frac{D_i n_i s_i}{s_i \Lambda + \frac{4D_i}{\bar{v}_i}} \quad (14)$$

where

$$\Lambda = \left[\left(\frac{\pi}{Z_p} \right)^2 + \left(\frac{2.405}{R_p} \right)^2 \right]^{-1/2}. \quad (15)$$

The diffusion coefficient is calculated separately for neutrals and ions. For positive and negative ions, the diffusion coefficient is the coefficient of ambipolar diffusion in electronegative plasma, as proposed by Stoeffels *et al.* in [2].

$$D_i = \begin{cases} D_i^{\text{free}} & \text{neutrals,} \\ D_+^{\text{free}} \frac{1 + \gamma(1 + 2\alpha)}{1 + \alpha\gamma} & \text{+ions,} \\ 0 & \text{-ions.} \end{cases} \quad (16)$$

Here, $\gamma = T_e/T_i$ and $\alpha = \sum n_{i-}/n_e$. $D_i = 0$ for negative ions implies that no negative ions are reaching the plasma boundaries and therefore there are no negative ion diffusion losses. This approximation is justified by the positive plasma potential trapping the negative ions in the plasma bulk. It should be noted that the stated ambipolar diffusion coefficients are only valid for the case of $\alpha \ll \mu_e/\mu_i$, where μ are mobilities of electrons and ions respectively. The free diffusion coefficient for heavy

species is calculated as

$$D_i^{\text{free}} = \frac{\pi}{8} \lambda_i \bar{v}_i. \quad (17)$$

The mean free path λ_i for all heavy species is

$$\frac{1}{\lambda_i} = \sum_k n_k \sigma_{ik}^m (1 - \delta_{ik}), \quad (18)$$

where σ_{ik}^m is the momentum transfer cross section, and the mean speed \bar{v}_i is the mean thermal speed

$$\bar{v}_i = \begin{cases} \left(\frac{8k_B T_g}{\pi M_i} \right)^{1/2} & \text{neutrals,} \\ \left(\frac{8k_B T_i}{\pi M_i} \right)^{1/2} & \text{ions,} \end{cases} \quad (19)$$

where the ion temperature is approximated, as proposed by Lee and Lieberman in [3],

$$T_i = \begin{cases} (5800 - T_g) \frac{0.133}{p} + T_g & p > 0.133 \text{ Pa,} \\ 5800 & p \leq 0.133 \text{ Pa.} \end{cases} \quad (20)$$

The momentum transfer cross section σ_{ik}^m is for the purpose of this model crudely approximated with hard sphere model for neutral–neutral and ion–neutral collisions, and with momentum transfer for Rutherford scattering (as proposed by Lieberman and Lichtenberg in [4]) for the case of ion–ion collisions:

$$\sigma_{ik}^m = \begin{cases} (\sigma_i^{\text{LJ}} + \sigma_k^{\text{LJ}})^2 & i = i_+, i_-, i_0, \text{ and } k = k_0, \\ & i = i_0, \text{ and } k = k_+, k_-, \\ \pi b_0^2 \ln \left(\frac{2\lambda_{\text{De}}}{b_0} \right) & i = i_+, i_-, \text{ and } k = k_+, k_-, \end{cases} \quad (21)$$

with Debye length

$$\lambda_{\text{De}} = \left(\frac{\epsilon_0 T_e}{en_e} \right)^{1/2}, \quad (22)$$

classical distance of closest approach

$$b_0 = \frac{q_i q_k e^2}{2\pi \epsilon_0 m_R v_R^2}, \quad (23)$$

reduced mass

$$m_R = \frac{m_i m_k}{m_i + m_k}, \quad (24)$$

and the relative speed being approximated by the mean thermal speed

$$v_R = \bar{v}_i. \quad (25)$$

The δ_{ik} term filters out self-collisions, as collisions between the same species do not affect the species collective behaviour

$$\delta_{ik} = \begin{cases} 1 & \text{for } i = k, \\ 0 & \text{for } i \neq k. \end{cases} \quad (26)$$

Finally, the free diffusion coefficient for positive ions is approximated by

$$D_+^{\text{free}} = \overline{D_{i_+}^{\text{free}}}. \quad (27)$$

Boundary sources

Each k -th species which is lost (or *stuck*) to the plasma boundary can get returned as i -th species, introducing the boundary sources

$$\left(\frac{\delta n_i}{\delta t} \right)_{\text{diff}}^{\text{in}} = - \sum_k r_{ik} \left(\frac{\delta n_k}{\delta t} \right)_{\text{diff}}^{\text{out}} = \frac{A}{V} \sum_k \frac{D_k n_k s_k r_{ik}}{s_k \Lambda + \frac{4D_k}{\bar{v}_k}}. \quad (28)$$

Minimal Allowed Species Density

To prevent the ODE solver from *overshooting* into unphysical negative densities, an additional *artificial* term is added to the right-hand side of (1), ensuring a finite minimal value of particle densities (which can be considered zero). This correction term takes form of

$$\left(\frac{\delta n_i}{\delta t} \right)_{n_{\min}} = \begin{cases} \frac{n_{\min} - n_i}{\tau} & n_i < n_{\min}, \\ 0 & n_i \geq n_{\min}. \end{cases} \quad (29)$$

Here, n_{\min} is set to 1 particle/m³ and τ to 1.0×10^{-10} s, ensuring an adequately fast response.

Electron Energy Density Balance

The balance equation for the electron energy density consists of contributions from the absorbed power, elastic and inelastic electron collisions, electron production and consumption and contribution from diffusion losses of electrons and ions.

$$\frac{d\rho_e}{dt} = \frac{P}{Ve} - \left(\frac{\delta\rho_e}{\delta t}\right)_{\text{el/inel}} - \left(\frac{\delta\rho_e}{\delta t}\right)_{\text{gen/loss}} - \left(\frac{\delta\rho_e}{\delta t}\right)_{\text{el}\rightarrow\text{walls}} - \left(\frac{\delta\rho_e}{\delta t}\right)_{\text{ion}\rightarrow\text{walls}} \quad (30)$$

Contribution of Elastic and Inelastic Collisions

Electron energy density loss rate due to electron collisions is described as

$$\left(\frac{\delta\rho_e}{\delta t}\right)_{\text{el/inel}} = \sum_j R_j \Delta E_{e,j}, \quad (31)$$

with the electron energy loss for j -th reaction $\Delta E_{e,j}$ being

$$\Delta E_{e,j} = \begin{cases} \Delta E_{e,j}^{\text{inel}} & \text{inelastic collisions,} \\ 3 \frac{m_e}{M_{\text{cp},j}} (T_e - T_g) & \text{elastic collisions,} \\ 0 & \text{heavy species collisions or } a_{ej}^{\text{R}} = 0, \end{cases} \quad (32)$$

and

$$T_e = \frac{2}{3} \frac{\rho_e}{n_e}. \quad (33)$$

The electron density n_e is not resolved explicitly, but rather calculated from plasma charge neutrality

$$n_e = \sum_i n_i q_i. \quad (34)$$

For that reason, T_e might reach non-physically low values, when ρ_e governed directly by (30) is much greater than $\sum_i n_i q_i$. A correction is introduced to T_e in the form

of

$$T_e = \max \left(T_g, \frac{2}{3} \frac{\rho_e}{n_e} \right), \quad (35)$$

to help with the solution stability, as the electron temperature will converge to the neutral gas temperature at low electron energy, due to dominance of elastic processes.

Electron Generation and Loss Contribution

Rate of change of electron energy density due to generation and loss of electrons is described by

$$\left(\frac{\delta \rho_e}{\delta t} \right)_{\text{gen/loss}} = \frac{3}{2} T_e \sum_j (a_{ej}^R - a_{ej}^L) R_j. \quad (36)$$

Energy Loss by Electron Transport

Under a Maxwellian energy distribution assumption, each electron lost through the plasma boundary sheath takes away $2k_B T_e$ of energy with it [4], which gives

$$\left(\frac{\delta \rho_e}{\delta t} \right)_{\text{el} \rightarrow \text{walls}} = -2T_e \left(\frac{\delta n_e}{\delta t} \right)_{\text{walls}}, \quad (37)$$

while the total charge flux needs to be zero, yielding

$$\left(\frac{\delta n_e}{\delta t} \right)_{\text{walls}} = \sum_i q_i \left(\frac{\delta n_i}{\delta t} \right)_{\text{diff}}. \quad (38)$$

Energy Loss by Ion Transport

If it is assumed, that ions leave the plasma boundary sheath with the Bohm velocity, each positive ion removed from the plasma takes away $\frac{1}{2}k_B T_e$ of kinetic energy, as well as sheath voltage acceleration energy [4]

$$\left(\frac{\delta \rho_e}{\delta t} \right)_{\text{ion} \rightarrow \text{walls}} = -\frac{1}{2} T_e \sum_{i_+} \left(\frac{\delta n_{i_+}}{\delta t} \right)_{\text{diff}} - \bar{V}_s \sum_{i_+} q_{i_+} \left(\frac{\delta n_{i_+}}{\delta t} \right)_{\text{diff}}. \quad (39)$$

The last open parameter in the system is the mean sheath voltage \overline{V}_s , which, according to Lieberman and Lichtenberg [4], can be approximated by

$$\overline{V}_s = T_e \cdot \ln \left(\frac{\overline{M}_{i+}}{2\pi m_e} \right)^{1/2}. \quad (40)$$

This value of \overline{V}_s is only consistent with ICP plasma sources.

Minimal Allowed Electron Energy Density

To prevent the ODE solver from *overshooting* into unphysical negative ϱ_e , an additional *artificial* term is added to the right-hand side of (30), ensuring a finite minimal value of electron energy density (which can be considered zero). This correction term takes form of

$$\left(\frac{\delta \varrho_e}{\delta t} \right)_{\varrho_{\min}} = \begin{cases} \frac{\varrho_{e,\min} - \varrho_e}{\tau} & \varrho_e < \varrho_{e,\min}, \\ 0 & \varrho_e \geq \varrho_{e,\min}. \end{cases} \quad (41)$$

Here, $\varrho_{e,\min}$ was set to 1 eV/m³ and τ to 1.0×10^{-10} s, ensuring an adequately fast response.

References

- [1] S. Schröter, A. R. Gibson, M. J. Kushner, T. Gans, and D. O’Connell, “Numerical study of the influence of surface reaction probabilities on reactive species in an rf atmospheric pressure plasma containing humidity,” *Plasma Physics and Controlled Fusion*, vol. 60, p. 014035, Jan. 2018.
- [2] E. Stoffels, W. W. Stoffels, D. Vender, M. Haverlag, G. M. W. Kroesen, and F. J. de Hoog, “Negative Ions in Low Pressure Discharges,” *Contributions to Plasma Physics*, vol. 35, no. 4-5, pp. 331–357, 1995.
- [3] C. Lee and M. A. Lieberman, “Global model of Ar, O₂, Cl₂, and Ar/O₂ high-density plasma discharges,” *Journal of Vacuum Science & Technology A: Vacuum, Surfaces, and Films*, vol. 13, pp. 368–380, Mar. 1995.
- [4] M. A. Lieberman and A. J. Lichtenberg, *Principles of plasma discharges and materials processing*. Hoboken, N.J: Wiley-Interscience, 2nd ed ed., 2005. OCLC: ocm56752658.