

Week 6 — Law of Large Numbers, Duality, Conic Programming

AD7031 Optimization Under Uncertainty • Instructor: Prof. H. Park • Oct 3, 2026

1. Law of large numbers

$\tilde{\xi}_1, \tilde{\xi}_2, \tilde{\xi}_3, \dots$: an infinite sequence of **independent and identically distributed (i.i.d.)** random variables,

each with expected value μ

Law of large numbers (strong law, SLLN)


$$\mathbb{P}\left(\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i \in [N]} \tilde{\xi}_i = \mu\right) = 1$$

sample average $\longrightarrow$ expected value

Example portfolio: $\tilde{\xi}_i$ = return of an asset on day i , i.i.d. over days

asset	expected return μ	σ
bond	0.06	0.10
tech	0.10	0.20
energy	0.15	0.30

$$\frac{1}{N} \sum_{i \in [N]} \tilde{\xi}_i = \text{average return over the first } N \text{ days}$$



2. Duality theory

in this course, we focus on **convex** functions.

Primal Problem

$$\begin{aligned} \min \quad & f_0(x) \\ \text{s.t.} \quad & x \in \mathbb{R}^N \\ & f_i(x) \leq 0 \quad \forall i \in [I] \end{aligned} \tag{P}$$

Dual Problem

$$\begin{aligned} \max_{\theta} \min_{x \in \mathbb{R}^N} \quad & f_0(x) + \sum_{i \in [I]} \theta_i f_i(x) \\ \text{s.t.} \quad & \theta \in \mathbb{R}^I \\ & \theta \geq 0 \end{aligned} \tag{D}$$

$\begin{aligned} \min \quad & f_0(x) \\ \text{s.t.} \quad & x \in \mathbb{R}^N \\ & f_i(x) \leq 0 \quad \forall i \in [I] \end{aligned}$
$= \min_{x \in \mathbb{R}^N} f_0(x) + \underbrace{\max_{\theta \in \mathbb{R}_+^I} \sum_{i \in [I]} \theta_i f_i(x)}_{\leq 0}$

if $f_i(x) \leq 0 \quad \forall i$, then $= 0$

if $\exists i : f_i(x) > 0$, then $= +\infty$

$$\geq \max_{\theta \in \mathbb{R}_+^I} \min_{x \in \mathbb{R}^N} f_0(x) + \sum_{i \in [I]} \theta_i f_i(x)$$

Properties:

— **weak duality** $\min(P) \geq \max(D)$ always holds.

— **strong duality:** $\min(P) = \max(D)$ holds

when the problem (P) is convex and the Slater's condition holds

$$\exists x \in \mathbb{R}^N, \text{ s.t. } f_i(x) < 0 \quad \forall i \in [I]$$

In this course, we assume that strong duality holds for convex problem!!

3. Example: linear program

Primal LP:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & x \in \mathbb{R}^N \\ & Ax \leq b \end{array} \qquad \begin{array}{ll} f_0(x) = c^\top x \\ f_i(x) = a_i^\top x - b_i & \forall i \in [I] \end{array}$$

$$\begin{aligned} f_i(x) &\leq 0 \\ \iff a_i^\top x - b_i &\leq 0 \\ \iff a_i^\top x &\leq b_i \end{aligned}$$

Dual LP:

$$\begin{aligned} & \max_{\theta \in \mathbb{R}_+^I} \min_{x \in \mathbb{R}^N} c^\top x + \sum_{i \in [I]} \theta_i (a_i^\top x - b_i) \\ &= \max_{\theta \in \mathbb{R}_+^I} \min_{x \in \mathbb{R}^N} c^\top x + \sum_{i \in [I]} \theta_i a_i^\top x - \sum_{i \in [I]} \theta_i b_i \\ &= \max_{\theta \in \mathbb{R}_+^I} - \sum_{i \in [I]} \theta_i b_i + \underbrace{\min_{x \in \mathbb{R}^N} \left(c + \sum_{i \in [I]} \theta_i a_i \right)^\top x}_{= \begin{cases} 0 & \text{if } c + \sum_{i \in [I]} \theta_i a_i = 0 \\ -\infty & \text{if otherwise.} \end{cases}} \\ &= \max_{\substack{\theta \in \mathbb{R}_+^I \\ A^\top \theta = -c}} -b^\top \theta \end{aligned}$$

LP strong duality : if there exists $x \in \mathbb{R}^N : Ax \leq b$

$$\text{or } \theta \in \mathbb{R}^I : A^\top \theta = -c, \theta \geq 0$$

then strong duality holds


4. Conic programming

a conic program is a convex optimization problem.

— a **linear** objective function.

— a feasible set defines through the intersection of halfspaces $Ax \leq b$ or a proper cone C .

Def (cone) a set C is a cone if $\forall x \in C$, we have $\lambda x \in C \quad \forall \lambda \geq 0$



Def (proper cone) $\mathbb{R}^N$ is a cone but not pointed.

C is a proper cone if it's **convex, closed, pointed**


if $x, -x \in C$ then $x = 0$.

Generic conic program:

$$\begin{aligned} \min \quad & c^\top x \\ \text{s.t.} \quad & x \in \mathbb{R}^N \\ & Ax \preceq_C b \end{aligned} \quad \Longleftrightarrow \quad b - Ax \in C$$


Example:

1) **Linear program:** $C = \mathbb{R}_+^K$



$$\begin{aligned} b - Ax &\in \mathbb{R}_+^K \\ \Longleftrightarrow b - Ax &\geq 0 \quad \Longleftrightarrow Ax \leq b \end{aligned}$$

2) **Second-order cone:** $C = \text{SOC}^{N+1} = \{(x, t) \in \mathbb{R}^N \times \mathbb{R} : \|x\|_2 \leq t\}$ “ice-cream” cone



3) **Semidefinite program:** $C = \mathbb{S}_+^K$


$\mathbb{S}^K$: set of symmetric matrices in $\mathbb{R}^{K \times K}$ ($A^\top = A$)

$\mathbb{S}_+^K$: set of positive semidefinite (symmetric) matrices in $\mathbb{R}^{K \times K}$

$$A \in \mathbb{S}_+^K \text{ if } z^\top A z \geq 0 \quad \forall z \in \mathbb{R}^K$$

Generic semidefinite program:

$$\begin{aligned} \min \quad & c^\top x \\ \text{s.t.} \quad & x \in \mathbb{R}^N \\ & A_1 x_1 + A_2 x_2 + \cdots + A_N x_N \preceq_{\mathbb{S}_+^K} B \end{aligned} \quad \rightarrow \text{usually ignored}$$



4) **Several cones at once:** $C = C_1 \times C_2$ is again a proper cone

$$b_1 - A_1 x \in C_1 \text{ and } b_2 - A_2 x \in C_2 \iff \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} - \begin{bmatrix} A_1 \\ A_2 \end{bmatrix} x \in C_1 \times C_2$$

Example: portfolio

$$\begin{aligned} \max \quad & \mu^\top x \\ \text{s.t.} \quad & x \in \mathbb{R}^N \\ & e^\top x \leq 1 \iff 1 - e^\top x \in \mathbb{R}_+ && \text{budget} \\ & x^\top \Sigma x \leq \bar{\sigma}^2 \iff (L^\top x, \bar{\sigma}) \in \text{SOC}^{N+1} && \text{variance bound} \\ & x \geq 0 \iff x \in \mathbb{R}_+^N && \text{nonnegative weights} \end{aligned}$$

$$\Sigma = LL^\top \text{ (Cholesky)} \Rightarrow x^\top \Sigma x = \|L^\top x\|_2^2, \text{ so } x^\top \Sigma x \leq \bar{\sigma}^2 \iff \|L^\top x\|_2 \leq \bar{\sigma}$$

$$\Rightarrow C = \mathbb{R}_+^{N+1} \times \text{SOC}^{N+1} \text{ — a SOCP} \quad (\max \mu^\top x = -\min(-\mu)^\top x, \text{ so } c = -\mu)$$

$$\text{the coding session, } N = 2: \quad \mu = \begin{bmatrix} 0.10 \\ 0.15 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} 0.01 & 0.001 \\ 0.001 & 0.04 \end{bmatrix}, \quad \bar{\sigma} = 0.13$$

$$L = \begin{bmatrix} 0.1 & 0 \\ 0.01 & 0.19975 \end{bmatrix}, \quad L^\top x = \begin{bmatrix} 0.1x_1 + 0.01x_2 \\ 0.19975x_2 \end{bmatrix}, \quad \|L^\top x\|_2^2 = 0.01x_1^2 + 0.002x_1x_2 + 0.04x_2^2 \checkmark$$

$$\underbrace{\begin{bmatrix} 1 - x_1 - x_2 \\ x_1 \\ x_2 \end{bmatrix}}_{\in \mathbb{R}_+^3} \quad \underbrace{\begin{bmatrix} 0.1x_1 + 0.01x_2 \\ 0.19975x_2 \\ 0.13 \end{bmatrix}}_{\in \text{SOC}^3} \implies b - Ax \in \mathbb{R}_+^3 \times \text{SOC}^3$$

5. Conic programming duality

Def (*dual cone*)

$$C^* = \{ y \in \mathbb{R}^K : \langle y, x \rangle \geq 0 \quad \forall x \in C \}$$

Self dual cones ($C = C^*$)

— $C = \mathbb{R}_+^K$

— $C = \text{second-order cone}$

— $C = \mathbb{S}_+^K$

— products of these: $C = C_1 \times C_2, \quad C^* = C_1^* \times C_2^* = C$

Primal conic program:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & x \in \mathbb{R}^N \\ & Ax \preceq_C b \end{array}$$

Dual conic program:

$$\begin{array}{ll} \max & -\theta^\top b \\ \text{s.t.} & \theta \in C^* \\ & A^\top \theta = -c \end{array}$$