

# Exercise 7 & 8

## Probability Theory 2020 Autumn

Hanmo Chen  
2020214276

December 11, 2020

### 1 Problem 1

Denote the entries in  $U$  as  $u_{ij}$  and entries in  $Y$  as  $Y_{ij}$ , so

$$Y_{ij} = \sum_{r,s} u_{ri} u_{sj} X_{rs} \quad (1)$$

Also we have,

$$\text{Cov}(X_{ij}, X_{mn}) = \begin{cases} 2, & i = j = m = n \\ 1, & (i, j) = (m, n) \text{ or } (i, j) = (n, m), i \neq j \\ 0, & \text{otherwise} \end{cases} \quad (2)$$

Thus

$$\begin{aligned} \text{Cov}(Y_{ij}, Y_{mn}) &= \text{Cov} \left( \sum_{r,s} u_{ri} u_{sj} X_{rs}, \sum_{p,q} u_{pm} u_{qn} X_{pq} \right) \\ &= 2 \sum_{r=1}^n u_{ri} u_{rj} u_{rm} u_{rn} + \sum_{r \neq s} u_{ri} u_{sj} u_{rm} u_{sn} + \sum_{r \neq s} u_{ri} u_{sj} u_{rn} u_{sm} \\ &= \sum_{r,s} [u_{ri} u_{sj} u_{rm} u_{sn} + u_{ri} u_{sj} u_{rn} u_{sm}] \\ &= \left( \sum_r u_{ri} u_{rm} \right) \left( \sum_s u_{sj} u_{sn} \right) + \left( \sum_r u_{ri} u_{rn} \right) \left( \sum_s u_{sj} u_{sm} \right) \end{aligned} \quad (3)$$

Denote the column vectors in  $U$  as  $\mathbf{u}_i, i = 1, 2, 3, \dots, N$ , so  $\mathbf{u}_i \cdot \mathbf{u}_j = \sum_r u_{ri} u_{rj} = \delta_{ij}$ .

$$\text{Cov}(Y_{ij}, Y_{mn}) = \delta_{im} \delta_{jn} + \delta_{jm} \delta_{in} = \begin{cases} 2, & i = j = m = n \\ 1, & (i, j) = (m, n) \text{ or } (i, j) = (n, m), i \neq j \\ 0, & \text{otherwise} \end{cases} \quad (4)$$

Because  $X_{ij}, j \geq i$  are independent Gaussian variables, so the joint distribution of  $Y_{ij}$  is joint Gaussian distribution, which means,

$$\text{Cov}(Y_{ij}, Y_{mn}) = 0 \iff Y_{ij}, Y_{mn} \text{ are independent} \quad (5)$$

So all the entries on and above the diagonal of  $Y$  are independent, and  $Y_{ii} \sim N(0, 2), i = 1, 2, 3, \dots, N$  and  $Y_{ij} \sim N(0, 1), 1 \leq i < j \leq n$ . (It is easy to see that  $\mathbb{E}[Y_{ij}] = 0$ )

## 2 Problem 2

Notice that

1. If  $X_n = 1$ , then  $X_{n+1}, X_{n+2}, \dots$  are independent of  $X_1, X_2, \dots, X_n$
2. There is at least one 1 in any five-in-a-row  $X_i$ s as  $\{X_n, X_{n+1}, \dots, X_{n+4}\}$

So we can split  $X_1, X_2, \dots, X_n$  into a series of epsisodes, each episode  $L_j = [0, \dots, 0, 1]$  is consisted of  $n$  zeros ( $n$  can be 0, 1, 2, 3, 4) and 1 one. And  $L_j, j = 1, 2, \dots, m$  are independent. (For the last episode, if it is ended with 0, we can append 1 to its end and let  $n = n + 1$ .) Denote the length of each episode as  $l_j$ , so  $\sum_{j=1}^m l_j = n$ .

Consider the distribution of  $l_j$ , it can only take values in 1, 2, 3, 4, 5,

- $P(l_j = 1) = P(X_1 = 1) = 0.2$
- $P(l_j = 2) = P(X_1 = 0, X_2 = 1) = 0.16$
- $P(l_j = 3) = P(X_1 = 0, X_2 = 0, X_3 = 1) = 0.128$
- $P(l_j = 4) = P(X_1 = 0, X_2 = 0, X_3 = 0, X_4 = 1) = 0.1024$
- $P(l_j = 5) = P(X_1 = 0, X_2 = 0, X_3 = 0, X_4 = 0) = 0.4096$

So

$$\lim_{n \rightarrow \infty} \frac{S_n}{n} = \lim_{m \rightarrow \infty} \frac{m}{l_1 + l_2 + \dots + l_m} = \lim_{m \rightarrow \infty} \frac{1}{\frac{1}{m} \sum_{j=1}^m l_j} \quad (6)$$

According to Strong Law of Large Numbers,

$$\frac{1}{m} \sum_{j=1}^m l_j \xrightarrow{a.s.} E[l_j] = 3.3616 \quad (7)$$

So

$$\lim_{n \rightarrow \infty} \frac{S_n}{n} \xrightarrow{a.s.} \frac{1}{3.3616} \quad (8)$$