
ORBITS

NOTES ON CALCULATIONS ON CLASSICAL ORBITS UNDER GRAVITY

Hannah Michelle Ellis

November 23, 2025

Contents

1	Plane Polar Coordinate System	3
1.1	Polar unit vectors	3
1.2	Velocity and acceleration in polar coordinates . . .	5
1.2.1	Position	5
1.2.2	Velocity	5
1.2.3	Acceleration	5
2	Gravity	7
2.1	Gravity as a central force	7
3	Angular Momentum	8
3.1	Conservation Of Angular Momentum	8
4	Energy	10
4.1	Potential Energy	10
4.2	Kinetic Energy	10
4.2.1	Total Energy	11
4.3	Conservation of Total Energy	11
5	Geometry Of Ellipses	13
5.1	The Apses	13
5.2	Sum of distances from foci and P	14
5.3	Relationship between the distance from the focus and the Eccentric Anomaly	18
6	Kepler's Equation	20
7	Elliptical Orbits	24
7.1	Change of variables	24
7.2	Solution	26
8	Relationship between geometry and physics	30
8.1	Angular Momentum From Geometry	30
8.2	Total Energy From Geometry	31
8.3	Period	33

8.3.1	The Relationship between Mean Motion and Time Period	34
8.3.2	Period based on physical and geometric properties	34
9	In Plane Position and Velocity over time	36
9.1	Position	36
9.2	Velocity	36
9.2.1	Angular Velocity	36
9.2.2	Radial velocity	37

List of Tables

List of Figures

1	Position and unit vectors in plane polar coordinates	3
2	An Elipse with important values shown	13

1 Plane Polar Coordinate System

In this section we will derive the equations of motion for an object acted upon by gravity in plane polar coordinates. To do this we will first consider plane polar coordinates themselves and the derivatives of the position vector within the coordinate system.




Figure 1: Position and unit vectors in plane polar coordinates

1.1 Polar unit vectors

The unit vectors of plane polar coordinates can be determined from the standard unit vectors as

$$\hat{\mathbf{r}} = \hat{\mathbf{i}} \cos(\theta) + \hat{\mathbf{j}} \sin(\theta) \quad (1.1)$$

$$\hat{\boldsymbol{\theta}} = -\hat{\mathbf{i}} \sin(\theta) + \hat{\mathbf{j}} \cos(\theta) \quad (1.2)$$

Where r is the distance from the origin to the position or alternatively the length of the position vector, and θ is the angle between $\hat{\mathbf{i}}$ and $\mathbf{r}$. Since the unit vectors $\hat{\mathbf{r}}$ and $\hat{\boldsymbol{\theta}}$ depend on θ which changes with time, the unit vectors also change with time. We will begin by looking at the derivative of $\hat{\mathbf{r}}$ with respect to time.

$$\frac{d\hat{\mathbf{r}}}{dt} = \frac{d\theta}{dt} \frac{d}{d\theta} \hat{\mathbf{r}} = \frac{d\theta}{dt} [-\hat{\mathbf{i}} \sin(\theta) + \hat{\mathbf{j}} \cos(\theta)] = \frac{d\theta}{dt} \hat{\boldsymbol{\theta}} \quad (1.3)$$

where we have used equation 1.2 to replace the typical Cartesian coordinate unit vectors.

$$\frac{d\hat{\boldsymbol{\theta}}}{dt} = \frac{d\theta}{dt} \frac{d}{dt} \hat{\boldsymbol{\theta}} = \frac{d\theta}{dt} [-\hat{\mathbf{i}} \cos(\theta) - \hat{\mathbf{j}} \sin(\theta)] = -\frac{d\theta}{dt} \hat{\mathbf{r}} \quad (1.4)$$

Since $\frac{d\hat{\mathbf{r}}}{dt}$ too depends on θ so is also time dependent.

$$\frac{d^2 \hat{\mathbf{r}}}{dt^2} = \frac{d}{dt} \left[\frac{d\theta}{dt} \hat{\boldsymbol{\theta}} \right] = \frac{d^2 \theta}{dt^2} \hat{\boldsymbol{\theta}} + \frac{d\theta}{dt} \frac{d\hat{\boldsymbol{\theta}}}{dt} = \frac{d^2 \theta}{dt^2} \hat{\boldsymbol{\theta}} - \left(\frac{d\theta}{dt} \right)^2 \hat{\mathbf{r}} \quad (1.5)$$

Polar unit vectors summary

The polar coordinate unit vectors are

$$\begin{aligned} \hat{\mathbf{r}} &= \hat{\mathbf{i}} \cos(\theta) + \hat{\mathbf{j}} \sin(\theta) \\ \hat{\boldsymbol{\theta}} &= -\hat{\mathbf{i}} \sin(\theta) + \hat{\mathbf{j}} \cos(\theta) \end{aligned}$$

With the first derivatives being

$$\begin{aligned} \frac{d\hat{\mathbf{r}}}{dt} &= \omega \hat{\boldsymbol{\theta}} \\ \frac{d\hat{\boldsymbol{\theta}}}{dt} &= -\omega \hat{\mathbf{r}} \end{aligned}$$

And the second derivative of $\hat{\mathbf{r}}$ being

$$\frac{d^2 \hat{\mathbf{r}}}{dt^2} = \frac{d\omega}{dt} \hat{\boldsymbol{\theta}} - \omega^2 \hat{\mathbf{r}}$$

with $\omega = \frac{d\theta}{dt}$

1.2 Velocity and acceleration in polar coordinates

Here we will calculate the acceleration in terms of the polar unit vectors.

1.2.1 Position

Starting with the position vector

$$\mathbf{r} = r\hat{\mathbf{r}} \quad (1.6)$$

This is the length of the position vector in the direction of $\hat{\mathbf{r}}$.

1.2.2 Velocity

The first derivative will be the velocity of the position vector in terms of the polar unit vectors

$$\frac{d\mathbf{r}}{dt} = \frac{dr}{dt}\hat{\mathbf{r}} + r\frac{d\hat{\mathbf{r}}}{dt} = \frac{dr}{dt}\hat{\mathbf{r}} + r\omega\hat{\boldsymbol{\theta}} \quad (1.7)$$

The velocity is the sum of the radial velocity in the radial direction $\dot{r}$ and the angular part of the velocity $r\omega$ in the angular direction.

1.2.3 Acceleration

$$\begin{aligned} \frac{d^2\mathbf{r}}{dt^2} &= \frac{d^2r}{dt^2}\hat{\mathbf{r}} + \frac{dr}{dt}\frac{d\hat{\mathbf{r}}}{dt} + \frac{dr}{dt}\omega\hat{\boldsymbol{\theta}} + r\frac{d\omega}{dt}\hat{\boldsymbol{\theta}} + r\omega\frac{d\hat{\boldsymbol{\theta}}}{dt} \\ &= \frac{d^2r}{dt^2}\hat{\mathbf{r}} + \frac{dr}{dt}\omega\hat{\boldsymbol{\theta}} + \frac{dr}{dt}\omega\hat{\boldsymbol{\theta}} + r\frac{d\omega}{dt}\hat{\boldsymbol{\theta}} - r\omega^2\hat{\mathbf{r}} \\ &= \left[\frac{d^2r}{dt^2} - r\omega^2 \right] \hat{\mathbf{r}} + \left[2\frac{dr}{dt}\omega + r\frac{d\omega}{dt} \right] \hat{\boldsymbol{\theta}} \end{aligned} \quad (1.8)$$

Acceleration in polar coordinates summary

The position vector

$$\mathbf{r} = r\hat{\mathbf{r}}$$

Velocity

$$\frac{d\mathbf{r}}{dt} = \frac{dr}{dt}\hat{\mathbf{r}} + r\omega\hat{\boldsymbol{\theta}}$$

Acceleration

$$\frac{d^2\mathbf{r}}{dt^2} = \left[\frac{d^2r}{dt^2} - r\omega^2 \right] \hat{\mathbf{r}} + \left[2\frac{dr}{dt}\omega + r\frac{d\omega}{dt} \right] \hat{\boldsymbol{\theta}}$$

2 Gravity

2.1 Gravity as a central force

The force between two particles a distance r apart with mass M and m respectively is given by

$$\mathbf{F} = -\frac{GMm}{r^2}\hat{\mathbf{r}} \quad (2.1)$$

The acceleration on the particle of mass m due to the force of gravity between the two particles is

$$\ddot{\mathbf{r}} = \frac{\mathbf{F}}{m} = -\frac{GM}{r^2}\hat{\mathbf{r}} \quad (2.2)$$

Using equation 1.8 we can split the acceleration into radial and angular parts

$$-\frac{GM}{r^2} = \ddot{r} - r\omega^2 \quad (2.3)$$

$$0 = r\ddot{\theta} + 2\dot{r}\dot{\theta} \quad (2.4)$$

3 Angular Momentum

The angular momentum of a particle is given by

$$\begin{aligned}\mathbf{L} &= \mathbf{r} \times \mathbf{p} \\ &= m\mathbf{r} \times \frac{d\mathbf{r}}{dt} \\ &= mr\hat{\mathbf{r}} \times \left[\dot{r}\hat{\mathbf{r}} + r\omega\hat{\boldsymbol{\theta}} \right] \\ &= m \left[r\dot{r}\hat{\mathbf{r}} \times \hat{\mathbf{r}} + r^2\omega\hat{\mathbf{r}} \times \hat{\boldsymbol{\theta}} \right] \\ &= mr^2\omega\hat{\mathbf{k}}\end{aligned}\tag{3.1}$$

We can also write this as $\mathbf{L} = l\hat{\mathbf{k}}$ where

$$l = mr^2\omega\tag{3.2}$$

3.1 Conservation Of Angular Momentum

We can show now that angular momentum is conserved under a central force by taking the derivative with respect to time.

$$\frac{d\mathbf{L}}{dt} = \frac{dl\hat{\mathbf{k}}}{dt} = \frac{dl}{dt}\hat{\mathbf{k}}\tag{3.3}$$

$$\begin{aligned}\frac{dl}{dt} &= m \frac{d}{dt} [r^2\omega] \\ &= m \left[\frac{dr^2}{dt}\omega + r^2\frac{d\omega}{dt} \right] \\ &= m \left[\frac{dr^2}{dr} \frac{dr}{dt}\omega + r^2\frac{d\omega}{dt} \right] \\ &= m \left[2r\frac{dr}{dt}\omega + r^2\frac{d\omega}{dt} \right] \\ &= mr \left[2\frac{dr}{dt}\omega + r\frac{d\omega}{dt} \right]\end{aligned}\tag{3.4}$$

The part inside the brackets is the angular acceleration which for a central force is zero¹ and so

$$\begin{aligned}\frac{dl}{dt} &= mr \times 0 \\ &= 0\end{aligned}\tag{3.5}$$

Which means that the angular momentum is constant.

¹see equation 2.4

4 Energy

4.1 Potential Energy

The potential energy is the work done in moving the particle from a radius of ∞ to r . So the work done is given by

$$\begin{aligned}U(r) &= \int_{\infty}^r \mathbf{F}(r') \cdot d\mathbf{r}' \\U(r) &= \int_{\infty}^r -\frac{GMm}{r'^2} \hat{\mathbf{r}} \cdot dr' \hat{\mathbf{r}} \\U(r) &= -GMm \int_{\infty}^r \frac{dr'}{r'^2} \\U(r) &= -GMm \left[\frac{-1}{r'} \right]_{\infty}^r \\U(r) &= GMm \left[\frac{1}{\infty} - \frac{1}{r} \right] \\U(r) &= -\frac{GMm}{r}\end{aligned}\tag{4.1}$$

4.2 Kinetic Energy

The kinetic energy of a particle is given by

$$T(\dot{\mathbf{r}}) = \frac{1}{2} m \dot{\mathbf{r}} \cdot \dot{\mathbf{r}}\tag{4.2}$$

using equation 1.2.3 gives

$$\begin{aligned}T(\dot{\mathbf{r}}) &= \frac{1}{2} m \left[\dot{r} \hat{\mathbf{r}} + r\omega \hat{\boldsymbol{\theta}} \right] \cdot \left[\dot{r} \hat{\mathbf{r}} + r\omega \hat{\boldsymbol{\theta}} \right] \\&= \frac{1}{2} m \dot{r}^2 + \frac{1}{2} m r^2 \omega^2\end{aligned}\tag{4.3}$$

we can also make the substitution using using the magnitude of the angular momentum² $l = mr^2\omega$

²equation 3.2

$$\begin{aligned}
l &= mr^2\omega \\
\omega &= \frac{l}{mr^2}
\end{aligned}
\tag{4.4}$$

using equation 4.4 in equation 4.3 leads to

$$\begin{aligned}
T(\dot{r}) &= \frac{1}{2}m\dot{r}^2 + \frac{1}{2}mr^2\omega^2 \\
&= \frac{1}{2}m\dot{r}^2 + \frac{1}{2}mr^2\frac{l^2}{m^2r^4} \\
&= \frac{1}{2}m\dot{r}^2 + \frac{l^2}{2mr^2}
\end{aligned}
\tag{4.5}$$

4.2.1 Total Energy

Finally the total energy can be found by putting together the equation for kinetic energy and potential energy

$$\begin{aligned}
E(r, \dot{r}) &= T(\dot{r}) + U(r) \\
E(r, \dot{r}) &= \frac{1}{2}m\dot{r}^2 + \frac{l^2}{2mr^2} - \frac{GMm}{r}
\end{aligned}
\tag{4.6}$$

sometimes the last two terms of equation 4.6 are put together to form a new potential energy type term that is dependent only on the radius and the angular momentum.

4.3 Conservation of Total Energy

Just like with angular momentum we can show that total energy is conserved by taking the derivative with respect to time.

$$\begin{aligned}
\frac{dE}{dt} &= \frac{d}{dt} \left[\frac{1}{2} m \dot{r}^2 + \frac{l^2}{2mr^2} - \frac{GMm}{r} \right] \\
&= \frac{1}{2} m \frac{d\dot{r}^2}{dt} + \frac{l^2}{2m} \frac{dr^{-2}}{dt} - GMm \frac{dr^{-1}}{dt} \\
&= \frac{1}{2} m \frac{d\dot{r}^2}{d\dot{r}} \frac{d\dot{r}}{dt} + \frac{l^2}{2m} \frac{dr^{-2}}{dr} \frac{dr}{dt} - GMm \frac{dr^{-1}}{dr} \frac{dr}{dt} \\
&= \frac{1}{2} m (2\dot{r}) \ddot{r} + \frac{l^2}{2m} \left(\frac{-2}{r^3} \right) \dot{r} + \frac{GMm}{r^2} \dot{r} \\
&= m\dot{r}\ddot{r} - \frac{l^2}{mr^3} \dot{r} + \frac{GMm}{r^2} \dot{r}
\end{aligned} \tag{4.7}$$

using the magnitude of the angular momentum³ $l = mr^2\omega$

$$\begin{aligned}
\frac{dE}{dt} &= m\dot{r}\ddot{r} - \frac{l^2}{mr^3} \dot{r} + \frac{GMm}{r^2} \dot{r} \\
&= m\dot{r}\ddot{r} - \frac{m^2 r^4 \omega^2}{mr^3} \dot{r} + \frac{GMm}{r^2} \dot{r} \\
&= m\dot{r}\ddot{r} - mr\omega^2 \dot{r} + \frac{GMm}{r^2} \dot{r} \\
&= m\dot{r} \left[\ddot{r} - r\omega^2 + \frac{GM}{r^2} \right]
\end{aligned} \tag{4.8}$$

using the radial part of the acceleration⁴ $-\frac{GM}{r^2} = \ddot{r} - r\omega^2$ we get

$$\begin{aligned}
\frac{dE}{dt} &= m\dot{r} \left[-\frac{GM}{r^2} + \frac{GM}{r^2} \right] \\
&= m\dot{r} [0] \\
&= 0
\end{aligned} \tag{4.9}$$

This then shows that the total energy does not change with respect to time, so is a constant and also conserved.

³equation 3.2

⁴equation 2.3

5 Geometry Of Ellipses




Figure 2: An Ellipse with important values shown

In figure 2 a point P on the ellipse forms an angle E with the origin and the x axis, this angle is called the *Eccentric Anomaly*.

Also the angle θ is formed at the ellipse's focus between the point on the ellipse and the axis.

We also define the distances a and p as the apoapsis and periapsis distances respectively.

The semi major axis has been labeled as A , so that the total width of the ellipse is $2A$. Also shown on the diagram is the semi minor axis B , so the total height of the ellipse is $2B$.

For this text we will take the elliptical eccentricity to be defined as the ratio of the focus and the semi major axis.

$$e = \frac{f}{A} \quad (5.1)$$

5.1 The Apses

Before we get any further, we will look at the apses. From figure 2 we can see that

$$p = A - f = A - eA = A(1 - e) \quad (5.2)$$

and that

$$a = A + f = A + eA = A(1 + e) \quad (5.3)$$

also

$$a + p = A - f + A + f = 2A \quad (5.4)$$

$$a - p = A + f - A + f = 2f \quad (5.5)$$

which can give us the eccentricity as defined by the apses

$$e = \frac{f}{A} = \frac{2f}{2A} = \frac{a - p}{a + p} \quad (5.6)$$

5.2 Sum of distances from foci and P

One property of an ellipse is that the sum of the distances from any point on the ellipse and the foci is a constant. This fact allows you to draw an ellipse using two pins and a piece of string. We will use this fact and look at two special cases, before looking at the general case and deriving the equation for an ellipse. Before we do we will define what the sum is here.

$$S = d + r \quad (5.7)$$

Special Case 1: Point on the x axis When the point is at $(A, 0)$. The distance d is $f + A$ and the distance r is $A - f$ from the diagram above. So our sum becomes

$$S = d + r = f + A + A - f = 2A \quad (5.8)$$

Special Case 2: Point on the y axis When the point is on the y axis, it is at the point $(0, B)$. The distance d is $\sqrt{f^2 + B^2}$ and the distance r is also $\sqrt{f^2 + B^2}$ from the diagram above. So our sum is

$$S = 2\sqrt{f^2 + B^2} \quad (5.9)$$

and if the sum is a constant then this will be equal to the value obtained before⁵.

$$S = 2\sqrt{f^2 + B^2} = 2A \quad (5.10)$$

and so we now have the relationship between the semi major and minor axes and the focus

$$A^2 = f^2 + B^2 \quad (5.11)$$

This can also be used to get B in terms of A using the connection to the eccentricity⁶

$$\begin{aligned} B^2 &= A^2 - f^2 \\ &= A^2 - (eA)^2 \\ &= A^2(1 - e^2) \\ \Rightarrow B &= A\sqrt{1 - e^2} \end{aligned} \quad (5.12)$$

additionally, if we use the identity $x^2 - y^2 = (x + y)(x - y)$ we can also get the following

$$B = A\sqrt{1 - e^2} = \sqrt{A(1 + e) \times A(1 - e)} = \sqrt{ap} \quad (5.13)$$

where we have used equations 5.2⁷ and 5.3⁸ to substitute in for a and b respectively.

General case In general for a point at (x, y) we get the following

$$r^2 = (x - f)^2 + y^2 \quad (5.14)$$

$$d^2 = (x + f)^2 + y^2 \quad (5.15)$$

by looking at the triangles formed in figure 2 with hypotenuse being r and d respectively. The sum⁹ then becomes

$$\begin{aligned} S &= 2A = \sqrt{(x - f)^2 + y^2} + \sqrt{(x + f)^2 + y^2} \\ 2A - \sqrt{(x - f)^2 + y^2} &= \sqrt{(x + f)^2 + y^2} \end{aligned} \quad (5.16)$$

⁵equation 5.8 on page ??

⁶equation 5.1 on page 13

⁷on page 13

⁸on page 14

⁹equation 5.7 on page 14

squaring both sides gives

$$\begin{aligned}
\left(2A - \sqrt{(x-f)^2 + y^2}\right)^2 &= (x+f)^2 + y^2 \\
4A^2 - 4A\sqrt{(x-f)^2 + y^2} + (x-f)^2 + y^2 &= (x+f)^2 + y^2 \\
4A^2 - 4A\sqrt{(x-f)^2 + y^2} + x^2 - 2xf + f^2 &= x^2 + 2xf + f^2 \\
4A^2 - 4A\sqrt{(x-f)^2 + y^2} &= 4xf \\
4A^2 - 4xf &= 4A\sqrt{(x-f)^2 + y^2} \\
\frac{4A^2 - 4xf}{4A} &= \sqrt{(x-f)^2 + y^2} \\
\sqrt{(x-f)^2 + y^2} &= A - \frac{xf}{A} \quad (5.17)
\end{aligned}$$

Squaring both sides gives again gives

$$\begin{aligned}
(x-f)^2 + y^2 &= \left(A - \frac{xf}{A}\right)^2 \\
x^2 - 2xf + f^2 + y^2 &= A^2 - 2\frac{xfA}{A} + \frac{x^2f^2}{A^2} \\
x^2 - 2xf + f^2 + y^2 &= A^2 - 2xf + e^2x^2 \\
(1-e^2)x^2 + y^2 &= A^2 - f^2 \\
(1-e^2)x^2 + y^2 &= A^2(1-e^2) \\
\frac{(1-e^2)x^2}{A^2(1-e^2)} + \frac{y^2}{A^2(1-e^2)} &= 1 \\
\frac{x^2}{A^2} + \frac{y^2}{B^2} &= 1 \quad (5.18)
\end{aligned}$$

Which is the cartesian form of an ellipse, there by showing that the ellipse does indeed obey the constraint that the sum of the distances from the foci is a constant.

Polar Form Once again returning to figure 2 we can see that

$$x = f + r \cos \theta \quad (5.19)$$

$$y = r \sin \theta \quad (5.20)$$

The sum¹⁰ then becomes

$$\begin{aligned} S &= 2A = r + d \\ 2A - r &= d \end{aligned} \tag{5.21}$$

We can square both sides to get

$$\begin{aligned} (2A - r)^2 &= d^2 \\ 4A^2 - 4Ar + r^2 &= (x + f)^2 + y^2 \\ &= x^2 + 2fx + f^2 + y^2 \\ &= (f + r \cos \theta)^2 + 2fx + f^2 + r^2 \sin^2 \theta \\ &= f^2 + 2fr \cos \theta + r^2 \cos^2 \theta + 2f(f + r \cos \theta) + f^2 \\ &\quad + r^2 \sin^2 \theta \\ &= f^2 + 2fr \cos \theta + 2f^2 + 2fr \cos \theta + f^2 \\ &\quad + r^2(\sin^2 \theta + \cos^2 \theta) \\ &= 4f^2 + 4fr \cos \theta + r^2 \\ A^2 - Ar &= f^2 + fr \cos \theta \\ A^2 - f^2 &= (A + f \cos \theta) r \\ A^2 - e^2 A^2 &= (A + eA \cos \theta) r \\ A^2 (1 - e^2) &= A (1 + e \cos \theta) r \\ A (1 - e^2) &= (1 + e \cos \theta) r \end{aligned} \tag{5.22}$$

Which with some rearranging results in the polar form for the equation of an ellipse

$$r = \frac{A(1 - e^2)}{1 + e \cos \theta} \tag{5.23}$$

¹⁰equation 5.7 14

5.3 Relationship between the distance from the focus and the Eccentric Anomaly

Looking at figure 2 again we can see that x and y can also be given by

$$x = A \cos E \quad (5.24)$$

$$y = B \sin E = A\sqrt{1 - e^2} \sin E \quad (5.25)$$

We can also see that r can be given by

$$\begin{aligned} r^2 &= (x - f)^2 + y^2 \\ &= x^2 - 2xf + f^2 + y^2 \\ &= A^2 \cos^2 E - 2eA^2 \cos E + e^2 A^2 + A^2(1 - e^2) \sin^2 E \\ \left(\frac{r}{A}\right)^2 &= \cos^2 E - 2e \cos E + e^2 + (1 - e^2) \sin^2 E \\ \left(\frac{r}{A}\right)^2 &= \cos^2 E - 2e \cos E + e^2 + \sin^2 E - e^2 \sin^2 E \\ \left(\frac{r}{A}\right)^2 &= (\sin^2 E + \cos^2 E) - 2e \cos E + e^2(1 - \sin^2 E) \\ \left(\frac{r}{A}\right)^2 &= 1 - 2e \cos E + e^2 \cos^2 E \\ \left(\frac{r}{A}\right)^2 &= (1)^2 - 2(1)(e \cos E) + (e \cos E)^2 \\ \left(\frac{r}{A}\right)^2 &= (1 - e \cos E)^2 \end{aligned} \quad (5.26)$$

leading to

$$r = A(1 - e \cos E) \quad (5.27)$$

Summary

$$p = A(1 - e)$$

$$a = A(1 + e)$$

$$f = eA$$

$$A = \frac{a + p}{2}$$

$$f = \frac{a - p}{2}$$

$$e = \frac{a - p}{a + p}$$

$$\begin{aligned} B^2 &= A^2 - f^2 \\ &= A^2(1 - e^2) \\ &= A(1 + e)A(1 - e) \\ &= ap \\ B &= \sqrt{ap} \end{aligned}$$

$$\begin{aligned} r &= \frac{A(1 - e^2)}{1 + e \cos \theta} \\ &= A(1 - e \cos E) \end{aligned}$$

$$\begin{aligned} x &= A \cos E \\ &= f + r \cos \theta \end{aligned}$$

$$\begin{aligned} y &= B \sin E \\ &= r \sin \theta \end{aligned}$$

6 Kepler's Equation

Starting with equation 5.27 that gives the relationship between the distance from the focus and the eccentric anomaly

$$r = A(1 - e \cos E)$$

We can take the derivative with respect to time to get the relationship between the rate of change of distance with time and the eccentric anomaly.

$$\begin{aligned} \frac{dr}{dt} &= A \frac{d}{dt} (1 - e \cos E) \\ &= A \frac{dE}{dt} \frac{d}{dE} (1 - e \cos E) \\ &= eA \sin E \frac{dE}{dt} \end{aligned} \tag{6.1}$$

Using equation 5.23 for the equation of an ellipse in polar form $r = \frac{A(1-e^2)}{1+e \cos \theta}$. We can invert this to get

$$\frac{1}{r} = \frac{1 + e \cos \theta}{A(1 - e^2)} \tag{6.2}$$

which will make taking the derivative with respect to time easier.

$$\begin{aligned} \frac{d}{dt} \left(\frac{1}{r} \right) &= \frac{d}{dt} \left[\frac{1 + e \cos \theta}{A(1 - e^2)} \right] \\ \frac{dr}{dt} \frac{d}{dr} \left(\frac{1}{r} \right) &= \frac{1}{A(1 - e^2)} \frac{d\theta}{dt} \frac{d}{d\theta} [1 + e \cos \theta] \\ -\frac{1}{r^2} \frac{dr}{dt} &= \frac{-e \sin \theta}{A(1 - e^2)} \frac{d\theta}{dt} \end{aligned} \tag{6.3}$$

we can now substitute in equation 6.1 for $\frac{dr}{dt}$ to get

$$-\frac{1}{r^2} eA \sin E \frac{dE}{dt} = \frac{-e \sin \theta}{A(1 - e^2)} \frac{d\theta}{dt} \tag{6.4}$$

We can use equation 5.20 and 5.25 for y to obtain a relationship between $\sin \theta$ and $\sin E$

$$y = r \sin \theta = A\sqrt{1 - e^2} \sin E \quad (6.5)$$

We can use this relationship to replace the $\sin \theta$ term in equation 6.4

$$\begin{aligned} -\frac{1}{r^2} e A \sin E \frac{dE}{dt} &= \frac{-e \sin \theta}{A(1 - e^2)} \frac{d\theta}{dt} \\ &= \frac{-e A \sqrt{1 - e^2} \sin E}{A r (1 - e^2)} \frac{d\theta}{dt} \\ &= \frac{-e \sin E}{r \sqrt{1 - e^2}} \frac{d\theta}{dt} \\ \frac{-e A \sin E}{r^2} \frac{dE}{dt} &= \frac{-e \sin E}{r \sqrt{1 - e^2}} \frac{d\theta}{dt} \\ \frac{dE}{dt} &= \frac{-r^2}{e A \sin E} \frac{-e \sin E}{r \sqrt{1 - e^2}} \frac{d\theta}{dt} \\ \frac{dE}{dt} &= \frac{r}{A \sqrt{1 - e^2}} \frac{d\theta}{dt} \end{aligned} \quad (6.6)$$

Finally we can replace r with equation 5.27 $r = A(1 - e \cos E)$ to get

$$\frac{dE}{dt} = \frac{r}{A \sqrt{1 - e^2}} \frac{d\theta}{dt} \quad (6.7)$$

$$= \frac{A(1 - e \cos E)}{A \sqrt{1 - e^2}} \frac{d\theta}{dt} \quad (6.8)$$

$$= \frac{(1 - e \cos E)}{\sqrt{1 - e^2}} \frac{d\theta}{dt} \quad (6.9)$$

Now we have an equation that links the rate of change of E to the rate of change of θ in terms of just E and θ

Using Angular Momentum given by equation 3.2

$$l = m r^2 \frac{d\theta}{dt}$$

using equation 5.27 and rearranging we obtain

$$\begin{aligned}
 l &= mr^2 \frac{d\theta}{dt} \\
 \frac{d\theta}{dt} &= \frac{l}{mr^2} \\
 \frac{d\theta}{dt} &= \frac{l}{mA^2 (1 - e \cos E)^2}
 \end{aligned} \tag{6.10}$$

which we can now substitute in for $\frac{d\theta}{dt}$ term in equation 6.9

$$\begin{aligned}
 \frac{dE}{dt} &= \frac{(1 - e \cos E)}{\sqrt{1 - e^2}} \frac{d\theta}{dt} \\
 &= \frac{(1 - e \cos E)}{\sqrt{1 - e^2}} \frac{l}{mA^2 (1 - e \cos E)^2} \\
 &= \frac{l}{mA^2 \sqrt{1 - e^2}} \frac{1}{1 - e \cos E} \\
 (1 - e \cos E) \frac{dE}{dt} &= \frac{l}{mA^2 \sqrt{1 - e^2}}
 \end{aligned} \tag{6.11}$$

Now you can see that the right hand side is just a constant, which is defined by properties of the orbit. Namely the angular momentum, the mass of the orbiting object, the semi major axis and the eccentricity. An interesting thing to note is that the term $A^2 \sqrt{1 - e^2}$ can be rewritten as the product AB instead, so linking the major and minor axes. We will however keep it in it's original form for now. Since the right hand side is a constant we will simplify by choosing the constant k for it. This constant k is the *Mean Motion*, later¹¹ we will see that it's connected to the period of the orbit, and ends up being the mean angular velocity of the orbiting body.

$$k = \frac{\frac{l}{m}}{A^2 \sqrt{1 - e^2}} \tag{6.12}$$

¹¹see equation 8.8 on page 33

so finally all that is left to do is to integrate our equation with respect to time.

$$\begin{aligned}\int_{t_0}^t k dt &= \int_{t=t_0}^t (1 - e \cos E) \frac{dE}{dt} dt \\ k(t - t_0) &= \int_{E=0}^E (1 - e \cos E) dE \\ &= \int_0^E 1 dE - e \int_0^E \cos E dE \\ &= E - e \sin E\end{aligned}\tag{6.13}$$

$$\tag{6.14}$$

Note we have chosen t_0 to be the point in time when $E = 0$, which is the same as the statement of Kepler's equation.

$$E - e \sin E = k(t - t_0)\tag{6.15}$$

7 Elliptical Orbits

We can show that a general solution to the equations of motion is an ellipse.

7.1 Change of variables

Firstly we start by letting $r = \frac{1}{u}$, so that

$$\begin{aligned}\frac{dr}{dt} &= \frac{du^{-1}}{dt} \\ &= \frac{du^{-1}}{du} \frac{du}{dt} \\ &= -\frac{1}{u^2} \frac{du}{dt} \\ &= -r^2 \frac{du}{dt}\end{aligned}\tag{7.1}$$

next we introduce an ω by chain rule

$$\begin{aligned}\frac{dr}{dt} &= -r^2 \frac{du}{dt} \\ \frac{dr}{dt} &= -r^2 \frac{du}{d\theta} \frac{d\theta}{dt} \\ &= -r^2 \omega \frac{du}{d\theta}\end{aligned}$$

And using the magnitude of angular momentum¹² $l = mr^2\omega$ or rearranging $\omega = \frac{l}{mr^2}$

¹²equation 3.2

$$\begin{aligned}
\frac{dr}{dt} &= -r^2\omega \frac{du}{d\theta} \\
\frac{dr}{dt} &= -r^2 \frac{l}{mr^2} \frac{du}{d\theta} \\
\frac{dr}{dt} &= -\frac{l}{m} \frac{du}{d\theta} \\
\frac{dr}{dt} &= -\frac{l}{m} \frac{du}{d\theta}
\end{aligned} \tag{7.2}$$

Differentiating again with respect to time gives

$$\begin{aligned}
\frac{dr^2}{dt^2} &= -\frac{l}{m} \frac{d}{dt} \frac{du}{d\theta} \\
&= -\frac{l}{m} \frac{d\theta}{dt} \frac{d}{d\theta} \frac{du}{d\theta} \\
&= -\frac{l}{m} \omega \frac{d^2u}{d\theta^2}
\end{aligned} \tag{7.3}$$

We can substitute the magnitude of angular momentum¹³ $l = mr^2\omega$ again to get

$$\begin{aligned}
\frac{dr^2}{dt^2} &= -\frac{l}{m} \omega \frac{d^2u}{d\theta^2} \\
&= -\frac{mr^2\omega}{m} \omega \frac{d^2u}{d\theta^2} \\
&= -r^2\omega^2 \frac{d^2u}{d\theta^2}
\end{aligned} \tag{7.4}$$

We can now substitute this into the radial part of the acceleration¹⁴

¹³equation 3.2

¹⁴equation 2.3

$$\begin{aligned}
-\frac{GM}{r^2} &= \frac{d^2 r}{dt^2} - r\omega^2 \\
&= -r^2\omega^2 \frac{d^2 u}{d\theta^2} - r\omega^2 \\
&= -r^2\omega^2 \left[\frac{d^2 u}{d\theta^2} + \frac{1}{r} \right] \\
GM &= r^4\omega^2 \left[\frac{d^2 u}{d\theta^2} + \frac{1}{r} \right]
\end{aligned} \tag{7.5}$$

Next we recall that $u = \frac{1}{r}$ and that $\frac{l}{m} = r^2\omega$ and substitute these in

$$\begin{aligned}
GM &= \left(\frac{l}{m} \right)^2 \left[\frac{d^2 u}{d\theta^2} + u \right] \\
\frac{d^2 u}{d\theta^2} + u &= \frac{GMm^2}{l^2}
\end{aligned} \tag{7.6}$$

To simplify this equation we will use $\alpha = \frac{GMm^2}{l^2}$

7.2 Solution

The direct solution to

$$\frac{d^2 u}{d\theta^2} + u = \alpha \tag{7.7}$$

is $u = \alpha$

However because there is also a function that results in zero, we must include that

$$\begin{aligned}
\frac{d^2 u}{d\theta^2} + u &= 0 \\
\frac{d^2 u}{d\theta^2} &= -u
\end{aligned} \tag{7.8}$$

we can take the solution that $u = c \cos \theta$ which gives the general overall solution of

$$u(\theta) = \alpha + c \cos \theta \quad (7.9)$$

We can pick an initial condition when $\theta = 0$ the body is at periapsis or $r = p = A(1 - e)$ which gives $u = \frac{1}{A(1-e)}$ (see the ellipse notes for $p = A(1 - e)$)

$$\begin{aligned} \frac{1}{A(1 - e)} &= \alpha + c \\ \alpha &= \frac{1}{A(1 - e)} - c \end{aligned} \quad (7.10)$$

we could similarly have picked the condition that when $\theta = \pi$ the body is at apoapsis or $r = a = A(1 + e)$ which gives $u = \frac{1}{A(1+e)}$ (see ellipse notes for $a = A(1 + e)$)

$$\begin{aligned} \frac{1}{A(1 + e)} &= \alpha - c \\ \alpha &= \frac{1}{A(1 + e)} + c \end{aligned} \quad (7.11)$$

we can equate our two initial conditions to get c in terms of just a and p

$$\begin{aligned}
\alpha &= \frac{1}{A(1-e)} - c = \frac{1}{A(1+e)} + c \\
\frac{1}{A(1-e)} - \frac{1}{A(1+e)} &= 2c \\
\frac{A(1+e) - A(1-e)}{A^2(1+e)(1-e)} &= 2c \\
\frac{A + eA - A + eA}{A^2(1-e^2)} &= 2c \\
\frac{2eA}{A^2(1-e^2)} &= 2c \\
\frac{e}{A(1-e^2)} &= c
\end{aligned} \tag{7.12}$$

Now that we have c we can use this to determine α in terms of A and e

$$\begin{aligned}
\alpha &= \frac{1}{A(1+e)} + c \\
&= \frac{1}{A(1+e)} + \frac{e}{A(1-e)^2} \\
&= \frac{(1-e)}{A(1+e)(1-e)} + \frac{e}{A(1-e)^2} \\
&= \frac{1-e+e}{A(1-e^2)} \\
&= \frac{1}{A(1-e^2)}
\end{aligned}$$

So feeding all this back in we have

$$\begin{aligned}
u(\theta) &= \alpha + c \cos \theta \\
&= \frac{1}{A(1 - e^2)} + \frac{e}{A(1 - e^2)} \cos \theta \\
&= \frac{1 + e \cos \theta}{A(1 - e^2)}
\end{aligned} \tag{7.13}$$

Now we have a simple form for u we can recall that we chose $u = \frac{1}{r}$, so

$$r = \frac{1 + e \cos \theta}{A(1 - e^2)} \tag{7.14}$$

which is the equation for an ellipse¹⁵.

We also get the interesting result here that

$$\begin{aligned}
\alpha &= \frac{GMm^2}{l^2} = \frac{1}{A(1 - e^2)} \\
A(1 - e^2) &= \frac{l^2}{GMm^2}
\end{aligned} \tag{7.15}$$

which connects the semi major axis and the eccentricity to physical properties, like the masses and the angular momentum. This will help us to work out the geometry from the physics.

¹⁵equation 5.23

8 Relationship between geometry and physics

We saw in the previous chapter that the geometry and the physics were linked by equation 7.15.

8.1 Angular Momentum From Geometry

We can rearrange equation 7.15 to get

$$\begin{aligned}\frac{l^2}{GMm^2} &= A(1 - e^2) \\ \left(\frac{l}{m}\right)^2 &= GMA(1 - e^2)\end{aligned}\tag{8.1}$$

Which can be further rearranged using equations 5.12¹⁶ and 5.13¹⁷

$$\begin{aligned}\left(\frac{l}{m}\right)^2 &= GM\frac{A^2(1 - e^2)}{A} \\ &= GM\frac{B^2}{A} \\ &= GM\frac{ap}{A} \\ &= GM\frac{2ap}{a + p}\end{aligned}\tag{8.2}$$

With some further rearrangement we can obtain the angular momentum per unit mass of the orbiting body

¹⁶page 15

¹⁷page 15

$$\begin{aligned}
\frac{l^2}{GMm^2} &= \frac{2ap}{a+p} \\
\left(\frac{l}{m}\right)^2 &= GM \frac{2ap}{a+p} \\
\frac{l}{m} &= \sqrt{GM \frac{2ap}{a+p}}
\end{aligned} \tag{8.3}$$

8.2 Total Energy From Geometry

The total energy¹⁸ is given by

$$E(r, \dot{r}) = \frac{1}{2}m\dot{r}^2 + \frac{l^2}{2mr^2} - \frac{GMm}{r}$$

making this total energy per unit mass of the orbiting body instead

$$\frac{E}{m} = \frac{1}{2}\dot{r}^2 + \frac{1}{2r^2} \left(\frac{l}{m}\right)^2 - \frac{GM}{r} \tag{8.4}$$

¹⁸equation 4.6

If we choose one of the apses, say the periapsis we know that at this point $\dot{r} = 0$ so the total energy per unit mass is

$$\begin{aligned}
\frac{E}{m} &= \frac{1}{2p^2} \left(\frac{l}{m} \right)^2 - \frac{GM}{p} \\
&= \frac{1}{2p^2} \left(GM \frac{2ap}{a+p} \right) - \frac{GM}{p} \\
&= GM \left[\frac{2ap}{2p^2(a+p)} - \frac{1}{p} \right] \\
&= GM \left[\frac{2ap}{2p^2(a+p)} - \frac{2p(a+p)}{2p^2(a+p)} \right] \\
&= GM \left[\frac{2ap - 2p(a+p)}{2p^2(a+p)} \right] \\
&= GM \left[\frac{-2p^2}{2p^2(a+p)} \right] \\
&= -GM \left[\frac{1}{a+p} \right] \\
&= -\frac{GM}{2} \left[\frac{2}{a+p} \right] \\
&= -\frac{GM}{2A}
\end{aligned} \tag{8.5}$$

Note that energy is negative because we took zero to be when the particle is at infinity. We can rearrange equation 8.5 to obtain the semi major axis as

$$\begin{aligned}
\frac{E}{m} &= \tilde{E} = -\frac{GM}{2A} \\
2A\tilde{E} &= -GM \\
A &= -\frac{GM}{2\tilde{E}}
\end{aligned} \tag{8.6}$$

Where we are using $\tilde{E} = \frac{E}{m}$ to be the energy per unit mass of the orbiting body. This can then be fed back into our relationship for

angular momentum¹⁹.

$$\begin{aligned}
\tilde{l}^2 &= GMA(1 - e^2) \\
&= -\frac{(GM)^2(1 - e^2)}{2\tilde{E}} \\
-\frac{2\tilde{E}\tilde{l}^2}{(GM)^2} &= 1 - e^2 \\
e^2 - \frac{2\tilde{E}\tilde{l}^2}{(GM)^2} &= 1 \\
e^2 &= 1 + \frac{2\tilde{E}\tilde{l}^2}{(GM)^2} \\
e &= \sqrt{1 + \frac{2\tilde{E}\tilde{l}^2}{(GM)^2}} \tag{8.7}
\end{aligned}$$

Which means we can work out the semi major axis and the eccentricity based on the physics.

8.3 Period

An orbiting body will complete a complete orbit when it has gone through an angle $E = 2\pi$, we can then use Kepler's Equation²⁰ to compute the period. For ease of calculation we will assume that $t_0 = 0$.

$$\begin{aligned}
E - e \sin E &= kt \\
2\pi - e \sin(2\pi) &= kT \\
T &= \frac{2\pi}{k} \tag{8.8}
\end{aligned}$$

¹⁹equation 8.2

²⁰equation 6.15

8.3.1 The Relationship between Mean Motion and Time Period

As we have seen from equation 8.8, the mean motion is related to the orbital period

$$k = \frac{2\pi}{T} \quad (8.9)$$

which is how the constant gets the name *Mean Motion* as it is the mean angular velocity of the orbiting body. In this form Kelper's Equation²¹ becomes

$$E - e \sin E = 2\pi \frac{(t - t_0)}{T} \quad (8.10)$$

8.3.2 Period based on physical and geometric properties

The period can be related to other geometric and physical properties by substituting equation 6.12 in for k , we get

$$T = \frac{2\pi A^2 \sqrt{1 - e^2}}{\frac{l}{m}} \quad (8.11)$$

We can use equation 8.2²² to replace $\frac{l}{m}$.

$$\begin{aligned} T &= \frac{2\pi A^2 \sqrt{1 - e^2}}{\frac{l}{m}} \\ &= \frac{2\pi A^2 \sqrt{1 - e^2}}{\sqrt{GMA(1 - e^2)}} \\ &= \frac{2\pi A^2 \sqrt{1 - e^2}}{\sqrt{GM} \sqrt{A} \sqrt{1 - e^2}} \\ &= \frac{2\pi \sqrt{A^3}}{\sqrt{GM}} \end{aligned} \quad (8.12)$$

²¹equation 6.15 on page 23

²²page 30

We can get this in terms of the apses by using equation 5.4 on page 14 to substitute in for $A = \frac{a+p}{2}$.

$$\begin{aligned}
 T &= \frac{2\pi\sqrt{A^3}}{\sqrt{GM}} \\
 T^2 &= \frac{4\pi^2}{GM} A^3 \\
 &= \frac{4\pi^2}{GM} \left(\frac{a+p}{2}\right)^3
 \end{aligned} \tag{8.13}$$

Note that we have squared T to make the equation easier to work with, but this reproduces one of Kepler's Law, that the square of the period is proportional to the cube of the semi major axis²³.

²³or more usefully the average of the apses

9 In Plane Position and Velocity over time

We will assume that the apses are known as well as the central mass. From this the period is given by equation 8.13 on page 35

$$T = \sqrt{\frac{4\pi^2}{GM}} \left(\frac{a+p}{2} \right)^{\frac{3}{2}}$$

At any given time we can calculate the eccentric anomaly E using Kepler's Equation²⁴

$$E - e \sin E = 2\pi \frac{(t - t_0)}{T}$$

Since this is transcendental, it will need to be solved numerically, or by approximation. This text will not cover those methods, and assume that E can be determined as a function of t .

9.1 Position

Once E is known, then the position can be determined by

$$\begin{aligned} e &= \frac{a-p}{a+p} \\ r &= A(1 - e \cos E) \\ x &= r \cos \theta = A(\cos(E) - e) \\ y &= r \sin \theta = A\sqrt{1-e^2} \sin E \end{aligned}$$

9.2 Velocity

9.2.1 Angular Velocity

Using equation 8.2 on page 30 to get the angular momentum

$$\left(\frac{l}{m} \right) = \sqrt{GM} \sqrt{\frac{2ap}{a+p}}$$

²⁴equation 8.10 on page 34

and using equation ?? on page ??, we can get the angular velocity as

$$\omega = \frac{l}{m} \frac{1}{r^2} = \sqrt{GM} \sqrt{\frac{2ap}{a+p}} \frac{1}{r^2} \quad (9.1)$$

9.2.2 Radial velocity

The radial velocity we obtain from the geometry. Recall that we have

$$r = A(1 - e \cos E) \quad (9.2)$$

As we did for the derivation of Kepler's Equation, we can differentiate with respect to time. Which gives the result

$$\frac{dr}{dt} = eA \sin E \frac{dE}{dt} \quad (9.3)$$

And looking at Kepler's Equation again

$$\begin{aligned} E - e \sin E &= 2\pi \frac{t - t_0}{T} \\ t &= \frac{E - e \sin E}{2\pi} T + t_0 \\ \frac{dt}{dE} &= \frac{1 - e \cos E}{2\pi} T \end{aligned}$$

which gives us the result that

$$\frac{dE}{dt} = \frac{2\pi}{T} \frac{1}{1 - e \cos E} \quad (9.4)$$

We can put this back in to our earlier equation to get the radial velocity

$$\frac{dr}{dt} = \frac{2\pi}{T} \frac{eA \sin E}{1 - e \cos E} \quad (9.5)$$

or in terms of the apses

$$\frac{dr}{dt} = \frac{2\pi}{T} \frac{a - p}{2} \frac{\sin E}{1 - e \cos E} \quad (9.6)$$