

# Assignment 1

AI1110: Probability and Random Variables

Hannanur Rahman

BT22BTECH11007

A person buys a lottery ticket in 50 lotteries, in each of which his chance of winning a prize is  $\left(\frac{1}{100}\right)$ . What is the probability that he will win a prize (a) at least once (b) at least twice ?

**Answer:**

a.  $1 - \left(\frac{99}{100}\right)^{50}$

b.  $1 - \left(\frac{149}{100}\right)\left(\frac{99}{100}\right)^{49}$

**Solution:**  $X \in \{0, 1, 2, 3, \dots, 50\}$  .

We will solve this using CDF.

The trials are Bernoulli trials.

X has a binomial distribution with  $n=50$  and  $p=\frac{1}{100}$

$$\begin{aligned} \therefore q &= 1 - p = 1 - \frac{1}{100} = \frac{99}{100} \\ \therefore \Pr(X = x) &= \binom{n}{x} q^{n-x} p^x = \binom{50}{x} \left(\frac{99}{100}\right)^{50-x} \left(\frac{1}{100}\right)^x \end{aligned} \quad (1)$$

$$\begin{aligned} \text{(a) } \Pr(X \geq 1) &= 1 - F_X(0) = 1 - \Pr(X \leq 0) \\ &= 1 - \Pr(X = 0) \\ &= 1 - \binom{50}{0} \left(\frac{99}{100}\right)^{50} \\ &= 1 - \left(\frac{99}{100}\right)^{50} \end{aligned} \quad (2)$$

$$\begin{aligned} \text{(b) } \Pr(X \geq 2) &= 1 - F_X(1) = 1 - \Pr(X \leq 1) \\ &= 1 - \Pr(X \leq 1) \\ &= 1 - [\Pr(X = 0) + \Pr(X = 1)] \\ &= 1 - \Pr(X = 0) - \Pr(X = 1) \\ &= 1 - \left(\frac{99}{100}\right)^{50} - \frac{1}{2} \left(\frac{99}{100}\right)^{49} \\ &= 1 - \left(\frac{99}{100}\right)^{49} \left[ \frac{99}{100} + \frac{1}{2} \right] \\ &= 1 - \frac{149}{100} \left(\frac{99}{100}\right)^{49} \end{aligned} \quad (3)$$