

8 The Harmonic Oscillator

We want to solve the Schroedinger equation with a quadratic potential

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} + \frac{1}{2}m\omega^2 x^2 \psi = E\psi.$$

You've done this before with ladder operators. Now we do the brute-force approach as a practice for the Hydrogen Atom. By introducing the constants

$$\begin{aligned}\xi &\equiv \sqrt{\frac{m\omega}{\hbar}} x \\ K &\equiv \frac{2E}{\hbar\omega}\end{aligned}$$

the Schroedinger equation simplifies to

$$\frac{d^2\psi}{d\xi^2} = (\xi^2 - K) \psi.$$

The problem now is to look for allowed value of K . Let's begin by looking at the limit $\xi \gg K$. Then, the equation simplifies to

$$\frac{d^2\psi}{d\xi^2} = \xi^2 \psi.$$

This equation is approximately solved by

$$\psi(x) = Ae^{-\xi^2/2} + Be^{\xi^2/2}.$$

The second term blows up as $\xi \rightarrow \infty$, thus is not normalizable and not physical. The first term looks physical. That motivates us to try the following ansatz:

$$\psi(x) = h(\xi)e^{-\xi^2/2}$$

First, let's calculate the derivatives

$$\begin{aligned}\frac{d\psi}{d\xi} &= \left(\frac{dh}{d\xi} - \xi h \right) e^{-\xi^2/2} \\ \frac{d^2\psi}{d\xi^2} &= \left(\frac{d^2h}{d\xi^2} - 2\xi \frac{dh}{d\xi} + (\xi^2 - 1)h \right) e^{-\xi^2/2}\end{aligned}$$

So the Schroedinger equation from above becomes

$$\frac{d^2h}{d\xi^2} - 2\xi \frac{dh}{d\xi} + (K - 1)h = 0$$

Let us use yet another ansatz, this time in the form of a power series:

$$h(\xi) = \sum_{j=0}^{\infty} a_j \xi^j$$

The derivatives are:

$$\begin{aligned}\frac{dh}{d\xi} &= \sum_{j=1}^{\infty} j a_j \xi^{j-1} \\ \frac{d^2h}{d\xi^2} &= \sum_{j=2}^{\infty} j(j-1) a_j \xi^{j-2}\end{aligned}$$

Note that there is a typo in Griffiths. Putting everything back into the Schroedinger equation gives us

$$\sum_{j=0}^{\infty} [(j+1)(j+2)a_{j+2} - 2ja_j + (K-1)a_j] \xi^j = 0$$

Because of the uniqueness of the power series expansion, we can get rid of the sum and look at each term individually

$$(j+1)(j+2)a_{j+2} - 2ja_j + (K-1)a_j = 0$$

or equivalently

$$a_{j+2} = \frac{2j+1-K}{(j+1)(j+2)} a_j$$

This is still the Schroedinger equation! Once we specify a_0 all the even coefficients are determined. Once we specify a_1 all the odd coefficients are determined. Thus, let's write

$$\begin{aligned} h(\xi) &= h_{\text{even}}(\xi) + h_{\text{odd}}(\xi), & \text{where} \\ h_{\text{even}}(\xi) &= a_0 + a_2\xi^2 + \dots \\ h_{\text{odd}}(\xi) &= a_1\xi + a_3\xi^3 + \dots \end{aligned}$$

Remember, solutions have to be normalizable. So let's look at the limit for large j . We have $a_{j+2} \approx \frac{2}{j}a_j$, or up to some constant $a_j \approx \frac{C}{(j/2)!}$. Thus:

$$\begin{aligned} h(\xi) &\approx C \sum \frac{j}{(j/2)!} \xi^j \\ &\approx C \sum \frac{1}{j!} \xi^{2j} \approx C e^{\xi^2} \end{aligned}$$

Note that that would make the function ψ not normalizable. So something else has to be going on. We need to make sure the series of a 's will terminate. Thus, there is an $a_n = 0$. This ends the series of either the odd or the even part of h . The other one has to be 0 from the beginning ($a_1 = 0$ or $a_0 = 0$ depending on whether n is even or odd). Thus:

$$K = 2n + 1$$

Remember what K is to get

$$E_n = \left(n + \frac{1}{2}\right) \hbar \omega \quad \text{for } n = 0, 1, \dots$$

Using the allowed values for K , the recursion formula reads

$$a_{j+2} = \frac{-2(n-j)}{(j+1)(j+2)} a_j$$

Suppose $n = 0$, then there is only one term in the series (we have to set $a_1 = 0$ to kill h_{odd}). Thus:

$$h_0(\xi) = a_0$$

and hence

$$\psi_0(\xi) = a_0 e^{-\xi^2/2}$$

Similarly for $n = 1$, we take $a_0 = 0$, giving us

$$h_1(\xi) = a_1\xi$$

and hence

$$\psi_1(\xi) = a_1\xi e^{-\xi^2/2}$$

Now let's look at $n = 2$. This yields

$$\begin{aligned} a_2 &= -2a_0 \\ a_4 &= 0 \end{aligned}$$

hence

$$h_2(\xi) = a_0(1 - 2\xi^2)$$

and

$$\psi_2(\xi) = a_0(1 - 2\xi^2)e^{-\xi^2/2}$$

And so on. Probably time to stop with the algebra. Apart from the overall normalization, the solutions involve Hermite polynomials $H_n(\xi)$. The wavefunction has the general form of

$$\Phi_n(x) = \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} \frac{1}{\sqrt{2^n n!}} H_n(\xi) e^{-\xi^2/2}$$

where

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2} = \left(2x - \frac{d}{dx}\right)^n \cdot 1$$

The first part is called Rodrigues formula.

9 Quantum Mechanics in Three dimensions

9.1 Schroedinger Equation in Three dimensions

The time dependent Schroedinger equation is

$$i\hbar \frac{\partial \Phi}{\partial t} = H\Phi$$

Written in this way, this equation can already be interpreted as the three dimensional Schroedinger equation. We write the Hamiltonian as

$$H = \frac{1}{2m} (p_x^2 + p_y^2 + p_z^2) + V(x, y, z)$$

Using the standard definitions for the momentum operators we get

$$\begin{aligned} H &= \frac{-\hbar^2}{2m} (\partial_x^2 + \partial_y^2 + \partial_z^2) + V(x, y, z) \\ &= \frac{-\hbar^2}{2m} \nabla^2 + V(x, y, z). \end{aligned}$$

The operator ∇ is called the Laplacian, or `\nabla` in L^AT_EX. We again normalize the wave function for physical states. The integral is now a three dimensional integral:

$$\int |\Phi| d^3r = 1.$$

If the Hamiltonian is independent of time, we can factor of the time dependent exponential just as in the one dimensional case to get the three dimensional time independent Schroedinger Equation

$$\begin{aligned} H\psi &= E\psi \\ -\frac{\hbar^2}{2m}\nabla^2\psi + V\psi &= E\psi \end{aligned}$$

The time dependent solution is then

$$\Phi(x, y, z, t) = \sum c_n \Psi_n(x, y, z) e^{-iE_n t/\hbar}$$

9.2 Separation of variables

In most cases the potential is just a function of the radius

$$V(x, y, z) = V(r)$$

This motivates the introduction of spherical coordinates (r, θ, ϕ) . In spherical coordinates the Laplacian takes the following form

$$\nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \left(\frac{\partial^2}{\partial \phi^2} \right)$$

So let's look for solutions to the Schroedinger Equation in the form of

$$\psi(r, \theta, \phi) = R(r)Y(\theta, \phi)$$

Let's put this Ansatz in the Schroedinger Equation

$$-\frac{\hbar^2}{2m} \left(\frac{Y}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} R \right) + \frac{R}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} Y \right) + \frac{R}{r^2 \sin^2 \theta} \left(\frac{\partial^2}{\partial \phi^2} Y \right) \right) + VRY = ERY$$

Multiplying with $-2mr^2/(\hbar^2 RY)$ gives:

$$\frac{1}{R} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} R \right) + \frac{1}{Y \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} Y \right) + \frac{1}{Y \sin^2 \theta} \left(\frac{\partial^2}{\partial \phi^2} Y \right) - \frac{2mr^2}{\hbar^2} (V - E) = 0$$

Our Ansatz worked! Note that only the first and last term depend on R . Only the middle two terms depend on ϕ and θ . Therefore, the two terms have to be constants to cancel each other. The constant can be any complex number. W.l.o.g. let us call this constant $l(l+1)$ for reasons that will become obvious in a second. We now have two equations

$$\begin{aligned} \frac{1}{R} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} R \right) - \frac{2mr^2}{\hbar^2} (V - E) &= l(l+1) \\ \frac{1}{Y \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} Y \right) + \frac{1}{Y \sin^2 \theta} \left(\frac{\partial^2}{\partial \phi^2} Y \right) &= -l(l+1) \end{aligned}$$

Let's look at both of those equations individually.

9.3 The Angular Equation

The equation we are trying to solve is

$$\frac{1}{Y \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} Y \right) + \frac{1}{Y \sin^2 \theta} \left(\frac{\partial^2}{\partial \phi^2} Y \right) = -l(l+1)$$

or equivalently

$$\sin \theta \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} Y \right) + \left(\frac{\partial^2}{\partial \phi^2} Y \right) = -l(l+1)Y \sin^2 \theta$$

Let's try an Ansatz as follow

$$Y(\theta, \phi) = \Theta(\theta)\Phi(\phi)$$

This gives (after dividing by $\Theta\Phi$)

$$\frac{1}{\Theta} \sin \theta \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \Theta \right) + \frac{1}{\Phi} \left(\frac{\partial^2}{\partial \phi^2} \Phi \right) = -l(l+1) \sin^2 \theta$$

In the same way we defined the constant $l(l+1)$ in the radial equation, we can again see that this equation is only satisfied if we have a constant m^2 (arbitrary name) such that

$$\begin{aligned} \frac{1}{\Theta} \sin \theta \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \Theta \right) + l(l+1) \sin^2 \theta &= m^2 \\ \frac{1}{\Phi} \left(\frac{\partial^2}{\partial \phi^2} \Phi \right) &= -m^2 \end{aligned}$$

The last equation (the one for Φ) is easy to solve:

$$\Phi(\phi) = e^{im\phi}$$

Because of symmetry, we must have

$$\Phi(\phi + 2\pi) = \Phi(\phi)$$

It follows that only certain values of m are allowed

$$m = 0, \pm 1, \pm 2, \dots$$

Let's look at the other equation

$$\sin \theta \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \Theta \right) + \Theta (l(l+1) \sin^2 \theta - m^2) = 0$$

This is harder to solve, so I'm giving you the answer:

$$\Theta(\theta) = AP_l^m(\cos \theta)$$

The P_l^m 's are the associated Legendre functions, defined by

$$P_l^m(x) \equiv (1-x^2)^{|m|/2} \left(\frac{\partial}{\partial x} \right)^{|m|} P_l(x)$$

where P_l are the Legendre polynomials

$$P_l(x) \equiv \frac{1}{2^l l!} \left(\frac{\partial}{\partial x} \right)^l (x^2 - 1)^l$$

The first few examples are

$$\begin{aligned}P_0(x) &= 1 \\P_1(x) &= x \\P_3(x) &= \frac{3}{2}x^2 - \frac{1}{2}\end{aligned}$$

The associated Legendre polynomials are in general not a polynomial. Here are a few examples:

$$\begin{aligned}P_0^0 &= 1 \\P_1^0 &= x \\P_1^1 &= -\sqrt{1-x^2} \\P_1^{-1} &= \frac{1}{2}\sqrt{1-x^2}\end{aligned}$$

LECTURE 6 ENDS