

ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/lagb20>

Support τ -tilting modules over one-point extensions

Hanpeng Gao & Zongzhen Xie

To cite this article: Hanpeng Gao & Zongzhen Xie (2021) Support τ -tilting modules over one-point extensions, Communications in Algebra, 49:2, 739-746, DOI: [10.1080/00927872.2020.1817470](https://doi.org/10.1080/00927872.2020.1817470)

To link to this article: <https://doi.org/10.1080/00927872.2020.1817470>



Published online: 17 Sep 2020.



Submit your article to this journal [↗](#)



Article views: 58



View related articles [↗](#)



View Crossmark data [↗](#)



Support τ -tilting modules over one-point extensions

Hanpeng Gao^a and Zongzhen Xie^b

^aDepartment of Mathematics, Nanjing University, Nanjing, P.R. China; ^bDepartment of Mathematics and Computer Science, School of Biomedical Engineering and Informatics, Nanjing Medical University, Nanjing, P.R. China

ABSTRACT

Let B be the one-point extension algebra of A by an A -module X . We proved that every support τ -tilting A -module can be extended to be a support τ -tilting B -module by two different ways. As a consequence, it is shown that there is an inequality

$$|\text{supp } \tau\text{-tilt } B| \geq 2|\text{supp } \tau\text{-tilt } A|.$$

ARTICLE HISTORY

Received 19 May 2020
Revised 24 August 2020
Communicated by Scott Chapman

KEYWORDS

One-point extensions;
semibricks; support
 τ -tilting modules

2020 MATHEMATICS

SUBJECT

CLASSIFICATION

16E30; 16G20

1. Introduction

Tilting modules are very important in the representation theory of finite dimensional algebras. Mutation is an effective way to construct a new tilting module from a given one. Unfortunately, mutation of tilting modules may not be realized.

In 2014, Adachi et al. [1] introduced the concept of support τ -tilting modules as a generalization of tilting modules, and they showed that mutation of support τ -tilting modules is always possible. The authors also proved that support τ -tilting modules are in bijection with some important classes in representation theory (such as, functorially finite torsion classes, 2-term sifting complexes, and cluster-tilting objects in the cluster category).

A new (support τ)-tilting module can be constructed by algebra extensions. In [3], Assem et al. studied how to extend and restrict tilting modules for one-point extension algebras by a projective module. In [8], Suarez generalized this result for the context of support support τ -tilting modules. More precisely, let $B = A[P]$ be the one-point extension of an algebra A by a projective A -module P and e the identity of A . If M is a support τ -tilting A -module, then $\text{Hom}_B(eB, M) \oplus S_a$ is a support τ -tilting B -module, where S_a is the simple module corresponding to the new point a (see [8, Theorem A]). An example shown that $\text{Hom}_B(eB, M) \oplus S_a$ may not be a support τ -tilting B -module if P is not projective (see [8, Example 4.7]).

Bricks and semibricks are considered in [5, 6]. An A -module M is called *brick* if $\text{Hom}_A(M, M)$ is a k division. A *semibrick* is a set consisting of isoclasses of pairwise Hom-orthogonal bricks. Let \mathcal{S} be a semibrick and $T(\mathcal{S})$ the smallest torsion class containing \mathcal{S} . In [2], the author called a semibrick \mathcal{S} is *left finite* if $T(\mathcal{S})$ is functorially finite and he also proved that there exists a

bijection $\Phi : \text{s}\tau\text{-tilt } A \mapsto \text{f}_L - \text{sbrick } A$ between the set of support τ -tilting A -modules and the set of left finite semibricks of A .

In this paper, we construct semibricks over the one-point extension B of an algebra A by an A -module X (may not be projective) and use the bijection to get support τ -tilting B -modules.

Proposition 1.1. (see Proposition 3.2) *Let B be the one-point extension algebra of A by an A -module X and \mathcal{S} be a semibrick in $\text{mod } A$. Then both \mathcal{S} and $\mathcal{S} \cup S_a$ are semibricks in $\text{mod } B$, where S_a stands for the simple module corresponding to the extension point a .*

Moreover, it is shown that \mathcal{S} is left finite implies $\mathcal{S} \cup S_a$ is also. We say an A -module M is a support τ -tilting module with respect to the semibrick \mathcal{S} if $\Phi(M) = \mathcal{S}$. As an application, we can construct support τ -tilting modules over one-point extensions from support τ -tilting A -modules.

Proposition 1.2. (see Proposition 3.7) *Let B be the one-point extension algebra of A by an A -module X and $M \in \text{mod } A$ be a support τ -tilting module with respect to the semibrick \mathcal{S} . Then both $P(T(\mathcal{S}))$ and $P(T(\mathcal{S} \cup S_a))$ are support τ -tilting B -modules.*

As a consequence, we have the following inequality

Corollary 1.3. $|\text{s}\tau\text{-tilt } B| \geq 2|\text{s}\tau\text{-tilt } A|$.

Moreover, we have the following.

Theorem 1.4. (see Theorem 3.9) *Let B be the one-point extension algebra of A by an A -module X and M be a support τ -tilting module in $\text{mod } A$.*

- (1) M is a support τ -tilting B -module.
- (2) Assume that $M \in \text{mod } A$ is a support τ -tilting module with respect to the semibrick \mathcal{S} , then $P(T(\mathcal{S} \cup S_a))$ has M as direct summand.
- (3) If $X \in \text{Fac } M$, then $P_a \oplus M$ is a support τ -tilting B -module.
- (4) If $\text{Hom}_A(X, \text{Fac } M) = 0$, then $S_a \oplus M$ is a support τ -tilting B -module.

Throughout this paper, all algebras will be basic connected finite dimensional k -algebras over an algebraically closed field k and all modules are basic. Let A be an algebra. The category of finitely generated left A -modules will be denote by $\text{mod } A$ and the Auslander-Reiten translation of A will be denote by τ . For $M \in \text{mod } A$, we denote by $\text{ind}(M)$ the set of isoclasses of indecomposable direct summands of M , and by $\text{Fac } M$ the full subcategory of $\text{mod } A$ consisting of modules isomorphic to factor modules of finite direct sums of copies of M . For a finite set J , $|J|$ stands for the cardinality of J . In particular, we write $|M| = |\text{ind}(M)|$. \mathbb{N} will be the set of all natural numbers.

2. Preliminaries

Let A be an algebra. In this section, we recall some definitions about support τ -tilting modules and semibricks over $\text{mod } A$.

Definition 2.1. ([1, Definition 0.1]) Let $M \in \text{mod } A$.

- (1) M is called τ -rigid if $\text{Hom}_A(M, \tau M) = 0$.
- (2) M is called τ -tilting if it is τ -rigid and $|M| = |A|$.
- (3) M is called support τ -tilting if it is a τ -tilting $A/\langle e \rangle$ -module where e is an idempotent of A .

We will denote by $\tau\text{-tilt } A$ (respectively, $\text{s}\tau\text{-tilt } A$) the set of isomorphism classes of τ -tilting A -modules (respectively, support τ -tilting A -modules).

Definition 2.2. ([1, Definition 0.3]) Let (M, P) be a pair in $\text{mod } A$ with P projective.

- (1) The pair (M, P) is called a τ -rigid pair if M is τ -rigid and $\text{Hom}_A(P, M) = 0$.
- (2) The pair (M, P) is called a support τ -tilting pair if it is τ -rigid and $|M| + |P| = |A|$.

Note that (M, P) is a support τ -tilting pair if and only if M is a τ -tilting $A/\langle e \rangle$ -module, where $eA \cong P$ [1, Proposition 2.3]. Hence, M is a τ -tilting A -module if and only if $(M, 0)$ is a support τ -tilting pair.

The following result is very useful.

Lemma 2.3. ([4, Proposition 5.8]) *For $M \in \text{mod } A$, M is τ -rigid if and only if $\text{Ext}_A^1(M, \text{Fac} M) = 0$.*

Definition 2.4. ([2, Definition 2.1]) Let $\mathcal{S} \subseteq \text{mod } A$. \mathcal{S} is called a *semibrick* if

$$\text{Hom}_A(S_i, S_j) = \begin{cases} k\text{-division algebra} & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

for any $S_i, S_j \in \mathcal{S}$.

By Schur's Lemma, a set of isoclasses of some simple modules is a semibrick.

Let \mathcal{Y} be a full subcategory of $\text{mod } A$ and $M \in \text{mod } A$. A homomorphism $f_M : M \rightarrow Y_M$ is called left \mathcal{Y} -approximation of M with $Y_M \in \mathcal{Y}$ if any morphism $f : M \rightarrow Y$ with $Y \in \mathcal{Y}$ factors through f_M . We say that \mathcal{Y} is *covariantly finite* if for any $M \in \text{mod } A$, there exists a left \mathcal{Y} -approximation of M . Dually, we can define the concepts of right \mathcal{Y} -approximation of M and *contravariantly finite* subcategories. \mathcal{Y} is called *functorially finite* if it is both covariantly finite and contravariantly finite.

A *torsion class* of $\text{mod } A$ is a full subcategory of $\text{mod } A$ closed under images, direct sums, and extensions. Recall that a semibrick \mathcal{S} of $\text{mod } A$ is *left finite* [2] if $T(\mathcal{S})$ is functorially finite, where $T(\mathcal{S})$ is the smallest torsion class containing \mathcal{S} . The set of all left finite semibricks of $\text{mod } A$ will be denoted by $\text{f}_L\text{-sbrick } A$.

The following result states the relationship between $\text{st-tilt } A$ and $\text{f}_L\text{-sbrick } A$.

Theorem 2.5. [2, Theorem 1.3(2)] *there exists a bijection*

$$\Phi : \text{st-tilt } A \mapsto \text{f}_L\text{-sbrick } A$$

given by $M \mapsto \text{ind}(M/\text{rad}_\Gamma M)$ where $\Gamma = \text{End}_A(M)$.

Recall that $M \in \text{mod } A$ is called *sincere* if every simple A -module appears as a composition factor in M . A τ -tilting A -module is exactly a sincere support τ -tilting. We say a semibrick \mathcal{S} of $\text{mod } A$ is *sincere* if $T(\mathcal{S})$ is sincere. Let $\text{sf}_L\text{-sbrick } A$ stand for all sincere left finite semibricks of $\text{mod } A$. We have the following result due to Asai in [2].

Corollary 2.6. *There exists a bijection $\Phi : \text{st-tilt } A \mapsto \text{sf}_L\text{-sbrick } A$.*

3. Main results

Let $X \in \text{mod } A$. The *one-point extension* of A by X is defined as the following matrix algebra

$$B = \begin{pmatrix} A & X \\ 0 & k \end{pmatrix}$$

with the ordinary matrix addition and the multiplication induced by the module structure of X . We write $B := A[X]$ with a the extension point. All B -modules can be viewed as $\begin{pmatrix} M \\ k^n \end{pmatrix}_f$ where $M \in \text{mod } A$, $n \in \mathbb{N}$ and $f \in \text{Hom}_A(X \otimes_k k^n, M)$ (see, [7, XV.1]). In particular, $S_a = \begin{pmatrix} 0 \\ k \end{pmatrix}_0$ and $P_a = \begin{pmatrix} X \\ k \end{pmatrix}_1$. Moreover, the morphisms from $\begin{pmatrix} M \\ k^n \end{pmatrix}_f$ to $\begin{pmatrix} M' \\ k^{n'} \end{pmatrix}_{f'}$ are pairs of $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ such that the following diagram

$$\begin{array}{ccc}
 X \otimes_k k^n & \xrightarrow{f} & M \\
 X \otimes \beta \downarrow & & \downarrow \alpha \\
 X \otimes_k k^{n'} & \xrightarrow{f'} & M'
 \end{array}$$

commutes, where $\alpha \in \text{Hom}_\Lambda(M, M')$ and $\beta \in \text{Hom}_\Gamma(k^n, k^{n'})$. A sequence

$$0 \rightarrow \begin{pmatrix} M_1 \\ k^{n_1} \end{pmatrix}_{f_1} \xrightarrow{\begin{pmatrix} \alpha_1 \\ \beta_1 \end{pmatrix}} \begin{pmatrix} M_2 \\ k^{n_2} \end{pmatrix}_{f_2} \xrightarrow{\begin{pmatrix} \alpha_2 \\ \beta_2 \end{pmatrix}} \begin{pmatrix} M_3 \\ k^{n_3} \end{pmatrix}_{f_3} \rightarrow 0$$

in $\text{mod } B$ is exact if and only if

$$0 \rightarrow M_1 \xrightarrow{\alpha_1} M_2 \xrightarrow{\alpha_2} M_3 \rightarrow 0$$

is exact in $\text{mod } A$ and

$$0 \rightarrow k^{n_1} \xrightarrow{\beta_1} k^{n_2} \xrightarrow{\beta_2} k^{n_3} \rightarrow 0$$

is exact in $\text{mod } k$.

Lemma 3.1. *For any $M \in \text{mod } A$, we have*

- (1) $\text{Hom}_B(S_a, M) = 0$;
- (2) $\text{Hom}_B(M, S_a) = 0$.

Proof. It is clear since ${}_B M \cong \begin{pmatrix} M \\ 0 \end{pmatrix}_0$. Hence, $\text{Hom}_B(S_a, M) \cong \text{Hom}_B\left(\begin{pmatrix} 0 \\ k \end{pmatrix}_0, \begin{pmatrix} M \\ 0 \end{pmatrix}_0\right) = 0$. Similarly, we can get $\text{Hom}_B(M, S_a) = 0$. □

Proposition 3.2. *Let \mathcal{S} be a semibrick in $\text{mod } A$. Then both \mathcal{S} and $\mathcal{S} \cup S_a$ are semibricks in $\text{mod } B$.*

Proof. It follows from Lemma 3.1. □

Lemma 3.3. *Let \mathcal{S} be a semibrick in $\text{mod } A$. Then*

$$T(\mathcal{S} \cup S_a) = \left\{ \begin{pmatrix} M \\ k^n \end{pmatrix}_f \mid \forall n \in \mathbb{N}, M \in T(\mathcal{S}) \text{ and } f \in \text{Hom}_A(X \otimes_k k^n, M) \right\}.$$

Proof. Since \mathcal{S} and S_a belong to $T(\mathcal{S} \cup S_a)$, we have $\left\{ \begin{pmatrix} M \\ 0 \end{pmatrix}_0 \mid M \in T(\mathcal{S}) \right\} \subset T(\mathcal{S} \cup S_a)$ and $\begin{pmatrix} 0 \\ k^n \end{pmatrix} \in T(\mathcal{S} \cup S_a)$ for all $n \in \mathbb{N}$. Note that $\forall n \in \mathbb{N}, M \in T(\mathcal{S})$ and $f \in \text{Hom}_A(X \otimes_k k^n, M)$, there exists the following exact sequence in $\text{mod } B$

$$0 \rightarrow \begin{pmatrix} M \\ 0 \end{pmatrix}_0 \rightarrow \begin{pmatrix} M \\ k^n \end{pmatrix}_f \rightarrow \begin{pmatrix} 0 \\ k^n \end{pmatrix}_0 \rightarrow 0$$

this implies $\begin{pmatrix} M \\ k^n \end{pmatrix}_f \in T(\mathcal{S} \cup S_a)$. It is clear that $\left\{ \begin{pmatrix} M \\ k^n \end{pmatrix}_f \mid \forall n \in \mathbb{N}, M \in T(\mathcal{S}) \text{ and } f \in \text{Hom}_A(X \otimes_k k^n, M) \right\}$ is closed under image, direct sum and extension. Thus it is a torsion class.

Hence $T(\mathcal{S} \cup S_a) = \left\{ \begin{pmatrix} M \\ k^n \end{pmatrix}_f \mid \forall n \in \mathbb{N}, M \in T(\mathcal{S}) \text{ and } f \in \text{Hom}_A(X \otimes_k k^n, M) \right\}$. □

Proposition 3.4. *Let \mathcal{S} be a semibrick in $\text{mod } A$. If \mathcal{S} is left finite, then $\mathcal{S} \cup \mathcal{S}_a$ is also.*

Proof. We only show that $T(\mathcal{S} \cup \mathcal{S}_a)$ is covariantly finite. It is dually to prove $T(\mathcal{S} \cup \mathcal{S}_a)$ is contravariantly finite.

Let $\begin{pmatrix} M \\ k^n \end{pmatrix}_f \in \text{mod } B$. Then M has a left $T(\mathcal{S})$ -approximation $h_M : M \rightarrow Z_M$ in $\text{mod } A$ since $T(\mathcal{S})$ is covariantly finite. Take $g = h_M \circ f$. The following commutative diagram

$$\begin{array}{ccc} X \otimes_k k^n & \xrightarrow{f} & M \\ \parallel & & \downarrow h_M \\ X \otimes_k k^n & \xrightarrow{g} & Z_M \end{array}$$

implies that $\begin{pmatrix} f_M \\ 1 \end{pmatrix}$ is a morphism from $\begin{pmatrix} M \\ k^n \end{pmatrix}_f$ to $\begin{pmatrix} Z_M \\ k^n \end{pmatrix}_g$. Next, we will show that $\begin{pmatrix} f_M \\ 1 \end{pmatrix}$ is left $T(\mathcal{S} \cup \mathcal{S}_a)$ -approximation of $\begin{pmatrix} M \\ k^n \end{pmatrix}_f$. For any $\begin{pmatrix} M_1 \\ k^{n_1} \end{pmatrix}_{f_1} \in T(\mathcal{S} \cup \mathcal{S}_a)$ and morphism $\begin{pmatrix} a \\ b \end{pmatrix} : \begin{pmatrix} M \\ k^n \end{pmatrix}_f \rightarrow \begin{pmatrix} M_1 \\ k^{n_1} \end{pmatrix}_{f_1}$, there is a morphism $h' : Z_M \rightarrow M_1$ such that $a = h' \circ h_M$ since h_M is a left approximation. Note that there exists a commutative diagram

$$\begin{array}{ccc} X \otimes_k k^n & \xrightarrow{f} & M \\ X \otimes b \downarrow & & \downarrow a \\ X \otimes_k k^{n_1} & \xrightarrow{f_1} & M_1 \end{array}$$

that is $a \circ f = f_1 \circ (X \otimes b)$. Therefore,

$$f_1 \circ (X \otimes b) = a \circ f = h' \circ h_M \circ f = h' \circ g,$$

that is, the following diagram

$$\begin{array}{ccc} X \otimes_k k^n & \xrightarrow{g} & Z_M \\ X \otimes b \downarrow & & \downarrow h' \\ X \otimes_k k^{n_1} & \xrightarrow{f_1} & M_1 \end{array}$$

commutes. Hence, $\begin{pmatrix} h' \\ b \end{pmatrix}$ is a morphism from $\begin{pmatrix} Z_M \\ k^n \end{pmatrix}_g$ to $\begin{pmatrix} M_1 \\ k^{n_1} \end{pmatrix}_{f_1}$, and the following equation holds

$$\begin{pmatrix} h' \\ b \end{pmatrix} \circ \begin{pmatrix} h_M \\ 1 \end{pmatrix} = \begin{pmatrix} h' \circ h_M \\ b \end{pmatrix} = \begin{pmatrix} g \\ b \end{pmatrix}.$$

By Lemma 3.3, $\begin{pmatrix} Z_M \\ k^n \end{pmatrix}_g \in T(\mathcal{S} \cup \mathcal{S}_a)$ since $Z_M \in T(\mathcal{S})$. Thus, we were done. \square

The following result can be found immediately.

Corollary 3.5. *Let \mathcal{S} be a semibrick in $\text{mod } A$. If \mathcal{S} is sincere left finite, then $\mathcal{S} \cup \mathcal{S}_a$ is also.*

Let \mathcal{F} be a full subcategory of $\text{mod } A$. An A -module M is called **Ext-projective** in \mathcal{F} if $\text{Ext}_A^1(M, F) = 0$ for all $F \in \mathcal{F}$. If \mathcal{F} is functorially finite in $\text{mod } A$, then there are only finitely many indecomposable Ext-projective modules in \mathcal{F} up to isomorphism. In this case, we will denote by $P(\mathcal{F})$ the direct sum of all Ext-projective modules in \mathcal{F} up to isomorphism.

Definition 3.6. We say that an A -module M is a support τ -tilting module with respect to the semibrick \mathcal{S} if $\Phi(M) = \mathcal{S}$.

Now, we can construct support τ -tilting B -modules from support τ -tilting A -modules.

Proposition 3.7. *Let $M \in \text{mod } A$ be a support τ -tilting module with respect to the semibrick \mathcal{S} . Then both $P(T(\mathcal{S}))$ and $P(T(\mathcal{S} \cup \mathcal{S}_a))$ are support τ -tilting B -modules. Moreover, if M is τ -tilting, then $P(T(\mathcal{S} \cup \mathcal{S}_a))$ is also.*

Proof. Since M is a support τ -tilting module, we have \mathcal{S} is a left finite semibrick of $\text{mod } A$ by Theorem 2.5. Hence, \mathcal{S} is also a left finite semibrick of $\text{mod } B$. Moreover, we have $\mathcal{S} \cup \mathcal{S}_a$ is a left finite semibrick of $\text{mod } B$ by Proposition 3.4. Therefore, $T(\mathcal{S})$ and $T(\mathcal{S} \cup \mathcal{S}_a)$ are functorially finite torsion classes. Hence, $P(T(\mathcal{S}))$ and $P(T(\mathcal{S} \cup \mathcal{S}_a))$ are support τ -tilting B -module by [1, Theorem 2.7]). \square

As a consequence, we have the following inequality.

Corollary 3.8. $|\text{s}\tau\text{-tilt } B| \geq 2|\text{s}\tau\text{-tilt } A|$.

Applying Proposition 3.7, we can give those forms of support τ -tilting B -module under certain conditions.

Theorem 3.9. *Let B be the one-point extension of A by X and M be a support τ -tilting module in $\text{mod } A$.*

- (1) M is a support τ -tilting B -module.
- (2) Assume that $M \in \text{mod } A$ is a support τ -tilting module with respect to the semibrick \mathcal{S} , then $P(T(\mathcal{S} \cup \mathcal{S}_a))$ has M as direct summand.
- (3) If $X \in \text{Fac } M$, then $P_a \oplus M$ is a support τ -tilting B -module.
- (4) If $\text{Hom}_A(X, \text{Fac } M) = 0$, then $S_a \oplus M$ is a support τ -tilting B -module.

Proof. Assume that $M \in \text{mod } A$ is a support τ -tilting module with respect to the semibrick \mathcal{S} , then $T(\mathcal{S}) = \text{Fac } M$ [2, Lemma 2.5(5)].

Note that $\forall n \in \mathbb{N}$, $M' \in T(\mathcal{S})$ and $f \in \text{Hom}_A(X \otimes_k k^n, M')$, there exists the following exact sequence in $\text{mod } B$

$$0 \rightarrow \begin{pmatrix} M' \\ 0 \end{pmatrix}_0 \rightarrow \begin{pmatrix} M' \\ k^n \end{pmatrix}_f \rightarrow \begin{pmatrix} 0 \\ k^n \end{pmatrix}_0 \rightarrow 0. \quad (1.1)$$

For any $Y \in \text{mod } B$, applying the functor $\text{Hom}_B(Y, -)$ to (1.1), we have the following exact sequence

$$\text{Ext}_B^1\left(Y, \begin{pmatrix} M' \\ 0 \end{pmatrix}_0\right) \rightarrow \text{Ext}_B^1\left(Y, \begin{pmatrix} M' \\ k^n \end{pmatrix}_f\right) \rightarrow \text{Ext}_B^1\left(Y, \begin{pmatrix} 0 \\ k^n \end{pmatrix}_0\right) = 0. \quad (1.2)$$

- (1) By Proposition 3.7, $P(T(\mathcal{S})) = P(\text{Fac } M) = M$ is a support τ -tilting B -module.
- (2) Putting $Y = {}_B M \cong \begin{pmatrix} M \\ 0 \end{pmatrix}_0$ in (1.2), we have $\text{Ext}_B^1\left(M, \begin{pmatrix} M' \\ 0 \end{pmatrix}_0\right) \cong \text{Ext}_A^1(M, M') = 0$ by Lemma 2.3, and hence $\text{Ext}_B^1\left(M, \begin{pmatrix} M' \\ k^n \end{pmatrix}_f\right) = 0$. By Lemma 3.3, M is a Ext-projective object in $T(\mathcal{S} \cup \mathcal{S}_a)$. Therefore, $P(T(\mathcal{S} \cup \mathcal{S}_a))$ has M as direct summand.
- (3) If $X \in \text{Fac } M$, then $P_a \in T(\mathcal{S} \cup \mathcal{S}_a)$ by Lemma 3.3. Hence, $P_a \oplus M$ is a direct summand of $P(T(\mathcal{S} \cup \mathcal{S}_a))$ by (2). In particular, $P_a \oplus M$ is a τ -rigid B -module. Suppose that (M, P) is a support τ -tilting pair in $\text{mod } A$. Hence, $\text{Hom}_A(P, \text{Fac } M) = 0$ because $\text{Hom}_A(P, M) = 0$.

This implies $\text{Hom}_B(P, P_a) \cong \text{Hom}_A(P, X) = 0$. Therefore, $(P_a \oplus M, P)$ is a support τ -tilting pair in $\text{mod } B$ since $|P_a \oplus M| + |P| = 1 + |A| = |B|$.

(4) Note that there is an exact sequence in $\text{mod } B$,

$$0 \rightarrow \begin{pmatrix} X \\ 0 \end{pmatrix} \cong X \xrightarrow{f} P_a \rightarrow S_a \rightarrow 0.$$

For any $Y' \in \text{Fac } M$, applying $\text{Hom}_B(-, Y')$ to it, we have the following exact sequence,

$$\text{Hom}_B(X, Y') \rightarrow \text{Ext}_B^1(S_a, Y') \rightarrow \text{Ext}_B^1(P_a, Y') = 0.$$

Since $\text{Hom}_A(X, \text{Fac } M) = 0$, we have $\text{Hom}_B(X, Y') = 0$. Hence, $\text{Ext}_B^1(S_a, Y') = 0$. Thus, $\text{Ext}_B^1(S_a, \text{Fac } M) = 0$. Putting $Y = S_a$ in (1.2), we have $\text{Ext}_B^1\left(S_a, \begin{pmatrix} M' \\ k^n \end{pmatrix}_f\right) = 0$. By Lemma 3.3, S_a is a Ext-projective object in $T(\mathcal{S} \cup S_a)$. Therefore, $P(T(\mathcal{S} \cup S_a))$ has $S_a \oplus M$ as direct summand. This implies $S_a \oplus M$ is a τ -rigid B -module. Suppose that (M, P) is a support τ -tilting pair in $\text{mod } A$. It is easy to get $(S_a \oplus M, P)$ is a support τ -tilting pair in $\text{mod } B$ since $\text{Hom}_B(P, S_a) = 0$ and $|S_a \oplus M| + |P| = 1 + |A| = |B|$. Hence, $S_a \oplus M$ is a support τ -tilting B -module. \square

Corollary 3.10. *Let B be the one-point extension of A by X and $M \in \text{mod } A$ be a τ -tilting module.*

- (1) *If $X \in \text{Fac } M$, then $P_a \oplus M$ is a τ -tilting B -module.*
- (2) *If $\text{Hom}_A(X, \text{Fac } M) = 0$, then $S_a \oplus M$ is a τ -tilting B -module.*

Example 3.11. Let A be a finite dimensional k -algebra given by the quiver

$$2 \rightarrow 3.$$

Considering the one-point extension of A by the simple module corresponding to the point 2, the algebra $B = A[2]$ is given by the quiver


$$1 \xrightarrow{\alpha} 2 \xrightarrow{\beta} 3$$

with the relation $\alpha\beta = 0$. The Hasse quiver of A is as follows (semibricks be remarked by red).

$$\begin{array}{ccccc} Q(\text{st-tilt } A): & T_1 = \begin{smallmatrix} 2 \\ 3 \end{smallmatrix} \text{ } 3 & \longrightarrow & T_2 = \text{ } 3 & \longrightarrow & T_3 = 0 \\ & \downarrow & & & \nearrow & \\ & T_4 = \begin{smallmatrix} 2 \\ 3 \end{smallmatrix} \text{ } 2 & \longrightarrow & T_5 = \text{ } 2. & & \end{array}$$

- (1) All support τ -tilting A -modules $T_i (i = 1, 2, 3, 4, 5)$ are support τ -tilting B -modules by Theorem 3.9(1).
- (2) Since $2 \in \text{Fac } T_i (i = 1, 4, 5)$, we have three support τ -tilting B -modules $P_1 \oplus T_1, P_1 \oplus T_4, P_1 \oplus T_5$ by Theorem 3.9(3). Moreover, $P_1 \oplus T_1, P_1 \oplus T_4$ are τ -tilting B -modules since T_1, T_4 are τ -tilting A -modules by Corollary 3.10.
- (3) Since $\text{Hom}_A(2, \text{Fac } T_i) = 0 (i = 2, 3)$, we have two support τ -tilting B -modules $S_1 \oplus T_2, S_1 \oplus T_3$ by Theorem 3.9(4).

In fact, the Hasse quiver $Q(\text{st-tilt } B)$ is as follows.



Acknowledgment

The authors would like to thank Professor Zhaoyong Huang and Dong Yang for continuous encouragement.

Funding

This work was partially supported by National Natural Science Foundation of China (Grant No. 11971225).

References

- [1] Adachi, T., Iyama, O., Reiten, I. (2014). τ -tilting theory. *Compos. Math.* 150(3):415–452. DOI: [10.1112/S0010437X13007422](https://doi.org/10.1112/S0010437X13007422).
- [2] Asai, S. (2020). Semibricks. *Int. Math. Res. Not. IMRN*. 2020(16):4993–5054. DOI: [10.1093/imrn/rny150](https://doi.org/10.1093/imrn/rny150)
- [3] Assem, I., Happel, D., Trepode, S. (2007). Extending tilting modules to one-point extensions by projectives. *Commun. Algebra* 35(10):2983–3006. DOI: [10.1080/00927870701404556](https://doi.org/10.1080/00927870701404556).
- [4] Auslander, M., Smalø, S. O. (1981). Almost split sequences in subcategories. *J. Algebra* 69(2):426–454. DOI: [10.1016/0021-8693\(81\)90214-3](https://doi.org/10.1016/0021-8693(81)90214-3).
- [5] Gabriel, P. (1962). Des catégories abéliennes (French). *Bul. Soc. Math. France*. 79:323–448. DOI: [10.24033/bsmf.1583](https://doi.org/10.24033/bsmf.1583).
- [6] Ringel, C.M. (1976). Representations of K -species and bimodules. *J. Algebra* 41(2):269–302. DOI: [10.1016/0021-8693\(76\)90184-8](https://doi.org/10.1016/0021-8693(76)90184-8).
- [7] Simson, D., Skowroński, A. (2007). *Elements of the Representation Theory of Associative Algebras*. Vol. 3, Representation-Infinite Tilted Algebras, London Math. Soc. Stud. Texts 72. Cambridge, MA: Cambridge University Press.
- [8] Suarez, P. (2018). τ -Tilting modules over one-point extensions by a projective module. *Algebr. Represent. Theor.* 21(4):769–786. DOI: [10.1007/s10468-017-9737-5](https://doi.org/10.1007/s10468-017-9737-5).