

A note on the Hasse quiver of τ -tilting modules

Hanpeng Gao

*Department of Mathematics, Nanjing University
No. 22 Hankou Road, Nanjing, Jiangsu 210093, P. R. China
hpgao07@163.com*

Received 1 July 2020

Accepted 23 August 2020

Published 30 September 2020

Communicated by S. R. López-Permouth

Let Λ be an algebra with an indecomposable projective-injective module. Adachi gave a method to construct the Hasse quiver of support τ -tilting Λ -modules. In this paper, we will show that it can be restricted to τ -tilting modules.

Keywords: τ -tilting modules; Hasse quiver; support τ -tilting quiver.

Mathematics Subject Classification 2020: 16G20, 16G60

1. Introduction

In 2014, Adachi *et al.* [2] introduced the concept of support τ -tilting modules as a generalization of tilting modules. They showed that, in contrast to tilting modules, it is always possible to exchange a given indecomposable summand of a support τ -tilting module for a unique other indecomposable and obtain a new support τ -tilting module. This process, called mutation, is essential in cluster theory. In the same paper, the authors also showed that the support τ -tilting modules are in bijection with several other important classes in representation theory including functorially finite torsion classes introduced in [3], 2-term silting complexes introduced in [4], and cluster-tilting objects in the cluster category.


Let Λ be a finite dimensional K -algebra over an algebraically closed field K . A Λ -module M is called τ -tilting if $\text{Hom}_{\Lambda}(M, \tau M) = 0$ and $|M| = |\Lambda|$. A module is called *support τ -tilting* if it is a τ -tilting $\Lambda/\Lambda e\Lambda$ -module for some idempotent e of Λ . We will denote by $\tau\text{-tilt } \Lambda$ (respectively, $s\tau\text{-tilt } \Lambda$) the set of isomorphism classes of basic τ -tilting (respectively, support τ -tilting) Λ -modules. For any two support τ -tilting Λ -modules M, N , we write $M \geq N$ if $\text{Fac}(M) \supseteq \text{Fac}(N)$. Then \geq gives a partial order on support τ -tilting Λ -modules. The associated Hasse quiver (support

τ -tilting quiver) $H(\Lambda)$ is as follows:

- The set of vertices is all basic support τ -tilting Λ -modules.
- Draw an arrow from M to N if $M > N$ and there is no support τ -tilting Λ -module L such that $M > L > N$.

Let \mathcal{N} be a subposet of $H(\Lambda)$ and $\mathcal{N}' := H(\Lambda) \setminus \mathcal{N}$. Adachi define a new quiver $H(\Lambda)^{\mathcal{N}}$ from $H(\Lambda)$ as follows (see [1, Definition 3.2]):

- vertices: vertices in $H(\Lambda)$ and \mathcal{N}^+ where \mathcal{N}^+ is a copy of \mathcal{N} .
- arrows: $\{a_1 \rightarrow a_2 \mid a_1 \rightarrow a_2 \in \mathcal{N}'\} \coprod \{n_2 \rightarrow a_2 \mid n_2 \rightarrow a_2, n_2 \in \mathcal{N}, a_2 \in \mathcal{N}'\}$
 $\coprod \{n_1 \rightarrow n_2, n_1^+ \rightarrow n_2^+ \mid n_1 \rightarrow n_2 \in \mathcal{N}\} \coprod \{a_1 \rightarrow n_1^+ \mid a_1 \rightarrow n_1, n_1 \in \mathcal{N}, a_1 \in \mathcal{N}'\}$
 $\coprod \{n_1^+ \rightarrow n_1 \mid n_1 \in \mathcal{N}\}.$



Suppose that Λ is a basic finite dimensional K -algebra with an indecomposable projective-injective module Q , Adachi shown that $\text{Soc}(Q)$ is a two-sided ideal of Λ (see [1, Proposition 3.1(1)]). Let $\overline{\Lambda} := \Lambda / \text{Soc}(Q)$ and

$$\mathcal{N}_1 := \{N \in \text{st-tilt } \overline{\Lambda} \mid Q/\text{Soc}(Q) \in \text{add } N \text{ and } \text{Hom}_{\Lambda}(N, Q) = 0\}.$$

It is shown that there is an isomorphism of posets $H(\Lambda) \rightarrow H(\overline{\Lambda})^{\mathcal{N}_1}$ [1, Theorem 3.3].

In this paper, we will show that this result can be restricted to τ -tilting modules. More precisely, for an algebra Λ , let $Q(X)$ be the full subquiver of $H(\Lambda)$ consisting of those vertices in X for a subset X of $H(\Lambda)$. Now, considering the set

$$\mathcal{N} := \{N \in \text{st-tilt } \overline{\Lambda} \mid Q/\text{Soc}(Q) \in \text{add } N, \text{Hom}_{\Lambda}(N, Q) = 0 \text{ and } |N| = |\overline{\Lambda}| - 1\},$$

we have the following result.

Theorem 1.1. *Let Λ be an algebra with an indecomposable projective-injective module Q .*

(1) *If Q is simple, then there is an isomorphism*

$$Q(\tau\text{-tilt } \Lambda) \rightarrow Q(\tau\text{-tilt } \overline{\Lambda}).$$

(2) If Q is not simple, then there is an isomorphism

$$Q(\tau\text{-tilt } \Lambda) \rightarrow Q\left(\tau\text{-tilt } \overline{\Lambda} \coprod \mathcal{N}\right).$$

Moreover, if $Q/\text{Soc}(Q)$ has $\text{Soc}(Q)$ as a composition factor, then $\mathcal{N} = \emptyset$. Hence, we have an isomorphism $Q(\tau\text{-tilt } \Lambda) \rightarrow Q(\tau\text{-tilt } \overline{\Lambda})$.

As an application, we can calculate the number of τ -tilting modules over linearly Dynkin type algebras whose square radical are zero.

Throughout this paper, all algebras will be basic, connected, finite dimensional K -algebras over an algebraically closed field K . For an algebra Λ , we denote by $\text{mod } \Lambda$ the category of finitely generated right Λ -modules and by τ the Auslander–Reiten translation of Λ . For $M \in \text{mod } \Lambda$, we also denote by $|M|$ the number of pairwise nonisomorphic indecomposable summands of M and by $\text{add } M$ the full subcategory of $\text{mod } \Lambda$ consisting of direct summands of finite direct sums of copies of M . For a set X , we denote by $|X|$ the cardinality of X . For two sets X, Y , $X \coprod Y$ means the disjoint union.

2. Main Results

Let Λ be an algebra. We always assume that Λ has an indecomposable projective-injective module Q and $\overline{\Lambda} := \Lambda/\text{Soc}(Q)$. Considering the following functor:

$$\overline{(-)} := - \otimes_{\Lambda} \overline{\Lambda} : \text{mod } \Lambda \rightarrow \text{mod } \overline{\Lambda}.$$

Then we have $\overline{Q} = Q/\text{Soc}(Q)$ and $\overline{M} \cong M$ for all indecomposable Λ -modules M which are not isomorphism to Q by [1, Proposition 3.1(2)]. We will denote by $\alpha(M)$ a basic Λ -module such that $\text{add}(\alpha(M)) = \text{add } \overline{M}$.

We need the following lemma.

Lemma 2.1. *Assume Q is not simple and $U \in \text{mod } \overline{\Lambda}$ does not have \overline{Q} as a direct summand. Then*

- (1) $U \in \tau\text{-tilt } \Lambda$ if and only if $U \in \tau\text{-tilt } \overline{\Lambda}$.
- (2) $Q \oplus \overline{Q} \oplus U \in \tau\text{-tilt } \Lambda$ if and only if $\overline{Q} \oplus U \in s\tau\text{-tilt } \Lambda$ and $|\overline{Q} \oplus U| = |\Lambda| - 1$ if and only if $\overline{Q} \oplus U \in s\tau\text{-tilt } \overline{\Lambda}$, $|\overline{Q} \oplus U| = |\overline{\Lambda}| - 1$ and $\text{Hom}_{\Lambda}(\overline{Q} \oplus U, Q) = 0$.
- (3) $Q \oplus U \in \tau\text{-tilt } \Lambda$ if and only if $\overline{Q} \oplus U \in \tau\text{-tilt } \Lambda$ and $\text{Hom}_{\Lambda}(\overline{Q} \oplus U, Q) \neq 0$.

Proof. Note that a support τ -tilting Λ -module M is τ -tilting if and only if $|M| = |\Lambda|$. Hence (1), (2) and (3) follow from [1, Proposition 3.7]. \square

Lemma 2.2. *Assume Q is not simple. We have*

$$\{M \in \tau\text{-tilt } \Lambda \mid Q \notin \text{add } M, \overline{Q} \in \text{add } M\} = \emptyset.$$

Proof. Let $M \in \{M \in \tau\text{-tilt } \Lambda \mid Q \notin \text{add } M, \overline{Q} \in \text{add } M\}$. Write $M = \overline{Q} \oplus X$ where X does not have $Q \oplus \overline{Q}$ as a direct summand, we have $Q \notin \text{Fac } X$ since Q is

projective. By [1, Proposition 3.7], $Q \oplus \overline{Q} \oplus X \in \text{st-tilt } \Lambda$ and $|Q \oplus \overline{Q} \oplus X| = |\Lambda| + 1$ which implies $Q \in \text{Fac } X$. This is a contradiction. \square

Now, we decompose $\tau\text{-tilt } \Lambda$ as the following three parts.

$$\mathcal{M}_1 := \{M \in \tau\text{-tilt } \Lambda \mid Q, \overline{Q} \notin \text{add } M\},$$

$$\mathcal{M}_2 := \{M \in \tau\text{-tilt } \Lambda \mid Q, \overline{Q} \in \text{add } M\},$$

$$\mathcal{M}_3 := \{M \in \tau\text{-tilt } \Lambda \mid Q \in \text{add } M, \overline{Q} \notin \text{add } M\}.$$

If $X \in \tau\text{-tilt } \overline{\Lambda}$, then X is sincere by [2, Proposition 2.2(a)]. Hence, X is a sincere Λ -module. Thus $\text{Hom}_\Lambda(N, Q) \neq 0$ since Q is indecomposable injective. Therefore, $\{N \in \tau\text{-tilt } \overline{\Lambda} \mid \overline{Q} \in \text{add } N, \text{Hom}_\Lambda(N, Q) = 0\} = \emptyset$. The following proposition can be obtained by Lemma 2.1 immediately.

Proposition 2.1. *Assume that Q is not simple. Then there are bijections*

$$\mathcal{M}_1 \rightarrow \mathcal{N}_1, \quad \mathcal{M}_2 \rightarrow \mathcal{N}_2, \quad \mathcal{M}_3 \rightarrow \mathcal{N}_3$$

given by $M \rightarrow \alpha(M)$ where

$$\mathcal{N}_1 := \{N \in \tau\text{-tilt } \overline{\Lambda} \mid \overline{Q} \notin \text{add } N\},$$

$$\mathcal{N}_2 := \mathcal{N} = \{N \in \text{st-tilt } \overline{\Lambda} \mid \overline{Q} \in \text{add } N, \text{Hom}_\Lambda(N, Q) = 0 \text{ and } |N| = |\overline{\Lambda}| - 1\},$$

$$\mathcal{N}_3 = \{N \in \tau\text{-tilt } \overline{\Lambda} \mid \overline{Q} \in \text{add } N, \text{Hom}_\Lambda(N, Q) \neq 0\}.$$

In particular, there is a bijection

$$\alpha : \tau\text{-tilt } \Lambda \rightarrow \tau\text{-tilt } \overline{\Lambda} \coprod \mathcal{N}.$$

Corollary 2.1. *We have*

$$\tau\text{-tilt } \Lambda = \{N \mid N \in \mathcal{N}_1\} \coprod \{Q \oplus N \mid N \in \mathcal{N}_2\} \coprod \{Q \oplus (N/\overline{Q}) \mid N \in \mathcal{N}_3\}.$$

Now, we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. (1) It is clearly since $\tau\text{-tilt } \Lambda = \{Q \oplus M \mid M \in \tau\text{-tilt } \overline{\Lambda}\}$ where Q is a simple projective-injective Λ -module.

(2) By Proposition 2.2, we have a bijection

$$\alpha : \tau\text{-tilt } \Lambda \rightarrow \tau\text{-tilt } \overline{\Lambda} \coprod \mathcal{N}.$$

We only need to show that, for any $M, L \in \tau\text{-tilt } \Lambda$, $M \geq L$ if and only if $\alpha(M) \geq \alpha(L)$. In fact, if $M \geq L$, then $L \in \text{Fac } M$ and we have $\overline{L} \in \text{Fac } \overline{M}$ which implies $\alpha(M) \geq \alpha(L)$. Conversely, let $\alpha(M) \geq \alpha(L)$. If both M and L are in \mathcal{M}_i ($i = 1, 2, 3$), then it is clear that $M \geq L$. Otherwise,

Case 1. If $L \in \mathcal{M}_1$, then $\alpha(L) = L$. Note that $\overline{M} \in \text{Fac } M$, we have $M \geq \alpha(M) \geq \alpha(L) = L$.

Case 2. If $L \in \mathcal{M}_2$, then $M \in \mathcal{M}_3$ since \mathcal{N}_1 has no \overline{Q} as a direct summand. Thus $\alpha(L) = Q \oplus \overline{L} \in \text{Fac } M$ because \mathcal{M}_3 has Q as a direct summand.

Case 3. If $L \in \mathcal{M}_3$, then $M \notin \mathcal{M}_1$ since \mathcal{N}_1 has no \overline{Q} as a direct summand. Assume $M \in \mathcal{M}_2$. Then $\alpha(M)$ has no $\text{Soc}(Q)$ as a composition factor and $\alpha(L)$ has $\text{Soc}(Q)$ as a composition factor. This is a contradiction with $\alpha(L) \in \text{Fac } \alpha(M)$.

Thus the assertion follows. \square


We illustrate Theorem 1.1 with the following example.

Example 2.1. Let Λ be a finite dimensional K -algebra given by the quiver

$$1 \xrightarrow{\alpha} 2 \xrightarrow{\beta} 3$$

with the relation $\alpha\beta = 0$. Take $Q = P_1$. Then Q is an indecomposable projective-injective module. The algebra $\overline{\Lambda}$ given by the quiver $1 \xrightarrow{\beta} 2 \xrightarrow{\alpha} 3$.

We draw the Hasse quivers $H(\Lambda)$ and $H(\overline{\Lambda})$ as follows:



We draw those arrows in $Q(\tau\text{-tilt } \Lambda)$ and $Q(\tau\text{-tilt } \overline{\Lambda} \amalg \mathcal{N})$ by \rightsquigarrow and \mathcal{N} is marked by red.

Considering the following quivers:

$$A_n : n \longrightarrow n-1 \longrightarrow \cdots \longrightarrow 2 \longrightarrow 1$$

$$D_n : n \longrightarrow n-1 \longrightarrow \cdots \longrightarrow 3 \begin{matrix} \nearrow 1 \\ \searrow 2 \end{matrix}$$

Take $A_n^2 := KA_n/\text{rad}^2$ and $D_n^2 := KD_n/\text{rad}^2$. Applying our results, we can give a recurrence relation about the numbers of τ -tilting modules over A_n^2 and D_n^2 .

Theorem 2.1. *Let Λ_n^2 be an algebra (A_n^2 or D_n^2). Then we have*

$$|\tau\text{-tilt } \Lambda_n^2| = |\tau\text{-tilt } \Lambda_{n-1}^2| + |\tau\text{-tilt } \Lambda_{n-2}^2|.$$

Proof. Take $Q = P_n$ which is an indecomposable projective-injective Λ_n^2 -module. Since $\text{Soc}(Q) \cong S_{n-1}$, we have $\overline{\Lambda_n^2} = \Lambda_n^2/S_{n-1} \cong \Lambda_{n-1}^2 \times K$ and $Q/S_{n-1} \cong S_n$. Hence

$$\begin{aligned} \mathcal{N} &= \{N \in \text{s}\tau\text{-tilt } \overline{\Lambda_n^2} \mid Q/\text{Soc}(Q) \in \text{add } N, \text{Hom}_{\Lambda_n^2}(N, Q) = 0 \text{ and } |N| = n-1\} \\ &= \{N \in \text{s}\tau\text{-tilt } (\Lambda_{n-1}^2 \times K) \mid S_n \in \text{add } N, \text{Hom}_{\Lambda_n^2}(N, P_n) = 0 \text{ and } |N| = n-1\} \\ &= \{S_n \oplus L \mid L \in \text{silt } \Lambda_{n-1}^2, \text{Hom}_{\Lambda_n^2}(L, P_n) = 0 \text{ and } |L| = n-2\} \\ &= \{S_n \oplus L \mid L \in \text{silt } \Lambda_{n-1}^2, \text{Hom}_{\Lambda_{n-1}^2}(L, S_{n-1}) = 0 \text{ and } |L| = n-2\} \\ &= \{S_n \oplus L \mid L \in \text{silt } \Lambda_{n-2}^2 \text{ and } |L| = n-2\} \\ &= \{S_n \oplus L \mid L \in \tau\text{-tilt } \Lambda_{n-2}^2\}. \end{aligned}$$

By Theorem 1.1, there is a bijection $Q(\tau\text{-tilt } \Lambda_n^2) \rightarrow Q(\tau\text{-tilt } (\Lambda_{n-1}^2 \times K) \amalg \mathcal{N})$. Thus

$$|\tau\text{-tilt } \Lambda_n^2| = |\tau\text{-tilt } (\Lambda_{n-1}^2 \times K)| + |\mathcal{N}| = |\tau\text{-tilt } \Lambda_{n-1}^2| + |\tau\text{-tilt } \Lambda_{n-2}^2|. \quad \square$$

Corollary 2.2.

$$\begin{aligned} (1) \quad |\tau\text{-tilt } A_n^2| &= \frac{(1+\sqrt{5})^{n+1} - (1-\sqrt{5})^{n+1}}{\sqrt{5} \cdot 2^{n+1}}. \\ (2) \quad |\tau\text{-tilt } D_n^2| &= \frac{(2\sqrt{5}-1)(1+\sqrt{5})^{n-1} + (2\sqrt{5}+1)(1-\sqrt{5})^{n-1}}{\sqrt{5} \cdot 2^{n-1}}. \end{aligned}$$

Acknowledgments

The author would like to thank Professor Zhaoyong Huang for helpful discussions. He also thanks the referee for the useful and detailed suggestions. This work was partially supported by the National Natural Science Foundation of China (No. 11971225).

References

- [1] T. Adachi, The classification of τ -tilting modules over Nakayama algebras, *J. Algebra* **452** (2016) 227–262.
- [2] T. Adachi, O. Iyama and I. Reiten, τ -tilting theory, *Compos. Math.* **150** (2014) 415–452.
- [3] M. Auslander and S. Smalø, Almost split sequences in subcategories, *J. Algebra* **69** (1981) 426–454.
- [4] B. Keller and D. Vossieck, Aisles in derived categories, *Bull. Soc. Math. Belg. Sér. A* **40**(2) (1988) 239–253.