

Point	λ	δ_0	δ_1	δ_2	J	Test (95%)
Around the MSM optimizer solution						
Center	6.85×10^{-4}	-5.41	14.96	3.42	29,326	Reject
$\lambda \times 2$	1.37×10^{-3}	-6.37	14.96	3.42	29,306	Reject
$\lambda \div 2$	3.42×10^{-4}	-3.85	14.96	3.42	29,482	Reject
$\delta_1 + 1$	6.85×10^{-4}	-2.16	15.96	3.42	29,377	Reject
$\delta_1 - 1$	6.85×10^{-4}	-8.61	13.96	3.42	29,380	Reject
$\delta_2 + 0.2$	6.85×10^{-4}	-7.65	14.96	3.62	29,369	Reject
$\delta_2 - 0.2$	6.85×10^{-4}	-3.13	14.96	3.22	29,382	Reject
Around the ELVIS point						
Center	5.43×10^{-7}	5.25	4.30	0.54	32,940	Reject
$\lambda \times 2$	1.09×10^{-6}	5.22	4.30	0.54	33,033	Reject
$\lambda \div 2$	2.71×10^{-7}	5.25	4.30	0.54	32,910	Reject
$\delta_1 + 1$	5.43×10^{-7}	8.70	5.30	0.54	34,220	Reject
$\delta_1 - 1$	5.43×10^{-7}	1.79	3.30	0.54	31,206	Reject
$\delta_2 + 0.2$	5.43×10^{-7}	2.72	4.30	0.74	30,589	Reject
$\delta_2 - 0.2$	5.43×10^{-7}	7.82	4.30	0.34	35,783	Reject

Real data (lag_m, trim 0.5%). Naive MSM (Tim Conley's variant): ε drawn from the corporate pool, $S = 250$, same draws at all points; δ_0 profiled at each point. $J = n\hat{L}_n$ compared with $\chi^2_{5,.95} = 11.07$.