

Connectedness

Stepan Kuznetsov

Computer Science Department, Higher School of Economics

Outline

Connectedness of Random Graphs

Random Graph

- Recall that $G(n, p)$ is the Erdős – Rényi random graph on n vertices, where each edge is planted with probability p , independently from others.

Random Graph

- Recall that $G(n, p)$ is the Erdős – Rényi random graph on n vertices, where each edge is planted with probability p , independently from others.
- The probability p could depend on n .

Random Graph

- Recall that $G(n, p)$ is the Erdős – Rényi random graph on n vertices, where each edge is planted with probability p , independently from others.
- The probability p could depend on n .
- We start with $p = \frac{1}{2}$.

Random Graph

- Recall that $G(n, p)$ is the Erdős – Rényi random graph on n vertices, where each edge is planted with probability p , independently from others.
- The probability p could depend on n .
- We start with $p = \frac{1}{2}$.
- Recall that the probability of $G(n, \frac{1}{2})$ to have an isolated vertex tends to 0 when $n \rightarrow \infty$.

Connectedness of $G(n, \frac{1}{2})$

- Let us generalize that computation and estimate the probability of $G(n, \frac{1}{2})$ to be *connected*.

Connectedness of $G(n, \frac{1}{2})$

- Let us generalize that computation and estimate the probability of $G(n, \frac{1}{2})$ to be *connected*.
- $P(G(n, \frac{1}{2}) \text{ is not connected}) \leq$

$$\sum_{\emptyset \neq V' \subsetneq V} P(V' \text{ is isolated}) = \sum_{m=1}^{n-1} \frac{\binom{n}{m}}{2^{m(n-m)}}$$

Connectedness of $G(n, \frac{1}{2})$

- Let us generalize that computation and estimate the probability of $G(n, \frac{1}{2})$ to be *connected*.
- $P(G(n, \frac{1}{2}) \text{ is not connected}) \leq$
$$\sum_{\emptyset \neq V' \subsetneq V} P(V' \text{ is isolated}) = \sum_{m=1}^{n-1} \frac{\binom{n}{m}}{2^{m(n-m)}}$$
- Oops... this looks more like 1 (binomial formula).

Connectedness of $G(n, \frac{1}{2})$

- Let us generalize that computation and estimate the probability of $G(n, \frac{1}{2})$ to be *connected*.
- $P(G(n, \frac{1}{2}) \text{ is not connected}) \leq$
$$\sum_{\emptyset \neq V' \subsetneq V} P(V' \text{ is isolated}) = \sum_{m=1}^{n-1} \frac{\binom{n}{m}}{2^{m(n-m)}}$$
- Oops... this looks more like 1 (binomial formula).
- The problem is that our cases overlap too much.

Connectedness of $G(n, \frac{1}{2})$

- Let us do something stronger: show that the probability that *any two vertices have a common neighbour* tends to 1.

Connectedness of $G(n, \frac{1}{2})$

- Let us do something stronger: show that the probability that *any two vertices have a common neighbour* tends to 1.
- This implies connectedness, so connectedness will also have desired probability.

Connectedness of $G(n, \frac{1}{2})$

- $P(\text{there are two vertices without a common neighbour}) \leq \sum_{i \neq j} P(v_i \text{ and } v_j \text{ do not have a common neighbour}) =$

Connectedness of $G(n, \frac{1}{2})$

- $P(\text{there are two vertices without a common neighbour}) \leq$
 $\sum_{i \neq j} P(v_i \text{ and } v_j \text{ do not have a common neighbour}) =$
 $n(n-1)(\frac{3}{4})^{n-2}$

Connectedness of $G(n, \frac{1}{2})$

- $P(\text{there are two vertices without a common neighbour}) \leq$
 $\sum_{i \neq j} P(v_i \text{ and } v_j \text{ do not have a common neighbour}) =$
 $n(n-1)(\frac{3}{4})^{n-2} \rightarrow 0$

Connectedness of $G(n, \frac{1}{2})$

- $P(\text{there are two vertices without a common neighbour}) \leq \sum_{i \neq j} P(v_i \text{ and } v_j \text{ do not have a common neighbour}) = n(n-1)(\frac{3}{4})^{n-2} \rightarrow 0$
- Indeed, v_i and v_j do not have a common neighbour if and only if for any w from the remaining $n-2$ vertices one of the edges $\langle w, v_i \rangle$ and $\langle w, v_j \rangle$ was not drawn.

Connectedness of $G(n, \frac{1}{2})$

- $P(\text{there are two vertices without a common neighbour}) \leq \sum_{i \neq j} P(v_i \text{ and } v_j \text{ do not have a common neighbour}) = n(n-1)(\frac{3}{4})^{n-2} \rightarrow 0$
- Indeed, v_i and v_j do not have a common neighbour if and only if for any w from the remaining $n-2$ vertices one of the edges $\langle w, v_i \rangle$ and $\langle w, v_j \rangle$ was not drawn. This happens with probability $\frac{3}{4}$.

Boundaries for Connectedness

- Now we know that $G(n, \frac{1}{2})$ is almost surely connected, for $n \rightarrow \infty$.

Boundaries for Connectedness

- Now we know that $G(n, \frac{1}{2})$ is almost surely connected, for $n \rightarrow \infty$.
- On the other hand, as shown before, $G(n, 2^{-n})$ almost surely has an isolated vertex, thus, it is disconnected.

Boundaries for Connectedness

- Now we know that $G(n, \frac{1}{2})$ is almost surely connected, for $n \rightarrow \infty$.
- On the other hand, as shown before, $G(n, 2^{-n})$ almost surely has an isolated vertex, thus, it is disconnected.
- The exact boundaries are as follows:

Boundaries for Connectedness

- Now we know that $G(n, \frac{1}{2})$ is almost surely connected, for $n \rightarrow \infty$.
- On the other hand, as shown before, $G(n, 2^{-n})$ almost surely has an isolated vertex, thus, it is disconnected.
- The exact boundaries are as follows:
 - if $p < \frac{(1-\varepsilon) \ln n}{n}$, then $G(n, p)$ is almost surely disconnected;

Boundaries for Connectedness

- Now we know that $G(n, \frac{1}{2})$ is almost surely connected, for $n \rightarrow \infty$.
- On the other hand, as shown before, $G(n, 2^{-n})$ almost surely has an isolated vertex, thus, it is disconnected.
- The exact boundaries are as follows:
 - if $p < \frac{(1-\varepsilon) \ln n}{n}$, then $G(n, p)$ is almost surely disconnected;
 - if $p > \frac{(1+\varepsilon) \ln n}{n}$, then $G(n, p)$ is almost surely connected.