

Combinations

Vladimir Podolskii

Computer Science Department, Higher School of Economics

Outline

Unordered Pairs

Combinations

Ranking problem

Ranking Problem

Suppose we have n texts and we need to rank them by relevance. One of the standard approaches requires to compare each text with each other. How many comparisons we need to make?

Ranking problem

Solution attempt:

- There are n documents

Ranking problem

Solution attempt:

- There are n documents
- We need to compare each document to each of $n - 1$ other documents

Ranking problem

Solution attempt:

- There are n documents
- We need to compare each document to each of $n - 1$ other documents
- By the rule of product there will be $n \times (n - 1)$ comparisons

Ranking problem

Solution attempt:

- There are n documents
- We need to compare each document to each of $n - 1$ other documents
- By the rule of product there will be $n \times (n - 1)$ comparisons
- Is everything right?

Ranking problem

Consider $n = 3$ and denote texts by a, b and c

- Our solution gives $3 \times 2 = 6$ comparisons

Ranking problem

Consider $n = 3$ and denote texts by a , b and c

- Our solution gives $3 \times 2 = 6$ comparisons
- But actually, there are only 3: a vs. b , a vs. c and b vs. c

Ranking problem

Consider $n = 3$ and denote texts by a , b and c

- Our solution gives $3 \times 2 = 6$ comparisons
- But actually, there are only 3: a vs. b , a vs. c and b vs. c
- Let's review our solution

Ranking problem

Consider $n = 3$ and denote texts by a , b and c

- Our solution gives $3 \times 2 = 6$ comparisons
- But actually, there are only 3: a vs. b , a vs. c and b vs. c
- Let's review our solution
- For each text we compare it with 2 others

Ranking problem

Consider $n = 3$ and denote texts by a , b and c

- Our solution gives $3 \times 2 = 6$ comparisons
- But actually, there are only 3: a vs. b , a vs. c and b vs. c
- Let's review our solution
- For each text we compare it with 2 others
- That is, a is compared to b and c ; b is compared to a and c ; c is compared to a and b

Ranking problem

Consider $n = 3$ and denote texts by a , b and c

- Our solution gives $3 \times 2 = 6$ comparisons
- But actually, there are only 3: a vs. b , a vs. c and b vs. c
- Let's review our solution
- For each text we compare it with 2 others
- That is, a is compared to b and c ; b is compared to a and c ; c is compared to a and b
- We counted comparisons ab, ac, ba, bc, ca, cb

Ranking problem

Consider $n = 3$ and denote texts by a , b and c

- Our solution gives $3 \times 2 = 6$ comparisons
- But actually, there are only 3: a vs. b , a vs. c and b vs. c
- Let's review our solution
- For each text we compare it with 2 others
- That is, a is compared to b and c ; b is compared to a and c ; c is compared to a and b
- We counted comparisons ab, ac, ba, bc, ca, cb
- We counted each comparison twice!

Ranking problem

Now we can fix our solution:

- There are n documents

Ranking problem

Now we can fix our solution:

- There are n documents
- We need to compare to each of $n - 1$ other documents

Ranking problem

Now we can fix our solution:

- There are n documents
- We need to compare to each of $n - 1$ other documents
- By the rule of product there will be $n \times (n - 1)$ comparisons

Ranking problem

Now we can fix our solution:

- There are n documents
- We need to compare to each of $n - 1$ other documents
- By the rule of product there will be $n \times (n - 1)$ comparisons
- We have counted every a vs. b comparison twice: as a vs. b and as b vs. a

Ranking problem

Now we can fix our solution:

- There are n documents
- We need to compare to each of $n - 1$ other documents
- By the rule of product there will be $n \times (n - 1)$ comparisons
- We have counted every a vs. b comparison twice: as a vs. b and as b vs. a
- Let's just divide the result by 2

Ranking problem

Now we can fix our solution:

- There are n documents
- We need to compare to each of $n - 1$ other documents
- By the rule of product there will be $n \times (n - 1)$ comparisons
- We have counted every a vs. b comparison twice: as a vs. b and as b vs. a
- Let's just divide the result by 2
- The answer is $n(n - 1)/2$

What we have learned

- We have counted the number of unordered pairs of n objects

What we have learned

- We have counted the number of **unordered pairs** of n objects
- The answer is $n(n - 1)/2$

What we have learned

- We have counted the number of **unordered pairs** of n objects
- The answer is $n(n - 1)/2$
- When counting make sure that you count each object once

What we have learned

- We have counted the number of **unordered pairs** of n objects
- The answer is $n(n - 1)/2$
- When counting make sure that you count each object once
- If you counted each object k times, just divide the result by k

Outline

Unordered Pairs

Combinations

Road Trip

Road Trip

You are planning a road trip. You have 5 friends, but you can fit only 3 of them in your car. How many ways do you have to do it?

- Basically, we want to pick a subset of size 3 in the set of your 5 friends
- We are interested in how many subsets of size 3 are there
- Recall that elements of subset are unordered

Road Trip

Solution attempt:

- We can pick the first friend in 5 ways
- We can pick the second friend in 4 ways
- We can pick the third friend in 3 ways
- By the rule of product there are $5 \times 4 \times 3 = 60$ possibilities
- Is there a problem?
- Again we counted each option several times

Road Trip

Fixing solution:

- Note, that we actually counted **ordered** sequences of friends

Road Trip

Fixing solution:

- Note, that we actually counted **ordered** sequences of friends
- Each (unordered) group $\{a, b, c\}$ is counted $3 \times 2 = 6$ times: abc, acb, bac, bca, cab and cba

Road Trip

Fixing solution:

- Note, that we actually counted **ordered** sequences of friends
- Each (unordered) group $\{a, b, c\}$ is counted $3 \times 2 = 6$ times: abc, acb, bac, bca, cab and cba
- We can just divide the previous result by 6

Road Trip

Fixing solution:

- Note, that we actually counted **ordered** sequences of friends
- Each (unordered) group $\{a, b, c\}$ is counted $3 \times 2 = 6$ times: abc, acb, bac, bca, cab and cba
- We can just divide the previous result by 6
- The actual result is $5 \times 4 \times 3 / (3 \times 2) = 10$

Combinations

- For a set S its k -combination is a subset of S of size k

Combinations

- For a set S its k -combination is a subset of S of size k
- The number of k -combinations of n element set is denoted $\binom{n}{k}$

Combinations

- For a set S its k -combination is a subset of S of size k
- The number of k -combinations of n element set is denoted $\binom{n}{k}$
- Pronounced ' n choose k '

Combinations

- For a set S its k -combination is a subset of S of size k
- The number of k -combinations of n element set is denoted $\binom{n}{k}$
- Pronounced ' n choose k '
- We have shown $\binom{5}{3} = 10$

3-Combinations and 3-Permutations

abc	abd	abe	acd	ace	ade	bcd	bce	bde	cde
acb	adb	aeb	adc	aec	aed	bdc	bec	bed	ced
bac	bad	bae	cad	cae	dae	cbd	cbe	dbe	dce
bca	bda	bea	cda	cea	dea	cdb	ceb	deb	dec
cba	dba	eba	dca	eca	eda	dcb	ecb	edb	edc
cab	dab	eab	dac	eac	ead	dbc	ebc	ebd	ecd

3-Combinations and 3-Permutations



abc	abd	abe	acd	ace	ade	bcd	bce	bde	cde
acb	adb	aeb	adc	aec	aed	bdc	bec	bed	ced
bac	bad	bae	cad	cae	dae	cbd	cbe	dbe	dce
bca	bda	bea	cda	cea	dea	cdb	ceb	deb	dec
cba	dba	eba	dca	eca	eda	dcb	ecb	edb	edc
cab	dab	eab	dac	eac	ead	dbc	ebc	ebd	ecd

3-Combinations and 3-Permutations

	abc	abd	abe	acd	ace	ade	bcd	bce	bde	cde
	acb	adb	aeb	adc	aec	aed	bdc	bec	bed	ced
	bac	bad	bae	cad	cae	dae	cbd	cbe	dbe	dce
3!	bca	bda	bea	cda	cea	dea	cdb	ceb	deb	dec
	cba	dba	eba	dca	eca	eda	dcb	ecb	edb	edc
	cab	dab	eab	dac	eac	ead	dbc	ebc	ebd	ecd

3-Combinations and 3-Permutations

	abc	abd	abe	acd	ace	ade	bcd	bce	bde	cde
	acb	adb	aeb	adc	aec	aed	bdc	bec	bed	ced
	bac	bad	bae	cad	cae	dae	cbd	cbe	dbe	dce
3!	bca	bda	bea	cda	cea	dea	cdb	ceb	deb	dec
	cba	dba	eba	dca	eca	eda	dcb	ecb	edb	edc
	cab	dab	eab	dac	eac	ead	dbc	ebc	ebd	ecd

$$3! \binom{5}{3} = \frac{5!}{(5-3)!}$$

Combinations

- Combinations is an important setting that is needed a lot

Combinations

- Combinations is an important setting that is needed a lot
- How can it arise in Data Science? Here is an example

Combinations

- Combinations is an important setting that is needed a lot
- How can it arise in Data Science? Here is an example
- Sometimes you need to use weak ML model that cannot in principal give a good prediction on your data

Combinations

- Combinations is an important setting that is needed a lot
- How can it arise in Data Science? Here is an example
- Sometimes you need to use weak ML model that cannot in principal give a good prediction on your data
- One possible solution: enrichment by feature products

Combinations

- Suppose you have 5 numerical features a, b, c, d and e in your setting

Combinations

- Suppose you have 5 numerical features a, b, c, d and e in your setting
- To enrich the set of features we can add new features consisting of products of any 3 distinct features

Combinations

- Suppose you have 5 numerical features a, b, c, d and e in your setting
- To enrich the set of features we can add new features consisting of products of any 3 distinct features
- For example, we add $a \cdot b \cdot c, b \cdot c \cdot e, d \cdot a \cdot c$ and all other triple products

Combinations

- Suppose you have 5 numerical features a, b, c, d and e in your setting
- To enrich the set of features we can add new features consisting of products of any 3 distinct features
- For example, we add $a \cdot b \cdot c, b \cdot c \cdot e, d \cdot a \cdot c$ and all other triple products
- After that we can run our weak model on the new set of features

Combinations

- Suppose you have 5 numerical features a, b, c, d and e in your setting
- To enrich the set of features we can add new features consisting of products of any 3 distinct features
- For example, we add $a \cdot b \cdot c, b \cdot c \cdot e, d \cdot a \cdot c$ and all other triple products
- After that we can run our weak model on the new set of features
- In many cases it gives much better results

Combinations

- But how many features we have added?

Combinations

- But how many features we have added?
- Basically, for each new feature we pick k -combination of initial features

Number of Combinations

There is the following short formula for the number of combinations

Theorem

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Proof

- Need to count the number of subsets of size k of a set of size n

Proof

- Need to count the number of subsets of size k of a set of size n
- There are n choices of the first element, $(n - 1)$ choices of the second element, \dots , $(n - k + 1)$ choices of the k -th element

Proof

- Need to count the number of subsets of size k of a set of size n
- There are n choices of the first element, $(n - 1)$ choices of the second element, \dots , $(n - k + 1)$ choices of the k -th element
- This gives $n(n - 1) \cdots (n - k + 1) = \frac{n!}{(n-k)!}$

Proof

- Need to count the number of subsets of size k of a set of size n
- There are n choices of the first element, $(n - 1)$ choices of the second element, \dots , $(n - k + 1)$ choices of the k -th element
- This gives $n(n - 1) \dots (n - k + 1) = \frac{n!}{(n-k)!}$
- But this is the number of k -permutations rather than the number of k -combinations

Proof

- In other words , we counted ordered subset instead of unordered

Proof

- In other words , we counted ordered subset instead of unordered
- Each subset is counted as many times, as there are ways to order it

Proof

- In other words , we counted ordered subset instead of unordered
- Each subset is counted as many times, as there are ways to order it
- There are $k!$ ways to order a subset of size k

Proof

- In other words , we counted ordered subset instead of unordered
- Each subset is counted as many times, as there are ways to order it
- There are $k!$ ways to order a subset of size k
- Thus, each subset is counted $k!$ times

Proof

- In other words , we counted ordered subset instead of unordered
- Each subset is counted as many times, as there are ways to order it
- There are $k!$ ways to order a subset of size k
- Thus, each subset is counted $k!$ times
- This finally gives $\frac{n!}{k!(n-k)!}$

Conclusion

- In this week we started to study combinatorics

Conclusion

- In this week we started to study combinatorics
- We started with simple observations

Conclusion

- In this week we started to study combinatorics
- We started with simple observations
- We built on them more complicated constructions

Conclusion

- In this week we started to study combinatorics
- We started with simple observations
- We built on them more complicated constructions
- We reached a rather non-trivial question of counting combinations

Conclusion

- In this week we started to study combinatorics
- We started with simple observations
- We built on them more complicated constructions
- We reached a rather non-trivial question of counting combinations
- Next week we will proceed with combinations and will see that they have nice and interesting properties