AMRITA VISHWA VIDYAPEETHAM
DEEMED TO BE UNIVERSITY UNDER SECTION 3 OF UGC ACT, 1956

SCHOOL OF ARTIFICIAL INTELLIGENCE COIMBATORE

CardioCog: An Explainable AI Framework for Cognitive Health Assessment and Early Prediction of Cognitive Decline Using Cardiovascular, Genetic, and Clinical Factors

Final Review – B.Tech Project

Akula Akhil Vighnesh – CB.SC.U4AIE23303

Vada Gouri Hansika Reddy – CB.SC.U4AIE23304

Pilli Venkata Bhargav Reddy – CB.SC.U4AIE23329

Marella Venkata Mahidhar Reddy – CB.SC.U4AIE23354

Team D07

Guide: Dr. Manoj Bhat K, Assistant Professor | Co-guide: Amrutha Veluppal, Assistant Professor

Introduction & Motivation

Domain: AI / Healthcare Analytics / XAI / Biomedical Data Science

Introduction

- Dementia and Alzheimer's disease are increasing worldwide, affecting memory, thinking and quality of life.
- Hypertension, diabetes, obesity and stroke are associated with a higher risk of cognitive decline.
- Clinics already record cognition, vascular history, genetics and sometimes PET, but rarely combine them into one explainable estimate of near-term risk.

Motivation

- Decline is hard to reverse once noticeable, so the years before diagnosis are the window for intervention.
- Early identification allows low-cost steps and gives families time to plan.
- In NACC, **14.5%** of people without dementia converted within three years; among those with MCI, **41.5%**.

Problem Statement

Clinical, cardiovascular, genetic and PET information is rarely combined into one explainable estimate of near-term decline, and it is unclear which source actually adds information.

For a person without dementia at a visit, CardioCog estimates over the next three years:

- the risk of **dementia**
- the risk of a **CDR-SB rise ≥ 1**
- the expected **change in CDR-SB**

In scope	**Out of scope**
3-year dementia and CDR-SB decline risk	Clinical diagnosis
Demographic, cardiovascular, genetic, cognitive, clinical-stage and PET data	Treatment recommendation
SHAP explanations and quoted published evidence (RAG)	Post-diagnosis prognosis
Validation at 9 held-out NACC centres	Validation on a non-NACC cohort

Research Questions

RQ	Question
1	How accurately can dementia and CDR-SB decline be predicted, within normal and MCI groups separately?
2	Does cardiovascular history add to demographics, genetics and current cognition?
3	Do amyloid and tau PET add to the clinical model?
4	Do deep models, including a modality-encoder fusion network, beat classical ones?
5	Do earlier visits help?
6	Can one model cope with whole categories of missing information?
7	Can each estimate be explained and supported with published evidence?

RQ2 is the project's original hypothesis: that cardiovascular history improves the prediction. The study tests it rather than assuming it.

Literature Review

Work	Core method	Relevance / limitation
Livingston et al. (2020)	Lancet Commission, 12 modifiable risk factors	Vascular risk matters over the life course; mostly midlife exposure
Alaka et al. (2025)	CAIDE + resting heart rate, random forest, NACC	Same database; AUC 0.80–0.91, no within-stage results
Stern & Linial (2026)	CatBoost on UK Biobank codes + APOE	AUROC 0.773 without cognitive inputs
Aschwanden et al. (2020)	Random survival forests, 52 predictors	Cardiovascular factors less important than expected
Yi et al. (2023)	XGBoost + SHAP for diagnosis	CDR-SB most important; diagnosis, not prediction
Gorishniy et al. (2021)	FT-Transformer for tabular data	No deep model consistently beats boosted trees
Lewis et al. (2020)	Retrieval-augmented generation	Grounds generated text in a fixed library

Research Gap and Our Approach

Research gap

- Few studies measure what cardiovascular factors add **once current cognition is known**.
- Results are rarely reported **within** normal and MCI groups, where the clinical question lies.
- Many models lack held-out sites and checks for information leakage.

Our approach

- One leakage-controlled NACC cohort: 19,470 people, nine whole centres held out.
- Category-by-category ablation, with results within diagnostic groups.
- 17 independent verification checks on the whole pipeline.

Objectives

- Build a leakage-free three-year cohort from NACC with person-level splits and nine whole centres held out.
- Predict dementia and CDR-SB decline, and estimate the three-year CDR-SB change, from 48 features in five categories.
- Measure what cardiovascular history, genetics, PET and earlier visits add once current cognition is known.
- Compare classical models, deep tabular models and a modality-encoder fusion network that copes with missing categories.
- Explain each estimate with SHAP and quoted published evidence in a research prototype, and verify the pipeline independently.

Changes After the First Review

First-review prototype

3D CNN on raw amyloid PET + 34 clinical features; outcome: 1-year CDR-SB rise ≥ 0.5. Mean AUROC 0.786 at the review, 0.652 when the same notebook was re-run.

Diagnosis

Only 198 people and 17 events, so a handful of participants decides the score. PET time frames were mixed, with no registration, and SHAP was fitted on all data.

Decision

Move to the full clinical cohort with three-year outcomes and thousands of events, and treat PET images as a separate, validated pipeline.

Datasets: Clinical Tables and PET

Source (NACC Freeze 74)	Size	Used for
Clinical visits (UDS forms)	217,598 visits, 57,038 participants, 46 centres, 2005–2026	All 48 features and both outcomes
Amyloid PET summary (PET core)	About 5,100 participants; Centiloid value and amyloid status	PET late-fusion model
Tau PET summary (PET core)	2,724 participants; meta-temporal and entorhinal SUVR	PET late-fusion model
Raw PET images (DICOM)	599 scans (318 amyloid, 232 tau, 49 FDG) from 446 participants, about 100 GB	Image pipeline: SUVR check and 3D CNN

Clinical table

2,649 columns, mostly empty; 114 variables mapped by hand, giving 48 features. The test battery changed in 2015, so scores were converted to adjusted z-scores.

PET images

Only 5 of the 446 people with images later developed dementia, so images were used to check measurements, not to predict dementia.

Data: NACC Freeze 74 and the Cohort


Step	Participants
All participants	57,038
First visit, in person, UDS v1–3	55,461
not demented at that visit	38,107
outcome known at 3 years	19,470
converted to dementia	2,818 (14.5%)

Three groups, very different risk

Normal: 12,377 (1.6% convert)


MCI: 5,993 (41.5%)

Impaired, not MCI: 1,100 (12.3%)



Person-level split 70 / 15 / 15; 9 of 46 centres held out; everything learned from data is fitted on training only.

Outcomes and Censoring



Primary: dementia within 3 years

1 if diagnosed within 3 years; 0 if not and followed for 3 years; otherwise unknown and left out.

Secondary outcomes


CDR-SB rise ≥ 1 within 3 years (20,355 people, 5,035 events) and the 3-year change in CDR-SB. A Cox model that keeps censored people gives the same conclusions.

48 Features in Five Categories


Category	Features
Demographics	8
Cardiovascular and medical	25
Cognitive tests (z-scores)	9
Clinical stage (CDR-SB, diagnosis)	3
Genetics (APOE ε4, family history)	3
PET (second-stage model only)	4

Harmonising tests across form versions

Each score is compared with the expected score for the person's age, sex and education (fitted on cognitively normal people): z = (score − expected) / SD. MoCA 21 for a 72-year-old woman with 16 years of education: expected 25.8, z = −1.9.



Methodology: The CardioCog Pipeline



Rule for every step

Only information available at the index visit is used; anything learned from data is fitted on the training set; results are reported on the test set and on centres never seen in training.

Techniques & Tools

Models compared on the same 48 features

- Logistic regression, random forest, **XGBoost** (main model), MLP, FT-Transformer
- Modality-encoder fusion network with attention, risk and progression heads

Evaluation

- AUROC, AUPRC, Brier score, calibration, net benefit; 1,000 bootstrap resamples


Explainability and evidence

- SHAP, category ablation, Grad-CAM for the PET CNN
- Hybrid retrieval: MiniLM embeddings + BM25, merged by rank fusion

Data and platforms

- NACC Freeze 74: UDS clinical visits (v1–3), PET core values and PET images
- Python (scikit-learn, XGBoost, PyTorch, SHAP, lifelines, SimpleITK); FastAPI + React prototype

Modality-Encoder Fusion Network



- Each category has its own encoder; attention combines the five tokens into one summary.
- A missing category gets a learned token; whole categories are hidden with probability 0.35 in training.
- Loss: cross-entropy for risk + 0.5 × Huber loss for the CDR-SB change.

Results: Dementia Within Three Years

Model	Test AUROC (95% CI)	AUPRC	Brier	Held-out AUROC	Brier
Logistic regression	0.936 (0.924–0.949)	0.721	0.066	0.945	0.060
Random forest	0.932 (0.919–0.944)	0.704	0.070	0.940	0.064
XGBoost	**0.938 (0.925–0.948)**	0.713	0.066	**0.946**	0.059
MLP	0.938 (0.926–0.949)	0.721	0.066	0.946	0.059
FT-Transformer	0.939 (0.927–0.949)	0.720	0.066	0.945	0.061

Test set: 2,286 people, 332 events. Held-out centres: 4,064 people, 545 events.

What it shows

All five models discriminate well and are well calibrated, at new centres as well as on the test set.

What it does not show

The intervals overlap almost completely: deep models give no gain over logistic regression or boosted trees.

Is 0.94 Too Good? Results Within Groups

Group	Test	Held-out
All not demented	0.937	0.946
Cognitively normal	0.884	0.876
MCI	0.835	0.846

Converters: normal 24 / 37, MCI 289 / 496 (test / held-out).

Critical check

Removing all current-cognition information drops AUROC from 0.937 to 0.746.


Interpretation

Diagnosis alone gives AUROC 0.84, because MCI converts 25 times as often as normal cognition. Within MCI the model still ranks risk at 0.84.

Meaning


The model estimates short-term progression given the current clinical stage. It is not leakage: inputs come from the index visit, the label from later visits.

What Drives the Prediction?



Category	Alone	Gain added last (test)	Gain (held-out)
Cognitive tests	0.896	+0.022	+0.019
Clinical stage	0.897	+0.015	+0.021
Demographics	0.671	+0.010	+0.009
Genetics (APOE)	0.645	+0.003	+0.001
Cardiovascular / medical	0.626	+0.001	+0.001

The Cardiovascular Question (RQ2)




- Alone: AUROC **0.626**.
- After cognition: **+0.001** for dementia, +0.0004 for CDR-SB decline.
- Cox model: hypertension HR 0.995; diabetes and stroke ≈ 1.

Interpretation

Limited added value in this cohort, not evidence that vascular health is unimportant: median age 71, midlife exposure unobserved, and past damage already shows in the cognitive scores.


Explainability: Shapley Values



Category	SHAP share
Clinical stage	40.8%
Cognitive tests	35.0%
Demographics	15.5%
Cardiovascular	5.1%
Genetics	3.5%

- Each prediction is split into contributions from each feature.
- Rankings are stable over 8 refits (Spearman 0.89–0.97).
- Low BMI raises the estimated risk: weight loss before diagnosis, not a causal effect.

Cognitive Decline as the Outcome




CDR-SB rise ≥ 1

20,355 people, 5,035 events. AUROC 0.858 test, 0.885 held-out; within normal cognition 0.76–0.83.

Predicted 3-year CDR-SB change

Mean absolute error 0.69 → 0.60 (test), 0.82 → 0.72 (held-out); MCI 1.90 → 1.59; no gain for normal cognition.

Fusion Network: Results




AUROC	Fusion net	XGBoost
Test	0.934	0.937
Held-out	0.943	0.946
MCI (test)	0.831	0.835

Finding


As good as XGBoost but no better. Attention is 18–22% for every category (uniform = 20%), so it does not explain predictions; SHAP and ablation are used instead.

Earlier Visits and PET Measures



Visit history (RQ5)


Using up to all earlier visits adds +0.000 to +0.005 AUROC.



Amyloid and tau PET (RQ3)

In MCI, 34% of amyloid-positive vs 7% of negative people convert within 2 years, but adding PET to the model gives no demonstrable gain (243 people, 33 events).

PET Image Pipeline: DICOM to SUVR




Group averages only (318 amyloid, 232 tau, 49 FDG scans).

- **Convert:** 599 dynamic DICOM series → one static volume.
- **Register:** to MNI152, then to a study PET template.
- **Quantify:** SUVR = mean cortex ÷ mean cerebellum.

Validation against the NACC PET core

Pearson r = 0.977 over 318 scans (0.96–0.99 per tracer); mean difference −0.043.

3D CNN for Amyloid Status



Method (292 scans, 84 positive)	AUROC
Cortical SUVR (one number)	0.982
Regional logistic model	0.971
3D CNN	0.960


Interpretation

The image model works, but the label itself comes from cortical SUVR, so one number is hard to beat.

Limitation

Grad-CAM maps were diffuse, and only 5 imaged people converted, so images were not used to predict dementia.

Validation: Calibration, Usefulness, Missing Data



- Predicted 14.5% = observed 14.5% (test).
- At 20% risk: sensitivity 83%, specificity 88%.
- Useful (net benefit) from 2% to 60% risk.


Brier score	Standard	Dropout	Specialist
Everything	0.066	0.067	0.066
No clinical stage	0.185	0.076	0.074
No cognition at all	0.441	0.115	0.111
Demographics only	0.444	0.120	0.118

Modality dropout keeps one model close to a specialist model.

Research Prototype for Clinicians

Example: 72-year-old woman, 16 years of education, MCI, CDR-SB 2, MoCA 21.


- FastAPI + React; the form is generated from the 48 trained features.
- Risk shown next to the observed rate of similar people, with SHAP by category.
- Plausibility checks flag contradictory entries.
- 27 of 28 behaviour checks pass.



Plain-Language Report

- **Summary:** risk next to the observed rate of similar people (39% vs 40%).
- **Main reasons:** largest SHAP contributions in plain words.
- **Reliability:** missing categories and warnings.
- **Published evidence:** quoted passages with their source.
- **Limits:** a fixed statement that it is not a diagnosis.

Evidence is quoted, never generated, so the report cannot state anything the model or the literature did not provide.



SUMMARY

The estimated chance of a dementia diagnosis within the next three years is 39%. For comparison, 40% of 1,283 research participants aged 65-74 with a status of MCI were diagnosed with dementia within three years.

MAIN REASONS

The estimate was raised mainly by a CDR Sum of Boxes of 2; a current status of MCI; screening test (MoCA / MMSE) below expected for age, sex and education.

It was lowered by age 72.

Overall, the model relied most on the clinician's rating of daily function and cognitive status.

RELIABILITY

Not available, so not used: genes. The estimate is less certain than with complete information.


PUBLISHED EVIDENCE

"We aimed to quantify the relative risk of progression from mild cognitive impairment (MCI) to dementia in people with and without diabetes, and with and without the MetS (MetS); and to identify potential modifiers of the risk of progression from MCI to dementia in people with diabetes or MetS.MethodsWe searched ..."
MCI progression with diabetes meta-analysis 2018 (2018)

"Risk factors in early life (education), midlife (hypertension, obesity, hearing loss, traumatic brain injury, and alcohol misuse) and later life (smoking, depression, physical inactivity, social isolation, diabetes, and air pollution) can contribute to increased dementia risk (table 1). Good evidence exists for all ..."
Lancet Commission 2020 (2020)

This is a research estimate from a model trained on volunteers in the NACC research cohort. It is not a diagnosis, and it does not say what treatment would change the risk. The factors listed describe what the model used, not causes.

Evidence Assistant (RAG)



- MiniLM embeddings + BM25 keyword search, merged by rank fusion.
- If nothing is similar enough, it says the library has no evidence.

Test (11 papers, 447 passages)	Result
Correct source first	32 / 32
Personal-advice questions flagged	8 / 8
Out-of-scope questions declined	6 / 6

Independent Verification

Check	Result
No person or held-out centre in more than one partition	0 shared
Features unchanged when all later visits are deleted	Identical (1,500 people × 67 features)
No PET value newer than the visit that uses it	0 of 1,481 amyloid, 713 tau
Labels recomputed independently	0 disagreements in 11,403 visits
Shuffled labels	AUROC 0.508 (chance)
Retrain with a new seed	0.937 / 0.946 (reported 0.938 / 0.946)

Errors the checks found, and we fixed

Education and race filled from a later visit (0.1% of people); PET scans matched to visits weeks before the scan. After fixing, every model was retrained; results moved only in the third decimal. 17 of 17 checks pass.

What Each Research Question Found

Research question	Answer
RQ1 Prediction accuracy	Dementia 0.94 (0.84 in MCI); decline 0.86 (0.75 in MCI)
RQ2 Cardiovascular history	+0.001 AUROC once cognition is known
RQ3 Amyloid and tau PET	Strong association, no demonstrable gain yet
RQ4 Deep models / fusion net	No gain over logistic regression or XGBoost
RQ5 Earlier visits	+0.000 to +0.005 AUROC
RQ6 Missing categories	Modality dropout keeps calibration close to specialist models
RQ7 Explanation and evidence	SHAP, plausibility checks, report with quoted evidence

Final takeaway

Current cognitive and functional status carries most of the three-year signal; vascular and genetic history add little once it is known.

Limitations & Future Work

- Held-out centres are still within NACC; validation on another cohort is the next step.
- NACC is a research cohort, mostly White and highly educated; about half of eligible people lacked a 3-year outcome.
- Few cognitively normal converters; normal cognition needs a 5- to 10-year model.
- Richer vascular data (repeated blood pressure, treatment, imaging) may capture what yes/no history misses.
- PET registered without MRI and tau not quantified; longer follow-up will allow a proper PET test.
- Evidence library of 11 papers; a larger curated library would help.

Industry / Research Relevance

- **Clinical screening:** flags elevated short-term risk for earlier specialist follow-up, with reasons and evidence, without making a diagnosis.
- **Clinical trials:** within MCI the model ranks risk at AUROC 0.84, which helps select people likely to progress.
- **Research:** a reusable method for measuring what each data source adds and for checking a pipeline against information leakage.

Team Roles

Members	Responsibility
Bhargav, Hansika	Cardiovascular and demographic features, baseline models
Akhil, Bhargav	Visit-history models, CDR-SB progression regression
Mahidhar, Hansika, Bhargav	PET image pipeline and 3D CNN, modality-encoder fusion network
Akhil, Hansika, Mahidhar	RAG evidence assistant, evaluation framework, ablations

References

[1] G. Livingston et al., "Dementia prevention, intervention, and care: 2020 report of the Lancet Commission," The Lancet, vol. 396, pp. 413–446, 2020.

[2] S. A. Alaka et al., "Enhancing the validity of CAIDE dementia risk scores with resting heart rate and machine learning: an analysis from the National Alzheimer's Coordinating Center," Alzheimer's & Dementia, vol. 21, e70442, 2025.

[3] A. Stern and M. Linial, "Integrative machine learning approach to risk prediction for dementia and Alzheimer's disease," GeroScience, vol. 48, pp. 3007–3023, 2026.

[4] D. Aschwanden et al., "Predicting cognitive impairment and dementia: a machine learning approach," J. Alzheimer's Disease, vol. 75, pp. 717–728, 2020.

[5] F. Yi et al., "XGBoost-SHAP-based interpretable diagnostic framework for Alzheimer's disease," BMC Med. Inform. Decis. Mak., vol. 23, 137, 2023.

[6] Y. Gorishniy et al., "Revisiting deep learning models for tabular data," NeurIPS, vol. 34, 2021.

[7] P. Lewis et al., "Retrieval-augmented generation for knowledge-intensive NLP tasks," NeurIPS, vol. 33, 2020.

[8] T. Chen and C. Guestrin, "XGBoost: a scalable tree boosting system," Proc. ACM SIGKDD, pp. 785–794, 2016.

[9] S. M. Lundberg and S.-I. Lee, "A unified approach to interpreting model predictions," NeurIPS, vol. 30, 2017.

[10] W. E. Klunk et al., "The Centiloid Project: standardizing quantitative amyloid plaque estimation by PET," Alzheimer's & Dementia, vol. 11, pp. 1–15, 2015.

AMRITA VISHWA VIDYAPEETHAM
DEEMED TO BE UNIVERSITY UNDER SECTION 3 OF UGC ACT, 1956

SCHOOL OF
ARTIFICIAL INTELLIGENCE
COIMBATORE

Thank You

Questions are welcome.

Data: National Alzheimer's Coordinating Center (NACC), Freeze 74, used under the NACC data use agreement. All results are aggregates.