





gas = [1, 2, 3, 4, 5] - cost = [3, 4, 5, 1, 2]

(15) >= (15)

(14) (15)



[totalGas = 0
totalCost = 0]

→ if (totalGas < totalCost) return -1

start = 0
tank = 0

for (i = 0; i < n; i++) {
 tank += gas[i] - cost[i];
 if (tank < 0) {
 start = i + 1;
 tank = 0;
 }

[1, 2, 3, 4, 5]
[3, 4, 5, 1, 2]

-2 = 1 - 3
3 = 4 - 1

$\left. \begin{array}{l} \text{start} = 1 + 1 \\ \text{tank} = 0 \end{array} \right\}$

3 = 4

}

return start.

896 → monotonic array

[1, 2, 2, 3, 4] → true

[5, 4, 2, 1] → true

[1, 3, 2] → false

= dec || inc

3, 4, 5

$3 < 4 \rightarrow \text{dec} = \text{false}$

4. median of two sorted array

nums1 = [1, 3] , nums2 = [2]
m = 1

(n) = 2

$n + m$
= (3)

1	2	3
---	---	---


 → 3/2 → 1.5
0 1 2

$[1, 3, 5] \rightarrow n$ $[2, 4, 6] \rightarrow m$

$n = 3$
 $m = 3$

$\text{int newArr}[] = \text{new int}[\text{length}]$

$\text{length} = n + m$



① < 2
3 2
3 4
5 4

```
while (i < n & j < m) {
    if (nums1[i] < nums2[j]) {
        newArr[k] = nums1[i]
        i++
    } else {
        newArr[k] = nums2[j]
        j++
    }
    k++
}
```

```
while (i < n) {
    // empty box
}
```

```
while (j < m) {
    // empty box
}
```

③

1, 2, 3
1, [2, 3], 4

5/2

②.5

1, ②, ③, 4
0 1 2

4/2

2 + 3

$$\begin{array}{l} 1/ - \\ \textcircled{2} - 1 \\ \textcircled{2 + 3} \end{array}$$