

Matrix Algebra: Determinants

Vũ Nguyễn Sơn Tùng

Faculty of information technology
HANU

Two-by-two determinant

Let matrix A be 2×2 matrix. Then

$$\det A = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc.$$

is called the determinant of the 2×2 matrix A .

A is invertible if and only if its determinant is nonzero.

If A is invertible, the equation $Ax = b$ has the unique solution $x = A^{-1}b$.

If A is not invertible, then $Ax = b$ may have no solution or an infinite number of solutions.

When $\det A = 0$, we say that the matrix A is *singular*.

Three-by-three determinant

It is also straightforward to define the determinant for a 3×3 matrix. Let matrix A be 3×3 matrix. Consider the system of equations $Ax = 0$ and determine the condition for which $x = 0$ is the only solution:

$$\begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0.$$

We can do the messy algebra of elimination to solve for x_1 , x_2 , and x_3 , then find that $x_1 = x_2 = x_3 = 0$ is the only solution when $\det A = 0$, where the definition, apart from a constant, is given by

$$\det A = aei + bfg + cdh - ceg - bdi - afh.$$

An easy way to remember we can use the Rule of Sarrus (See details in the next slide).

$$\det A = aei + bfg + cdh - ceg - bdi - afh.$$

An easy way to remember we can use the **Rule of Sarrus** (the French mathematician Pierre Frederic Sarrus). Determinant of A can be computed by the following scheme.

$$\begin{pmatrix} a & b & c & a & b \\ d & e & f & d & e \\ g & h & i & g & h \end{pmatrix} - \begin{pmatrix} a & b & c & a & b \\ d & e & f & d & e \\ g & h & i & g & h \end{pmatrix}.$$

Write out the first two columns of the matrix to the right of the third column, giving five columns in a row. Then add the products of the diagonals going from top to bottom and subtract the products of the diagonals going from bottom to top.

$$\det A = aei + bfg + cdh - ceg - bdi - afh.$$

An easy way to remember we can use the **Rule of Sarrus** (the French mathematician Pierre Frederic Sarrus). Determinant of A can be computed by the following scheme.

$$\begin{pmatrix} a & b & c & a & b \\ d & e & f & d & e \\ g & h & i & g & h \end{pmatrix} - \begin{pmatrix} a & b & c & a & b \\ d & e & f & d & e \\ g & h & i & g & h \end{pmatrix}.$$

Write out the first two columns of the matrix to the right of the third column, giving five columns in a row. Then add the products of the diagonals going from top to bottom and subtract the products of the diagonals going from bottom to top.

Question: Can the determinant of a 4×4 matrix be calculated with the Rule of Sarrus?

$$\det A = aei + bfg + cdh - ceg - bdi - afh.$$

An easy way to remember we can use the **Rule of Sarrus** (the French mathematician Pierre Frederic Sarrus). Determinant of A can be computed by the following scheme.

$$\begin{pmatrix} a & b & c & a & b \\ d & e & f & d & e \\ g & h & i & g & h \end{pmatrix} - \begin{pmatrix} a & b & c & a & b \\ d & e & f & d & e \\ g & h & i & g & h \end{pmatrix}.$$

Write out the first two columns of the matrix to the right of the third column, giving five columns in a row. Then add the products of the diagonals going from top to bottom and subtract the products of the diagonals going from bottom to top.

Question: Can the determinant of a 4×4 matrix be calculated with the Rule of Sarrus?

Answer: Unfortunately, this mnemonic only works for three-by-three matrices.

Problems

1. Find the determinant of the three-by-three identity matrix.
2. Show that the three-by-three determinant changes sign when the first two rows are interchanged.
3. Let A and B be two-by-two matrices. Prove by direct computation that $\det AB = \det A \det B$.

Laplace expansion

There is a way to write the three-by-three determinant that generalizes. It is called a **Laplace expansion** (also called a cofactor expansion or expansion by minors):

$$\det A = \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix}$$

$$= aei + bfg + cdh - ceg - bdi - afh = a(ei - fh) - b(di - fg) + c(dh - eg)$$

$$= a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix} = a \det A_{11} - b \det A_{12} + c \det A_{13},$$

where A_{11} , A_{12} and A_{13} are obtained from A by deleting the first row and one of the three columns.

Evidently, the 3×3 determinant can be computed from lower-order 2×2 determinants, called minors. The rule here for a general $n \times n$ matrix is that one goes across the first row of the matrix, multiplying each element in the row by the determinant of the matrix obtained by crossing out that element's row and column, and adding the results with alternating signs.

In fact, this expansion in minors can be done across any row or down any column. When the minor A_{ij} is obtained by deleting the i th row and j th column, then the sign of the term is given by $(-1)^{i+j}$.

An easy way to remember the signs is to form a checkerboard pattern, exhibited here for the 3×3 and 4×4 matrices:

$$\begin{pmatrix} + & - & + \\ - & + & - \\ + & - & + \end{pmatrix}, \quad \begin{pmatrix} + & - & + & - \\ - & + & - & + \\ + & - & + & - \\ - & + & - & + \end{pmatrix}.$$

In fact, this expansion in minors can be done across any row or down any column. When the minor A_{ij} is obtained by deleting the i th row and j th column, then the sign of the term is given by $(-1)^{i+j}$.

An easy way to remember the signs is to form a checkerboard pattern, exhibited here for the 3×3 and 4×4 matrices:

$$\begin{pmatrix} + & - & + \\ - & + & - \\ + & - & + \end{pmatrix}, \quad \begin{pmatrix} + & - & + & - \\ - & + & - & + \\ + & - & + & - \\ - & + & - & + \end{pmatrix}.$$

For example: Compute the determinant of matrix

$$A = \begin{pmatrix} 1 & 5 & 0 \\ 2 & 4 & -1 \\ 0 & -2 & 0 \end{pmatrix}$$

In fact, this expansion in minors can be done across any row or down any column. When the minor A_{ij} is obtained by deleting the i th row and j th column, then the sign of the term is given by $(-1)^{i+j}$.

An easy way to remember the signs is to form a checkerboard pattern, exhibited here for the 3×3 and 4×4 matrices:

$$\begin{pmatrix} + & - & + \\ - & + & - \\ + & - & + \end{pmatrix}, \quad \begin{pmatrix} + & - & + & - \\ - & + & - & + \\ + & - & + & - \\ - & + & - & + \end{pmatrix}.$$

For example: Compute the determinant of matrix

$$A = \begin{pmatrix} 1 & 5 & 0 \\ 2 & 4 & -1 \\ 0 & -2 & 0 \end{pmatrix}$$

SOLUTION Compute $\det A = a_{11} \det A_{11} - a_{12} \det A_{12} + a_{13} \det A_{13}$:

$$\begin{aligned} \det A &= 1 \cdot \det \begin{bmatrix} 4 & -1 \\ -2 & 0 \end{bmatrix} - 5 \cdot \det \begin{bmatrix} 2 & -1 \\ 0 & 0 \end{bmatrix} + 0 \cdot \det \begin{bmatrix} 2 & 4 \\ 0 & -2 \end{bmatrix} \\ &= 1(0 - 2) - 5(0 - 0) + 0(-4 - 0) = -2 \end{aligned}$$



Definition of a determinant

We can now give a recursive definition of a determinant:

Definition of a determinant

We can now give a recursive definition of a determinant:

*For $n \geq 2$, the **determinant** of an $n \times n$ matrix $A = [a_{ij}]$ is the sum of n terms of the form $\pm a_{ij} \det A_{1j}$, with plus and minus signs alternating, where the entries $a_{11}, a_{12}, \dots, a_{1n}$ are from the first row of A . In symbols,*

$$\det A = a_{11} \det A_{11} - a_{12} \det A_{12} + \dots + (-1)^{1+n} a_{1n} \det A_{1n}.$$

Cofactor expansion

Given $A = [a_{ij}]$, the (i, j) -cofactor of A is the number C_{ij} given by

$$C_{ij} = (-1)^{i+j} \det A_{ij}.$$

We omit the proof of the following fundamental theorem:

Cofactor expansion

Given $A = [a_{ij}]$, the (i, j) -cofactor of A is the number C_{ij} given by

$$C_{ij} = (-1)^{i+j} \det A_{ij}.$$

We omit the proof of the following fundamental theorem:

Theorem

The determinant of an $n \times n$ matrix A can be computed by a cofactor expansion across any row or down any column. The expansion across the i th row is

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \dots + a_{in}C_{in}.$$

The cofactor expansion down the j th column is

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \dots + a_{nj}C_{nj}.$$

Cofactor expansion

Given $A = [a_{ij}]$, the (i, j) -cofactor of A is the number C_{ij} given by

$$C_{ij} = (-1)^{i+j} \det A_{ij}.$$

We omit the proof of the following fundamental theorem:

Theorem

The determinant of an $n \times n$ matrix A can be computed by a cofactor expansion across any row or down any column. The expansion across the i th row is

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \dots + a_{in}C_{in}.$$

The cofactor expansion down the j th column is

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \dots + a_{nj}C_{nj}.$$

The formula in the theorem above is called a **cofactor expansion across the first row** of A .

Cofactor expansion

Given $A = [a_{ij}]$, the (i, j) -cofactor of A is the number C_{ij} given by

$$C_{ij} = (-1)^{i+j} \det A_{ij}.$$

We omit the proof of the following fundamental theorem:

Theorem

The determinant of an $n \times n$ matrix A can be computed by a cofactor expansion across any row or down any column. The expansion across the i th row is

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \dots + a_{in}C_{in}.$$

The cofactor expansion down the j th column is

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \dots + a_{nj}C_{nj}.$$

The formula in the theorem above is called a **cofactor expansion across the first row** of A .

Note: If A is a triangular matrix, then $\det A$ is the product of the entries on the main diagonal of A .

Basic properties of determinants

The secret of determinants lies in how they change when row operations are performed. The following theorem is useful to find determinants efficiently.

Basic properties of determinants

The secret of determinants lies in how they change when row operations are performed. The following theorem is useful to find determinants efficiently.

Theorem

Let A be a square matrix.

- (a) If a multiple of one row of A is added to another row to produce a matrix B , then $\det B = \det A$.*
- (b) If two rows of A are interchanged to produce B , then $\det B = -\det A$.*
- (b) If one row of A is multiplied by k to produce B , then $\det B = k \det A$*

Basic properties of determinants

The secret of determinants lies in how they change when row operations are performed. The following theorem is useful to find determinants efficiently.

Theorem

Let A be a square matrix.

- (a) If a multiple of one row of A is added to another row to produce a matrix B , then $\det B = \det A$.*
- (b) If two rows of A are interchanged to produce B , then $\det B = -\det A$.*
- (b) If one row of A is multiplied by k to produce B , then $\det B = k \det A$*

Note: We can perform operations on the columns of a matrix in a way that is analogous to the row operations we have considered.

The next theorem shows that column operations have the same effects on determinants as row operations:

Basic properties of determinants

The secret of determinants lies in how they change when row operations are performed. The following theorem is useful to find determinants efficiently.

Theorem

Let A be a square matrix.

- (a) If a multiple of one row of A is added to another row to produce a matrix B , then $\det B = \det A$.*
- (b) If two rows of A are interchanged to produce B , then $\det B = -\det A$.*
- (b) If one row of A is multiplied by k to produce B , then $\det B = k \det A$*

Note: We can perform operations on the columns of a matrix in a way that is analogous to the row operations we have considered.

The next theorem shows that column operations have the same effects on determinants as row operations:

Theorem

If A is an $n \times n$ matrix, then $\det A^T = \det A$.

Example:

Compute $\det A$, where

$$A = \begin{bmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{bmatrix}.$$

Example:

Compute $\det A$, where

$$A = \begin{bmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{bmatrix}.$$

SOLUTION The strategy is to reduce A to echelon form and then to use the fact that the determinant of a triangular matrix is the product of the diagonal entries. The first two row replacements in column 1 do not change the determinant:

$$\det A = \begin{vmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{vmatrix} = \begin{vmatrix} 1 & -4 & 2 \\ 0 & 0 & -5 \\ -1 & 7 & 0 \end{vmatrix} = \begin{vmatrix} 1 & -4 & 2 \\ 0 & 0 & -5 \\ 0 & 3 & 2 \end{vmatrix}$$

An interchange of rows 2 and 3 reverses the sign of the determinant, so

$$\det A = - \begin{vmatrix} 1 & -4 & 2 \\ 0 & 3 & 2 \\ 0 & 0 & -5 \end{vmatrix} = -(1)(3)(-5) = 15$$

A common use of Theorem 3(c) in hand calculations is to *factor out a common multiple of one row* of a matrix. For instance,

$$\begin{vmatrix} * & * & * \\ 5k & -2k & 3k \\ * & * & * \end{vmatrix} = k \begin{vmatrix} * & * & * \\ 5 & -2 & 3 \\ * & * & * \end{vmatrix}$$

where the starred entries are unchanged. We use this step in the next example.

The main theorem

$$U = \begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & \blacksquare \end{bmatrix} \quad \det U \neq 0$$
$$U = \begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & 0 & \blacksquare \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \det U = 0$$

FIGURE 1

Typical echelon forms of square matrices.

Suppose a square matrix A has been reduced to an echelon form U by row replacements and row interchanges. (This is always possible. See the row reduction algorithm in Section 1.2.) If there are r interchanges, then Theorem 3 shows that

$$\det A = (-1)^r \det U$$

Since U is in echelon form, it is triangular, and so $\det U$ is the product of the diagonal entries u_{11}, \dots, u_{nn} . If A is invertible, the entries u_{ii} are all pivots (because $A \sim I_n$ and the u_{ii} have not been scaled to 1's). Otherwise, at least u_{nn} is zero, and the product $u_{11} \cdots u_{nn}$ is zero. See Figure 1. Thus

$$\det A = \begin{cases} (-1)^r \cdot \left(\text{product of} \right. \\ \quad \left. \text{pivots in } U \right) & \text{when } A \text{ is invertible} \\ 0 & \text{when } A \text{ is not invertible} \end{cases} \quad (1)$$

It is interesting to note that although the echelon form U described above is not unique (because it is not completely row reduced), and the pivots are not unique, the *product* of the pivots *is* unique, except for a possible minus sign.

Formula (1) not only gives a concrete interpretation of $\det A$ but also proves the main theorem of this section:

Theorem

A square matrix A is invertible if and only if $\det A \neq 0$.

This theorem adds the statement $\det A \neq 0$ to the Invertible Matrix Theorem.

Theorem (Multiplicative property)

If A and B are $n \times n$ matrices, then $\det(AB) = \det A \cdot \det B$.

Determinants and Matrix Products

Theorem (Multiplicative property)

If A and B are $n \times n$ matrices, then $\det(AB) = \det A \cdot \det B$.

Example: Compute $\det AB$, where

$$A = \begin{bmatrix} 6 & 1 \\ 3 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}.$$

Determinants and Matrix Products

Theorem (Multiplicative property)

If A and B are $n \times n$ matrices, then $\det(AB) = \det A \cdot \det B$.

Example: Compute $\det AB$, where

$$A = \begin{bmatrix} 6 & 1 \\ 3 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}.$$

SOLUTION

$$AB = \begin{bmatrix} 6 & 1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 25 & 20 \\ 14 & 13 \end{bmatrix}$$

and

$$\det AB = 25 \cdot 13 - 20 \cdot 14 = 325 - 280 = 45$$

Since $\det A = 9$ and $\det B = 5$,

$$(\det A)(\det B) = 9 \cdot 5 = 45 = \det AB$$



A Linearity Property

For an $n \times n$ matrix A , we can consider $\det A$ as a function of the n column vectors in A .

The determinant is a linear function of any row, holding all other rows fixed.

A Linearity Property

For an $n \times n$ matrix A , we can consider $\det A$ as a function of the n column vectors in A .

The determinant is a linear function of any row, holding all other rows fixed.

For example:

$$\begin{vmatrix} a + a' & b + b' \\ c & d \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} + \begin{vmatrix} a' & b' \\ c & d \end{vmatrix}.$$

This linearity property of the determinant turns out to have many useful consequences that are studied in more advanced courses.

Useful properties

- The determinant of the identity matrix is one;
- The determinant changes sign under row interchange;
- The determinant is a linear function of any row, holding all other rows fixed;
- If a matrix has two equal rows, then the determinant is zero;
- If we add k times row- i to row- j , the determinant doesn't change;
- The determinant of a matrix with a row of zeros is zero;
- A matrix with a zero determinant is not invertible;
- The determinant of a diagonal matrix is the product of the diagonal elements;
- The determinant of an upper or lower triangular matrix is the product of the diagonal elements;
- The determinant of the product of two matrices is equal to the product of the determinants;
- The determinant of the inverse matrix is equal to the reciprocal of the determinant;
- The determinant of the transpose of a matrix is equal to the determinant of the matrix.

Cramer's Rule


Cramer's rule is needed in a variety of theoretical calculations. For instance, it can be used to study how the solution of $A\mathbf{x} = \mathbf{b}$ is affected by changes in the entries of \mathbf{b} .

Cramer's Rule

Cramer's rule is needed in a variety of theoretical calculations. For instance, it can be used to study how the solution of $A\mathbf{x} = \mathbf{b}$ is affected by changes in the entries of \mathbf{b} .

For any $n \times n$ matrix A and any \mathbf{b} in \mathbb{R}^n , let $A_i(\mathbf{b})$ be the matrix obtained from A by replacing column i by the vector \mathbf{b} .

$$A_i(\mathbf{b}) = [\mathbf{a}_1 \quad \cdots \quad \mathbf{b} \quad \cdots \quad \mathbf{a}_n]$$


col i

Cramer's Rule

Cramer's rule is needed in a variety of theoretical calculations. For instance, it can be used to study how the solution of $A\mathbf{x} = \mathbf{b}$ is affected by changes in the entries of \mathbf{b} .

For any $n \times n$ matrix A and any \mathbf{b} in \mathbb{R}^n , let $A_i(\mathbf{b})$ be the matrix obtained from A by replacing column i by the vector \mathbf{b} .

$$A_i(\mathbf{b}) = [\mathbf{a}_1 \quad \cdots \quad \mathbf{b} \quad \cdots \quad \mathbf{a}_n]$$

\uparrow
col i

Theorem (Cramer's Rule)

Let A be an invertible $n \times n$ matrix. For any $\mathbf{b} \in \mathbb{R}^n$, the unique solution \mathbf{x} of $A\mathbf{x} = \mathbf{b}$ has entries given by

$$x_i = \frac{\det A_i(\mathbf{b})}{\det A}.$$

Example:

Use Cramer's rule to solve the system

$$3x_1 - 2x_2 = 6; \quad -5x_1 + 4x_2 = 8.$$

Example:

Use Cramer's rule to solve the system

$$3x_1 - 2x_2 = 6; \quad -5x_1 + 4x_2 = 8.$$

SOLUTION View the system as $A\mathbf{x} = \mathbf{b}$. Using the notation introduced above,

$$A = \begin{bmatrix} 3 & -2 \\ -5 & 4 \end{bmatrix}, \quad A_1(\mathbf{b}) = \begin{bmatrix} 6 & -2 \\ 8 & 4 \end{bmatrix}, \quad A_2(\mathbf{b}) = \begin{bmatrix} 3 & 6 \\ -5 & 8 \end{bmatrix}$$

Since $\det A = 2$, the system has a unique solution. By Cramer's rule,

$$x_1 = \frac{\det A_1(\mathbf{b})}{\det A} = \frac{24 + 16}{2} = 20$$

$$x_2 = \frac{\det A_2(\mathbf{b})}{\det A} = \frac{24 + 30}{2} = 27$$



Formula for the inverse matrix

Cramer's rule leads easily to a general formula for the inverse of an $n \times n$ matrix A . The j th column of A^{-1} is \mathbf{x} that satisfies $A\mathbf{x} = \mathbf{e}_j$, where \mathbf{e}_j is the j th column of the identity matrix, and the i th entry of \mathbf{x} is the (i, j) -entry of A^{-1} .

Let A_{ji} be the submatrix of A formed by deleting row j and column i . Then

$$\det A_i(\mathbf{e}_j) = (-1)^{i+j} \det A_{ji} = C_{ji},$$

where C_{ji} is a cofactor of A .

Formula for the inverse matrix

Cramer's rule leads easily to a general formula for the inverse of an $n \times n$ matrix A . The j th column of A^{-1} is \mathbf{x} that satisfies $A\mathbf{x} = \mathbf{e}_j$, where \mathbf{e}_j is the j th column of the identity matrix, and the i th entry of \mathbf{x} is the (i, j) -entry of A^{-1} .

Let A_{ji} be the submatrix of A formed by deleting row j and column i . Then

$$\det A_i(\mathbf{e}_j) = (-1)^{i+j} \det A_{ji} = C_{ji},$$

where C_{ji} is a cofactor of A . By Cramer's rule we have

$$\{(i, j)\text{-entry of } A^{-1}\} = x_i = \frac{\det A_i(\mathbf{e}_j)}{\det A},$$

$$A^{-1} = \frac{1}{\det A} \begin{bmatrix} C_{11} & C_{21} & \cdots & C_{n1} \\ C_{12} & C_{22} & \cdots & C_{n2} \\ \vdots & \vdots & & \vdots \\ C_{1n} & C_{2n} & \cdots & C_{nn} \end{bmatrix}.$$

Formula for the inverse matrix

Cramer's rule leads easily to a general formula for the inverse of an $n \times n$ matrix A . The j th column of A^{-1} is \mathbf{x} that satisfies $A\mathbf{x} = \mathbf{e}_j$, where \mathbf{e}_j is the j th column of the identity matrix, and the i th entry of \mathbf{x} is the (i, j) -entry of A^{-1} .

Let A_{ji} be the submatrix of A formed by deleting row j and column i . Then

$$\det A_i(\mathbf{e}_j) = (-1)^{i+j} \det A_{ji} = C_{ji},$$

where C_{ji} is a cofactor of A . By Cramer's rule we have

$$\{(i, j)\text{-entry of } A^{-1}\} = x_i = \frac{\det A_i(\mathbf{e}_j)}{\det A},$$

$$A^{-1} = \frac{1}{\det A} \begin{bmatrix} C_{11} & C_{21} & \cdots & C_{n1} \\ C_{12} & C_{22} & \cdots & C_{n2} \\ \vdots & \vdots & & \vdots \\ C_{1n} & C_{2n} & \cdots & C_{nn} \end{bmatrix}.$$

The matrix of cofactors on the right side of the formula above is called the **adjugate** (or **classical adjoint**) of A , denoted by $\text{adj } A$.

Formula for the inverse matrix

Cramer's rule leads easily to a general formula for the inverse of an $n \times n$ matrix A . The j th column of A^{-1} is \mathbf{x} that satisfies $A\mathbf{x} = \mathbf{e}_j$, where \mathbf{e}_j is the j th column of the identity matrix, and the i th entry of \mathbf{x} is the (i, j) -entry of A^{-1} .

Let A_{ji} be the submatrix of A formed by deleting row j and column i . Then

$$\det A_i(\mathbf{e}_j) = (-1)^{i+j} \det A_{ji} = C_{ji},$$

where C_{ji} is a cofactor of A . By Cramer's rule we have

$$\{(i, j)\text{-entry of } A^{-1}\} = x_i = \frac{\det A_i(\mathbf{e}_j)}{\det A},$$

$$A^{-1} = \frac{1}{\det A} \begin{bmatrix} C_{11} & C_{21} & \cdots & C_{n1} \\ C_{12} & C_{22} & \cdots & C_{n2} \\ \vdots & \vdots & & \vdots \\ C_{1n} & C_{2n} & \cdots & C_{nn} \end{bmatrix}.$$

The matrix of cofactors on the right side of the formula above is called the **adjugate** (or **classical adjoint**) of A , denoted by $\text{adj } A$.

Theorem (An inverse formula)

Let A be an invertible $n \times n$ matrix. Then

$$A^{-1} = \frac{1}{\det A} \text{adj } A.$$

Methods to Find Inverse of Matrix

The inverse of a matrix can be found using two methods:

- (1) By elementary row or column operations (Gauss-Jordan method);
- (2) By inverse of matrix formula (using the adjoint and determinant of matrix).

Finding the inverse matrix using inverse formula

The inverse of matrix A can be computed using the inverse of matrix formula:

$$A^{-1} = \frac{1}{\det A} \operatorname{adj} A,$$

i.e., by dividing the adjoint of a matrix by the determinant of the matrix. The inverse of a matrix can be calculated by following the given steps:

- Step 1. Calculate the minors of all elements of A ;
- Step 2. Then compute the cofactors of all elements and write the cofactor matrix by replacing the elements of A by their corresponding cofactors.
- Step 3. Find the adjoint of A (written as $\operatorname{adj} A$) by taking the transpose of the cofactor matrix of A .
- Step 4. Multiply $\operatorname{adj} A$ by the reciprocal of the determinant.

Example

Find the inverse of the matrix A :

$$\begin{bmatrix} 3 & 0 & 2 \\ 2 & 0 & -2 \\ 0 & 1 & 1 \end{bmatrix}.$$

Example

Find the inverse of the matrix A :

$$\begin{bmatrix} 3 & 0 & 2 \\ 2 & 0 & -2 \\ 0 & 1 & 1 \end{bmatrix}.$$

Solution. It needs 4 steps. It is all simple arithmetic but there is a lot of it, so try not to make a mistake!

Step 1. Matrix of minors.

The first step is to create a "Matrix of Minors", where each minor A_{ji} formed by deleting row j and column i . Here are the first two, and last two, calculations of the "Matrix of Minors" :

$$\begin{bmatrix} \bullet & 0 & 2 \\ 2 & 0 & -2 \\ 0 & 1 & 1 \end{bmatrix} \quad 0 \times 1 - (-2) \times 1 = 2$$
$$\begin{bmatrix} 3 & \bullet & 2 \\ 2 & 0 & -2 \\ 0 & 1 & 1 \end{bmatrix} \quad 2 \times 1 - (-2) \times 0 = 2$$

...

$$\begin{bmatrix} 3 & 0 & 2 \\ 2 & 0 & -2 \\ 0 & \bullet & 1 \end{bmatrix} \quad 3 \times -2 - 2 \times 2 = -10$$
$$\begin{bmatrix} 3 & 0 & 2 \\ 2 & 0 & -2 \\ 0 & 1 & \bullet \end{bmatrix} \quad 3 \times 0 - 0 \times 2 = 0$$

And here is the calculation for the whole matrix:

$$\begin{bmatrix} 0 \times 1 - (-2) \times 1 & 2 \times 1 - (-2) \times 0 & 2 \times 1 - 0 \times 0 \\ 0 \times 1 - 2 \times 1 & 3 \times 1 - 2 \times 0 & 3 \times 1 - 0 \times 0 \\ 0 \times (-2) - 2 \times 0 & 3 \times (-2) - 2 \times 2 & 3 \times 0 - 0 \times 2 \end{bmatrix} = \begin{bmatrix} 2 & 2 & 2 \\ -2 & 3 & 3 \\ 0 & -10 & 0 \end{bmatrix}$$

Matrix of Minors

Step 2. Matrix of Cofactors.

Just apply a "checkerboard" of minuses to the "Matrix of Minors". In other words, we need to change the sign of alternate cells, like this:

$$\begin{array}{ccc} \begin{bmatrix} 2 & 2 & 2 \\ -2 & 3 & 3 \\ 0 & -10 & 0 \end{bmatrix} & \rightsquigarrow & \begin{bmatrix} + & - & + \\ - & + & - \\ + & - & + \end{bmatrix} \\ \text{Matrix of Minors} & & \end{array}$$

$$\begin{array}{ccc} & & \rightsquigarrow & \begin{bmatrix} 2 & -2 & 2 \\ +2 & 3 & -3 \\ 0 & +10 & 0 \end{bmatrix} \\ & & \text{Matrix of CoFactors} \end{array}$$

In general, the cofactor of A is $C_{ji} = (-1)^{i+j} \det A_{ji}$.

Step 3. Adjugate of A (also called Adjoint).

Now "Transpose" all elements of the previous matrix. In other words swap their positions over the diagonal (the diagonal stays the same):

$$\begin{bmatrix} 2 & 2 & 0 \\ -2 & 3 & 10 \\ 2 & -3 & 0 \end{bmatrix}$$

Step 4. Multiply by $\frac{1}{|A|}$.

Now find the determinant of the original matrix. This isn't too hard, because we already calculated the determinants of the smaller parts when we did "Matrix of Minors".

Using: Elements of top row: 3, 0, 2 and Minors for top row: 2, 2, 2, we end up with

$$\det A = 3 \cdot 2 - 0 \cdot 2 + 2 \cdot 2 = 10.$$

Now we have:

$$A^{-1} = \frac{1}{10} \begin{bmatrix} 2 & 2 & 0 \\ -2 & 3 & 10 \\ 2 & -3 & 0 \end{bmatrix} = \begin{bmatrix} 0.2 & 0.2 & 0 \\ -0.2 & 0.3 & 1 \\ 0.2 & -0.3 & 0 \end{bmatrix}$$

Adjugate *Inverse*

THANK YOU FOR YOUR ATTENTION!