

Quotient Category and Sub-Category

A short story

Hao Fan

Faculty of Mathematics
Undergraduate

Course Report, March 2023



Table of Contents

1 Quotient Category

2 Sub-Category



We have studied a special category (\mathbf{hTop}) , defined as follows:

$$\begin{aligned}\mathrm{Ob}(\mathbf{hTop}) &:= \mathrm{Ob}(\mathbf{Top}) \\ \mathrm{Hom}_{(\mathbf{hTop})}(X, Y) &:= \mathrm{Hom}_{(\mathbf{Top})}(X, Y) / \simeq,\end{aligned}$$

where \simeq satisfies:

$$f_0 \simeq f_1, g_0 \simeq g_1 \implies g_0 \circ f_0 \simeq g_1 \circ f_1.$$



Table of Contents

1 Quotient Category

2 Sub-Category



Quotient Category

As a generalization:

Definition

Let a category \mathcal{C} and (a family of) equivalence relations $\sim^{A,B}$ on each set $\text{Hom}_{\mathcal{C}}(A, B)$, where $A, B \in \text{Ob}(\mathcal{C})$ be given. We define a new category \mathcal{C}/\sim as follows:

$$\text{Ob}(\mathcal{C}/\sim) := \text{Ob}(\mathcal{C}),$$

$$\text{Hom}_{\mathcal{C}/\sim}(A, B) := \text{Hom}_{\mathcal{C}}(A, B)/\sim^{A,B}.$$

where \simeq satisfies: $f_0 \simeq f_1, g_0 \simeq g_1 \implies g_0 \circ f_0 \simeq g_1 \circ f_1$. It automatically has the identity and associativity holds.



Quotient Category

Remark

There is a "natural" full functor: $Q: \mathcal{C} \rightarrow \mathcal{C}/\sim$.



Quotient Category

Remark

There is a "natural" full functor: $Q: \mathcal{C} \rightarrow \mathcal{C}/\sim$.

Corollary

If $A \cong B$ in the category \mathcal{C} , then $A \cong B$ in \mathcal{C}/\sim .



Examples of Quotient Categories

Example

The category $(\mathbf{hTop}) = (\mathbf{Top}) / \simeq$, where \simeq is the homotopy relation.

Example

The category $(\mathbf{Lin}_{\mathbb{C}})$, with equivalence relations \sim defined as follows: for $f, g \in \mathcal{L}(X, Y)$, $f \sim g \iff \exists k \in \mathbb{C} \setminus \{0\}$ such that $f = kg$. Then we have a quotient category $(\mathbf{Lin}_{\mathbb{C}}) / \sim$.



Table of Contents

1 Quotient Category

2 Sub-Category



Definition

Let a category \mathcal{C} be given. If another category \mathcal{C}' satisfies:

- 1 $\text{Ob}(\mathcal{C}') \subseteq \text{Ob}(\mathcal{C})$;
- 2 $\forall A, B \in \text{Ob}(\mathcal{C}') : \mathcal{C}'(A, B) := \text{Hom}_{\mathcal{C}'}(A, B) \subseteq \mathcal{C}(A, B)$;
- 3 compositions in \mathcal{C}' are birestrictions of compositions in \mathcal{C} , i.e. the following diagram commutes;

$$\begin{array}{ccc} \mathcal{C}'(A, B) \times \mathcal{C}'(B, C) & \longrightarrow & \mathcal{C}'(A, C) \\ \downarrow & & \downarrow \\ \mathcal{C}(A, B) \times \mathcal{C}(B, C) & \longrightarrow & \mathcal{C}(A, C) \end{array}$$

- 4 $\forall A \in \text{Ob}(\mathcal{C}')$, the identity map of A (in \mathcal{C}), i.e. $1_A^{\mathcal{C}}$ lies in $\mathcal{C}'(A, A)$;
- then we say \mathcal{C}' is a sub-category of \mathcal{C} . denoted by $\mathcal{C}' \subset \mathcal{C}$.

Remark

To describe a sub-category of \mathcal{C} , it suffices to tell its object class and morphism sets. Since all the others are determined by \mathcal{C} .



Remark

To describe a sub-category of \mathcal{C} , it suffices to tell its object class and morphism sets. Since all the others are determined by \mathcal{C} .

Remark

There is a "natural" faithful functor $I: \mathcal{C}' \rightarrow \mathcal{C}$.



Sub-Category

Remark

To describe a sub-category of \mathcal{C} , it suffices to tell its object class and morphism sets. Since all the others are determined by \mathcal{C} .

Remark

There is a "natural" faithful functor $I: \mathcal{C}' \rightarrow \mathcal{C}$.

Corollary

If $A \cong B$ in the category \mathcal{C}' , then $A \cong B$ in \mathcal{C} .



Sub-Category

Remark

To describe a sub-category of \mathcal{C} , it suffices to tell its object class and morphism sets. Since all the others are determined by \mathcal{C} .

Remark

There is a "natural" faithful functor $I: \mathcal{C}' \rightarrow \mathcal{C}$.

Corollary

If $A \cong B$ in the category \mathcal{C}' , then $A \cong B$ in \mathcal{C} .

Definition

A sub-category is said to be **full**, if the functor $I: \mathcal{C}' \rightarrow \mathcal{C}$ is full.



Sub-Category

Remark

To describe a sub-category of \mathcal{C} , it suffices to tell its object class and morphism sets. Since all the others are determined by \mathcal{C} .

Remark

There is a "natural" faithful functor $I: \mathcal{C}' \rightarrow \mathcal{C}$.

Corollary

If $A \cong B$ in the category \mathcal{C}' , then $A \cong B$ in \mathcal{C} .

Definition

A sub-category is said to be **full**, if the functor $I: \mathcal{C}' \rightarrow \mathcal{C}$ is full.



The following are sub-categories relevant to our familiar categories:

Examples

- 1 $(\mathbf{FinSet}) \subset (\mathbf{Set})$, it is full;



The following are sub-categories relevant to our familiar categories:

Examples

- 1 $(\mathbf{FinSet}) \subset (\mathbf{Set})$, it is full;
- 2 $(\mathbf{FinLin}_{\mathbb{C}}) \subset (\mathbf{Lin}_{\mathbb{C}})$, it is full;



The following are sub-categories relevant to our familiar categories:

Examples

- 1 $(\mathbf{FinSet}) \subset (\mathbf{Set})$, it is full;
- 2 $(\mathbf{FinLin}_{\mathbb{C}}) \subset (\mathbf{Lin}_{\mathbb{C}})$, it is full;
- 3 $(\mathbf{Ab}) \subset (\mathbf{Grp})$, it is full;



The following are sub-categories relevant to our familiar categories:

Examples

- 1 $(\mathbf{FinSet}) \subset (\mathbf{Set})$, it is full;
- 2 $(\mathbf{FinLin}_{\mathbb{C}}) \subset (\mathbf{Lin}_{\mathbb{C}})$, it is full;
- 3 $(\mathbf{Ab}) \subset (\mathbf{Grp})$, it is full;
- 4 $(\mathbf{Haus}) \subset (\mathbf{Top})$, it is full;



Example

Let a fixed category \mathcal{C} satisfying $\text{Aut}(X, Y) \neq \text{Hom}(X, Y)$ for some $X, Y \in \text{Ob}(\mathcal{C})$ be given. Consider the sub-category $B\mathcal{C}$ defined as follows:

$$\text{Ob}(B\mathcal{C}) := \text{Ob}(\mathcal{C}), \text{Hom}_{B\mathcal{C}}(X, Y) := \text{Isom}_{\mathcal{C}}(X, Y).$$

Then is is a sub-category that is **not** full.

