

Computational aspects of modular parametrizations of elliptic curves

Hao Chen

University of Washington Ph.D. defense

Advisor: William Stein

April 26, 2016

Critical subgroups of elliptic curves

- 1 Critical subgroups of elliptic curves
 - Elliptic curves and modular curves
 - The critical subgroup and critical polynomials
 - Computing the critical subgroup
- 2 q -expansion of newforms at non-unitary cusps
 - Problem description
- 3 Chow-Heegner points
 - Even index
 - Computing Chow-Heegner points

Elliptic curves over \mathbb{Q}

Definition

An **elliptic curve** over \mathbb{Q} is a nonsingular projective curve $E \subseteq \mathbb{P}^2$ with defining equation

$$y^2z = x^3 + Axz^2 + Bz^3,$$

where $A, B \in \mathbb{Q}$ and $4A^3 + 27B^2 \neq 0$.

Theorem (Mordell-Weil)

$E(\mathbb{Q})$ is a finitely generated abelian group, i.e.,

$$E(\mathbb{Q}) \cong \mathbb{Z}^r \oplus T,$$

for some $r \geq 0$ and T finite.

r is called the **rank**. T is the **torsion subgroup**.

The BSD conjecture

There is an entire function $L(E, s)$ called the L -function of E .

The rank part of the **Birch and Swinnerton-Dyer (BSD) conjecture** is

$$\text{rank}(E(\mathbb{Q})) = \text{ord}_{s=1} L(E, s).$$

- $\text{ord}_{s=1} L(E, s)$ is the **analytic rank**, denoted by $r_{an}(E)$.
- The BSD conjecture is open when $r_{an}(E) > 1$.
- The proof of rank BSD for $r_{an}(E) \leq 1$ uses Heegner points.

Modular curves

Let $N \geq 1$ be an integer, consider the group

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z}) : N \mid c \right\}.$$

Let $\mathcal{H}^* := \{z \in \mathbb{C} : \text{im}(z) > 0\} \cup \mathbb{Q} \cup \infty$. $\Gamma_0(N)$ acts on \mathcal{H}^* .

Definition

$X_0(N) = \Gamma_0(N) \backslash \mathcal{H}^*$ is the modular curve of level N .

- $X_0(N)$ is a nonsingular projective curve.

The modularity theorem

Theorem (Modularity (Breuil, Conrad, Taylor fixme))

For every elliptic curve E/\mathbb{Q} , there exists an integer $N > 1$ and a surjective morphism $\varphi : X_0(N) \rightarrow E$ defined over \mathbb{Q} .

The smallest N is called the conductor of E .

Let $\omega = \varphi^*\left(\frac{dx}{y}\right)$. Then $\omega = cf(z)dz$, where f is the **modular form attached to E** .

We assume E is optimal. Then φ is unique up to sign.

Idea: use φ to find points on E from special points on $X_0(N)$.
– rational points on $X_0(N)$ – cusps. – Heegner points. – Ramification points. – Others??
Note: up to now, no known construction in ≥ 2 .

The critical subgroup $E_{crit}(\mathbb{Q})$

Let $R_\varphi \in \text{Div}(X_0(N))$ be the ramification divisor of φ .

Definition (Mazur, Swinnerton-Dyer)

The **critical subgroup** of E is

$$E_{crit}(\mathbb{Q}) = \langle \text{tr}(\varphi([z])) : [z] \in \text{supp } R_\varphi \rangle \subseteq E(\mathbb{Q}),$$

where $\text{tr}(P) = \sum_{\sigma: \mathbb{Q}(P) \rightarrow \bar{\mathbb{Q}}} P^\sigma$.

Question (Mazur and Swinnerton-Dyer, 1974)

Is there an elliptic curve E/\mathbb{Q} with $r_{an}(E) \geq 2$ and $\text{rank}(E_{crit}(\mathbb{Q})) > 0$?

Theorem (C.)

For all elliptic curves E of rank 2 and conductor $N < 1000$, the rank of $E_{\text{crit}}(\mathbb{Q})$ is 0.

To prove this, we compute $E_{\text{crit}}(\mathbb{Q})$ for each curve.

Critical j -polynomial

To help compute $E_{crit}(\mathbb{Q})$, we make the following definition.

Definition

Write $\text{div}(\omega) = \sum n_z[z]$. The **critical j -polynomial** of E is

$$F_{E,j}(x) = \prod_{z \in \text{supp div}(\omega), j(z) \neq \infty} (x - j(z))^{n_z}.$$

$F_{E,j}(x) \in \mathbb{Q}[x]$ and $\deg F_{E,j} \leq 2g - 2$ (equality holds if N is square free).

For $h \in \mathbb{Q}(X_0(N))$, can define $F_{E,h}(x)$.

Let H_d be the Hilbert class polynomial of disc d .

Example

$$F_{44a,j}(x) = H_{-44}(x)^2. \quad F_{37a,j}(x) = H_{-148}(x). \quad F_{37b,j}(x) = H_{-16}(x)^2.$$

Polynomial Relation (I)

Let

$$r := j(j - 1728) \frac{\omega}{dj}, \quad u := \frac{1}{j}.$$

Take $T \gg 0$ and let $P(x, y) = f_n(y)x^n + \cdots + f_1(y)x + f_0(y)$ be irreducible and such that $P(u, rj^T) = 0$.

Proposition (C.)

Then

$$F_{E,j}(x) = P\left(\frac{1}{x}, 0\right) \cdot x^A (x - 1728)^B$$

where A, B are explicitly computable.

So it suffices to compute the polynomial P . We can compute P using **linear algebra**, given the q -expansions of r and u .

The critical subgroup $E_{crit}(\mathbb{Q})$

Theorem (C.)

Suppose $r_{an}(E) \geq 2$, and assume at least one of the following holds:

(1) $F_{E,j} = \prod_{m=1}^k H_{D_m}^{s_i} \cdot F_0$, where $\mathbb{Q}(\sqrt{D_m}) \neq \mathbb{Q}(\sqrt{D_n})$ for all $m \neq n$, and F_0 is irreducible.

(2) $F_{E,h}$ is irreducible for some non-constant function $h \in \mathbb{Q}(X_0(N))$.

Then $\text{rank}(E_{crit}(\mathbb{Q})) = 0$.

Critical polynomials for elliptic curves of rank 2 and conductor < 1000 (I)

E	$g(X_0(N))$	h	Factorization of $F_{E,h}(x)$
389a	32	j	$H_{-19}(x)^2(x^{60} + \dots)$
433a	35	j	$x^{68} + \dots$
446d	55	j	$x^{108} + \dots$
563a	47	j	$H_{-43}(x)^2(x^{90} - \dots)$
571b	47	j	$H_{-67}(x)^2(x^{90} - \dots)$
643a	53	j	$H_{-19}(x)^2(x^{102} - \dots)$
664a	81	$\frac{\eta_4 \eta_8^2 \eta_{332}^5}{\eta_{166} \eta_{664}^6 \eta_2}$	$x^{160} - \dots$
655a	65	j	$x^{128} - \dots$
681c	75	j	$x^{148} - \dots$
707a	67	j	$x^{132} - \dots$

Critical polynomials for elliptic curves of rank 2 and conductor < 1000 (II)

E	$g(X_0(N))$	h	Factorization of $F_{E,h}(x)$
709a	58	j	$x^{114} - \dots$
718b	89	j	$H_{-52}(x)^2(x^{172} - \dots)$
794a	98	j	$H_{-4}(x)^2(x^{192} - \dots)$
817a	71	j	$x^{140} - \dots$
916c	113	j	$H_{-12}(x)^8(x^{216} + \dots)$
944e	115	$\frac{\eta_{16}^4 \eta_4^2}{\eta_8^6}$	$x^{224} - \dots$ ¹
997b	82	j	$H_{-27}(x)^2(x^{160} - \dots)$
997c	82	j	$x^{162} - \dots$

¹Here 4 of the critical points are cusps, so $\deg F = 2g - 6$.

Future work:

- Compute $E_{crit}(\mathbb{Q})$ for $E = \mathbf{5077a}$.
Current method will take roughly 5500/(number of cpus) hours.
- Prove or disprove that $\text{rank}(E_{crit}(\mathbb{Q})) = 0$ whenever $r_{an}(E)$ is even.
(For infinitely many?)

Motivation:

Modular forms

Let f be a function $f : \mathcal{H} \rightarrow \mathbb{C}$, $\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$, and let $k \in \mathbb{Z}$. The weight- k action of α on f is defined by

$$f|[\alpha]_k(z) := (cz + d)^{-k} f(\alpha z).$$

Definition

A **modular form** of weight k and level N is a holomorphic function $f : \mathcal{H} \rightarrow \mathbb{C}$ s.t.

- (1) $f(z) = f|[\alpha]_k(z)$, $\forall \alpha \in \Gamma_0(N)$ ($\Gamma_1(N)$).
- (2) f has holomorphic extension to all cusps of $X_0(N)$ ($X_1(N)$).

Cusp forms = modular forms that are zero at all cusps.

Modular forms have **q -expansions**: $f(q) = \sum_{n \geq 0} a_n q^n$, $q = \exp(2\pi iz)$.

The space of cusp forms = $S_k(N)$.

Operators on modular forms

- Hecke operators: a family $\{T_n, n \geq 1\} \cup \{\langle d \rangle : (d, N) = 1\}$ of commuting linear operators on $S_k(N)$.
- B_d and U_d operators: $B_d(\sum a_n q^n) = \sum a_n q^{nd}$,
 $U_d(\sum a_n q^n) = \sum a_{nd} q^n$.
- The Atkin-Lehner involution W_N . If f is a newform on $\Gamma_1(N)$, then

$$f|W_N = w(f)\bar{f}$$

$w(f) \in \mathbb{C}_1$ is called the **pseudo-eigenvalue** of f .

Newforms

- $S_k(N) = S_k(N)^{old} \oplus S_k(N)^{new}$.
- $S_k(N)^{new}$ has a basis of simultaneous eigenforms for **all** Hecke operators. These eigenforms are called **newforms**.

Fourier expansion

Let $f \in S_k(\Gamma_0(N))$ be a newform and let

$$\mathfrak{c} = \left[\frac{a}{c}\right] \in X_0(N)$$

be a cusp. Goal: compute the expansion of f at \mathfrak{c} . First, the expansion is only well-defined for $\text{denom}(\mathfrak{c})^2 \mid N$.

Let's denote the expansion by $f_{\mathfrak{c}}(q)$. Let $c' = \frac{N}{c}$.

Theorem (C.)

$$f_{\left[\frac{a}{c}\right]}(q) = \frac{w(f)}{\varphi(c')} \sum_{\chi: \text{cond}(\chi) | c'} \chi(-a) R(f, \chi)(q)$$

Here $R(f, \chi)(q)$ can be computed knowing $f \otimes \chi$ and $w(f \otimes \chi)$.

Algorithm for identifying $f \otimes \chi$

Lemma

Let ϵ be the character of f . Then the level of $f \otimes \chi$ divides $\text{lcm}(N, \text{cond}(\epsilon) \text{cond}(\chi), \text{cond}(\chi)^2)$.

Lemma

For every $N \geq 1$, there exists an integer $B = B(N)$ such that if g_1, g_2 be two normalised newforms of levels N_1, N_2 dividing N and

$$a_n(g_1) = a_n(g_2), \text{ for all } 1 \leq n \leq B \text{ such that } \gcd(n, N) = 1,$$

then $g_1 = g_2$.

Algorithm to compute $f \otimes \chi$

Algorithm 1 Identifying $f \otimes \chi$

Input: $f \in S_k(\Gamma_0(N))$ a normalized newform; χ – Dirichlet character of prime power conductor $Q = q^\beta$ ($Q^2 \mid N$).

Output: The newform $f \otimes \chi$.

for each $M \mid N$ **do**

 Compute a basis $\{g_1, \dots, g_s\}$ of $S_k(M, \chi^2)^{new}$.

$B :=$ the Sturm bound for $\Gamma_1(MQ^2)$.

for $1 \leq j \leq s$ **do**

if $a_n(g_j) = a_n(f)\chi(n)$ for all $1 \leq n \leq B, \gcd(n, q) = 1$ **then**

return g_j .

end if

end for

end for

return Error.

Computing the pseudo-eigenvalue

Recall that modular symbols are linear combinations of $\{\alpha, \beta\}$, $\alpha, \beta \in \mathbb{Q} \cup \infty$, and we put

$$\langle f, \{\alpha, \beta\} \rangle := \int_{\alpha}^{\beta} f dz.$$

Lemma (C.)

There exists a weight- k modular symbol M such that $W_N(M) = N^{k/2-1}M^$. Moreover, if $\langle f, M \rangle \neq 0$, then*

$$w(f) = \frac{\langle f, M \rangle}{\langle f, M \rangle}.$$

Examples (I)

Example

$E = 50a$. f_E is twist-minimal. Let $\mathfrak{c} = [\frac{1}{10}]$, write $\alpha_0 = 0$ and

$$x^4 + x^3 + x^2 - x + \frac{1}{5} = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)(x - \alpha_4).$$

Then

$$f_{\mathfrak{c}}(q) = \sum_{n \geq 1} \alpha_n \bmod 5 a_n(f) q^n.$$

Example

Let $E = 48a$ and let $\mathfrak{c} = [\frac{1}{12}]$. We computed that

$$f_{\mathfrak{c}}(q) = -2iq^2 + 2iq^6 + O(q^7).$$

Since the first coefficient vanishes, we conclude that the modular parametrization $\varphi : X_0(48) \rightarrow E$ is ramified at the cusp \mathfrak{c} .

Examples (II)

Definition

A newform f is *twist-minimal* if it is not a twist of a newform of lower level.

Example

Let $E = \mathbf{98a}$ and $\mathfrak{c} = [\frac{1}{14}]$. Then f_E is not twist-minimal. More precisely, if χ is the quadratic character modulo 7, then

$$f \otimes \chi(q) = q - q^2 - 2q^3 + q^4 + O(q^6)$$


is a newform of level 14. We computed numerically that

$$\begin{aligned} f_{\mathfrak{c}}(q) = & (-0.755 - 0.172i)q + (0.441 - 0.916i)q^2 + (1.392 + 1.110i)q^3 \\ & + (0.696 - 0.555i)q^4 + (1.510 - 0.344i)q^6 - 3.023iq^7 + O(q^8) \end{aligned}$$

Chow-Heegner points

- 1 Critical subgroups of elliptic curves
 - Elliptic curves and modular curves
 - The critical subgroup and critical polynomials
 - Computing the critical subgroup
- 2 q -expansion of newforms at non-unitary cusps
 - Problem description
- 3 Chow-Heegner points
 - Even index
 - Computing Chow-Heegner points

Definition: Chow-Heegner points



E, F : two non-isogeneous elliptic curves of same conductor N .

φ, ψ : modular parametrisations of E, F .

The **Chow-Heegner point** associated to the pair (E, F) is

$$P_{E,F} = \sum \varphi(\psi^*(Q)), \forall Q \in F(\mathbb{C})$$

- Facts: (1) $P_{E,F}$ is independent of the choice of Q ;
(2) $P_{E,F} \in E(\mathbb{Q})$.

Even index of Chow-Heegner points

Fact: $P_{E,F}$ is torsion when $r_{an}(E) \geq 2$.

When E has rank 1, numerical evidence suggests that the index $i_{E,F} := [E(\mathbb{Q})/\text{tors} : \mathbb{Z}P_{E,F}]$ is even, when it is finite.

Theorem (C.)

Let $\sigma_0(N)$ denote the number of distinct prime factors of N . If

$$\sigma_0(N) > \log_2(\#E[2](\mathbb{Q})) + \log_2(\#F[2](\mathbb{Q})) + 2,$$

then $P_{E,F} \in 2E(\mathbb{Q})$.

Computing Chow-Heegner points: previous work

There exist numerical algorithms to compute Chow-Heegner points.

- Darmon, Daub, Lichtenstein and Rotger – using (complex and p -adic) iterated integrals.
- Stein – using complex integration to lift points via modular parametrization.

We have developed an algebraic algorithm to compute the Chow-Heegner points, again using q -expansions.

Example

Example

$E = \mathbf{89a}$ and $F = \mathbf{89b}$. Let

$G_1(x) = x^4 + \frac{13}{4}x^3 + \frac{17}{4}x^2 + \frac{21}{4}x + \frac{9}{2} = \prod(x - a_i)$ and
 $b_i = -\frac{8}{9}a_i^3 - \frac{20}{9}a_i^2 - \frac{28}{9}a_i - \frac{10}{3}$. Then

$$P_{E,F} = \sum_{i=1}^4 P_i, \text{ where } P_i = (a_i, b_i).$$

We obtain $P_{E,F} = (\frac{3}{4}, -\frac{15}{8})$.

Future work

- Compute Chow-Heegner points for curves of different conductors.
- Prove $i_{E,F}$ is even without assumptions on N .
- Verify the numerical data of Chow-Heegner points in Stein's table and extend the table.

Thank you!