

PHY3110 SP23 Notes

Haoran Sun (haoransun@link.cuhk.edu.cn)

0 Introduction

Grading: 30% homework, 30% midterm, 40% final.

Textbooks:

- H. Goldstein, C. Poole, J. Safko, Classical Mechanics, 3rd Edition, Pearson.
- J.R. Taylor, Classical Mechanics, University Science Books.
- T.W.B. Kibble, F.H. Berkshire, Classical Mechanics, 5th Edition, Imperial College Press.
- 梁昆森, 力学 (下册) 理论力学, 4th Edition, 高等教育出版社.

Classical mechanics describe the motion of macroscopic objects, which are not extremely massive and not extremely fast.

1 Newtonian Mechanics

Vectorial quantities of motion: position \mathbf{r} , velocity \mathbf{v} , force \mathbf{F} , momentum $\mathbf{p} = m\mathbf{v}$, angular momentum $\mathbf{L} = \mathbf{r} \times \mathbf{p}$. Equations of motion are derived from those vector quantities.

Analytical mechanics uses scalar quantities of motion

- Kinetic energy $T = \frac{1}{2}m\mathbf{v}^2$
- Potential energy $V = V(\mathbf{r})$

Equations of motion are derived from those scalar quantities.

1.1 Newton's Laws

Newton's 2nd law

$$\mathbf{F} = \frac{d\mathbf{p}}{dt} = m\mathbf{a} \quad (1)$$

valid in an inertial frame. Angular momentum \mathbf{L} and torque \mathbf{N} are also related

$$\frac{d\mathbf{L}}{dt} = \frac{d}{dt}(\mathbf{r} \times \mathbf{p}) = \mathbf{r} \times \mathbf{F} = \mathbf{N} \quad (2)$$

Work done by external forces

$$W_{12} = \int_1^2 \mathbf{F} \cdot d\mathbf{s} = \int_1^2 m \frac{d\mathbf{v}}{dt} \cdot d\mathbf{s} = \int_1^2 m\mathbf{v} \cdot d\mathbf{v} = \left. \frac{1}{2}m\mathbf{v}^2 \right|_1^2 \quad (3)$$

Define a scalar function $V(\mathbf{r})$, then $\mathbf{F} = -\nabla V(\mathbf{r})$ is a conservative force.

$$\oint \mathbf{F} \cdot d\mathbf{s} = 0 \quad (4)$$

Center of mass of the system

$$\mathbf{R} = \frac{\sum_i m_i \mathbf{r}_i}{\sum_i m_i} = \frac{\sum_i m_i \mathbf{r}_i}{M} \quad (5)$$

Total momentum

$$\mathbf{P} = \sum_i m_i \mathbf{p}_i = M \dot{\mathbf{R}} \quad (6)$$

Hence \mathbf{P} is conserved if external force $\mathbf{F}^{(e)}$ is zero.

Total angular momentum

$$\frac{d\mathbf{L}}{dt} = \frac{d}{dt} \sum_i \mathbf{r}_i \times \mathbf{p}_i = \sum_i \mathbf{r}_i \times \left(\mathbf{F}_i^{(e)} + \sum_j \mathbf{F}_{ij} \right) = \sum_i \mathbf{r}_i \times \mathbf{F}_i^{(e)} + \sum_{ij} \mathbf{r}_i \times \mathbf{F}_{ij}$$

Since \mathbf{r}_{ij} parallel to \mathbf{F}_{ij} , then

$$\sum_{ij} \mathbf{r}_i \mathbf{F}_{ij} = \frac{1}{2} \sum_{ij} \mathbf{r}_{ij} \times \mathbf{F}_{ji} = 0 \quad (7)$$

Therefore

$$\frac{d\mathbf{L}}{dt} = \mathbf{N}^{(e)} \quad (8)$$

Decomposition of the angular momentum

$$\mathbf{L} = \sum_i \mathbf{r}_i \times \mathbf{p}_i = \sum_i (\mathbf{R} + \mathbf{r}_i) \times m_i (\mathbf{V} + \mathbf{v}_i') = \sum_i \mathbf{R} \times m_i \mathbf{V} + \sum_i \mathbf{r}_i' \times m_i \mathbf{v}_i' \quad (9)$$

1.2 Constraints

Holonomic constraint

$$f(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N, t) = 0 \quad (10)$$

Example: rigid body

$$(\mathbf{r}_i - \mathbf{r}_j)^2 - c_{ij}^2 = 0 \quad (11)$$

Example: non-sliding cylinder

$$\dot{x} - R\dot{\theta} = 0 \Rightarrow x - R\theta = \text{const}$$

A constraint of the form

$$\sum_i g_i(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) d\mathbf{x}_i = 0 \Rightarrow dG(\mathbf{x}_1, \dots) = 0 \Rightarrow G(\mathbf{x}_1, \dots) = \text{const} \quad (12)$$

Non-holonomic constraint: cannot be written in the form of holonomic constraint.

1.3 Generalized coordinates

Suppose we have a N -particle system, we will have $3N$ DOFs. With k constraints, we will have $3N - k$ DOFs. Define q_1, \dots, q_{3N-k} generalized coordinates, we have

$$\mathbf{r}_i = \mathbf{r}_i(q_1, \dots, q_{3N-k}, t) \quad (13)$$

2 Lagrange Formalism

2.1 D'Alembert's Principle

Hint from the rigid body: internal forces of constraints do not work.

Virtual displacement: $\delta \mathbf{r}_i$ is consistent with the constraints imposed on the system at a given time

$$\mathbf{r}_i \rightarrow \mathbf{r}_i + \delta \mathbf{r}_i \quad (14)$$

Consider a system in equilibrium

$$\mathbf{F}_i = 0 \Rightarrow \sum_i \mathbf{F}_i \cdot \delta \mathbf{r}_i = 0 \quad (15)$$

Separate $\mathbf{F}_i = \mathbf{F}_i^{(a)} + \mathbf{f}_i$ where \mathbf{f}_i is the constraint force. Hence

$$\sum_i (\mathbf{F}_i^{(a)} + \mathbf{f}_i) \cdot \delta \mathbf{r}_i = 0 \Rightarrow \sum_i \mathbf{F}_i^{(a)} \cdot \delta \mathbf{r}_i = 0 \quad (16)$$

For a system moving under external forces

$$\mathbf{F}_i - \dot{\mathbf{p}}_i = 0 \Rightarrow \sum_i (\mathbf{F}_i - \dot{\mathbf{p}}_i) \delta \mathbf{r}_i = 0 \Rightarrow \sum_i (\mathbf{F}_i^{(a)} - \dot{\mathbf{p}}_i) \delta \mathbf{r}_i = 0 \quad (17)$$

For holonomic constraints

$$\mathbf{r}_i = \mathbf{r}_i(q_1, \dots, q_n, t), \quad \mathbf{v}_i = \frac{d\mathbf{r}_i}{dt} = \frac{\partial \mathbf{r}_i}{\partial t} + \sum_j \frac{\partial \mathbf{r}_i}{\partial q_j} \dot{q}_j, \quad \delta \mathbf{r}_i = \sum_j \frac{\partial \mathbf{r}_i}{\partial q_j} \delta q_j \quad (18)$$

Generalized force Q_j

$$\sum_i \mathbf{F}_i \delta \mathbf{r}_i = \sum_{ij} \mathbf{F}_i \frac{\partial \mathbf{r}_i}{\partial q_j} \delta q_j = \sum_j Q_j \delta q_j \quad (19)$$

Then

$$\sum_i \dot{\mathbf{p}}_i \cdot \delta \mathbf{r}_i = \sum_{ij} m_i \ddot{\mathbf{r}}_i \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} \delta q_j = \sum_{ij} \left[\frac{d}{dt} \left(m_i \dot{\mathbf{r}}_i \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} \right) - m_i \dot{\mathbf{r}}_i \cdot \frac{d}{dt} \frac{\partial \mathbf{r}_i}{\partial q_j} \right] \delta q_j \quad (20)$$

$$= \sum_i \left[\frac{d}{dt} \left(\frac{\partial T}{\partial \dot{q}_j} \right) - \frac{\partial T}{\partial q_j} \right] \delta q_j = \sum_j Q_j \delta q_j \quad (21)$$

$$(22)$$

Hence $\forall j$ we have

$$\frac{d}{dt} \left(\frac{\partial T}{\partial \dot{q}_j} \right) - \frac{\partial T}{\partial q_j} - Q_j = 0 \quad (23)$$

Let the potential energy $V = V(\mathbf{r}_i, \dots) = V(q_j, \dots)$, then we have

$$Q_j = \sum_i \mathbf{F}_i \frac{\partial \mathbf{r}_i}{\partial q_j} = \sum_i -\nabla_i V \frac{\partial \mathbf{r}_i}{\partial q_j} = -\frac{\partial V}{\partial q_j} \quad (24)$$

Therefore

$$\frac{d}{dt} \left(\frac{\partial (T - V)}{\partial \dot{q}_j} \right) - \frac{\partial (T - V)}{\partial q_j} - Q_j = 0 \quad (25)$$