

UBC Virtual Physics Circle

A Hacker's Guide to Brownian motion

David Wakeham

June 25, 2020



Overview

Overview


- ▶ Our last topic in the hacker's guide is **Brownian motion**.

Overview

- ▶ Our last topic in the hacker's guide is **Brownian motion**. It's the other reason Einstein got a Nobel prize!


Overview

- ▶ Our last topic in the hacker's guide is **Brownian motion**. It's the other reason Einstein got a Nobel prize!



Overview


- ▶ Our last topic in the hacker's guide is **Brownian motion**. It's the other reason Einstein got a Nobel prize!



- ▶ We start with some **review**,

Overview


- ▶ Our last topic in the hacker's guide is **Brownian motion**. It's the other reason Einstein got a Nobel prize!



- ▶ We start with some **review**, then we do some **thermodynamics**.

Overview

- ▶ Our last topic in the hacker's guide is **Brownian motion**. It's the other reason Einstein got a Nobel prize!



- ▶ We start with some **review**, then we do some **thermodynamics**. Add it all up to get Brownian motion!

Review

Review: Stokes' law

Review: Stokes' law

- ▶ First, we need to remind ourselves of a result from the lecture on **dimensional analysis**.

Review: Stokes' law

- ▶ First, we need to remind ourselves of a result from the lecture on **dimensional analysis**.
- ▶ A sphere of **radius R**

Review: Stokes' law

- ▶ First, we need to remind ourselves of a result from the lecture on **dimensional analysis**.
- ▶ A sphere of **radius R** moves at (slow) **speed v**

Review: Stokes' law


- ▶ First, we need to remind ourselves of a result from the lecture on **dimensional analysis**.
- ▶ A sphere of **radius R** moves at (slow) **speed v** through a fluid of **viscosity η** .

Review: Stokes' law

- ▶ First, we need to remind ourselves of a result from the lecture on **dimensional analysis**.
- ▶ A sphere of **radius** R moves at (slow) **speed** v through a fluid of **viscosity** η . (Units: $[\eta] = M/LT$.)


Review: Stokes' law

- ▶ First, we need to remind ourselves of a result from the lecture on **dimensional analysis**.
- ▶ A sphere of **radius R** moves at (slow) **speed v** through a fluid of **viscosity η** . (Units: $[\eta] = M/LT$.)



Review: Stokes' law


- ▶ First, we need to remind ourselves of a result from the lecture on **dimensional analysis**.
- ▶ A sphere of **radius R** moves at (slow) **speed v** through a fluid of **viscosity η** . (Units: $[\eta] = M/LT$.)



- ▶ **Stokes' law** states that the drag force is

Review: Stokes' law

- ▶ First, we need to remind ourselves of a result from the lecture on **dimensional analysis**.
- ▶ A sphere of **radius R** moves at (slow) **speed v** through a fluid of **viscosity η** . (Units: $[\eta] = M/LT.$)



- ▶ **Stokes' law** states that the drag force is

$$F_{\text{drag}} = 6\pi\eta vR.$$

Review: random walks and collisions

Review: random walks and collisions

- ▶ Last lecture, we introduced random walks and collisions.

Review: random walks and collisions

- ▶ Last lecture, we introduced random walks and collisions.
- ▶ Walks:

Review: random walks and collisions

- ▶ Last lecture, we introduced random walks and collisions.
- ▶ Walks: A random walk of N steps with length ℓ wanders

Review: random walks and collisions

- ▶ Last lecture, we introduced **random walks** and **collisions**.
- ▶ **Walks:** A random walk of N steps with length ℓ wanders

$$d \sim \sqrt{N}\ell.$$

Review: random walks and collisions

- ▶ Last lecture, we introduced **random walks** and **collisions**.
- ▶ **Walks:** A random walk of N steps with length ℓ wanders

$$d \sim \sqrt{N}\ell.$$

- ▶ **Collisions:** If you have cross-section σ ,

Review: random walks and collisions

- ▶ Last lecture, we introduced **random walks** and **collisions**.
- ▶ **Walks:** A random walk of N steps with length ℓ wanders

$$d \sim \sqrt{N}\ell.$$

- ▶ **Collisions:** If you have cross-section σ , and collide with stuff of density n (number per unit volume),

Review: random walks and collisions

- ▶ Last lecture, we introduced **random walks** and **collisions**.
- ▶ **Walks:** A random walk of N steps with length ℓ wanders

$$d \sim \sqrt{N}\ell.$$



- ▶ **Collisions:** If you have cross-section σ , and collide with stuff of density n (number per unit volume), your **mean free path** λ between collisions is

Review: random walks and collisions

- ▶ Last lecture, we introduced **random walks** and **collisions**.
- ▶ **Walks**: A random walk of N steps with length ℓ wanders

$$d \sim \sqrt{N}\ell.$$

- ▶ **Collisions**: If you have cross-section σ , and collide with stuff of density n (number per unit volume), your **mean free path** λ between collisions is



Speed and diffusion

Speed and diffusion

- ▶ There is another useful piece of terminology.

Speed and diffusion

- ▶ There is another useful piece of terminology.
- ▶ In some time t , suppose a walk spreads a distance d .

Speed and diffusion

- ▶ There is another useful piece of terminology.
- ▶ In some time t , suppose a walk spreads a distance d .
The **diffusion constant** D is defined by

Speed and diffusion

- ▶ There is another useful piece of terminology.
- ▶ In some time t , suppose a walk spreads a distance d . The **diffusion constant** D is defined by

$$d = \sqrt{Dt}.$$

Speed and diffusion

- ▶ There is another useful piece of terminology.
- ▶ In some time t , suppose a walk spreads a distance d .
The **diffusion constant** D is defined by

$$d = \sqrt{Dt}.$$

- ▶ Assume the walker moves at **speed** v .

Speed and diffusion

- ▶ There is another useful piece of terminology.
- ▶ In some time t , suppose a walk spreads a distance d . The **diffusion constant** D is defined by

$$d = \sqrt{Dt}.$$

- ▶ Assume the walker moves at **speed** v . Each step takes time $\tau = \ell/v$, and N steps take time $t = N\tau$.

Speed and diffusion

- ▶ There is another useful piece of terminology.
- ▶ In some time t , suppose a walk spreads a distance d . The **diffusion constant** D is defined by

$$d = \sqrt{Dt}.$$

- ▶ Assume the walker moves at **speed** v . Each step takes time $\tau = \ell/v$, and N steps take time $t = N\tau$. Then

$$D = \frac{d^2}{t} = \frac{(\sqrt{N}\ell)^2}{t} = \frac{N\ell^2}{t} = \frac{\ell^2}{\tau} = v\ell.$$

Speed and diffusion

- ▶ There is another useful piece of terminology.
- ▶ In some time t , suppose a walk spreads a distance d . The **diffusion constant** D is defined by

$$d = \sqrt{Dt}.$$

- ▶ Assume the walker moves at **speed** v . Each step takes time $\tau = \ell/v$, and N steps take time $t = N\tau$. Then

$$D = \frac{d^2}{t} = \frac{(\sqrt{N}\ell)^2}{t} = \frac{N\ell^2}{t} = \frac{\ell^2}{\tau} = v\ell.$$

- ▶ So the **diffusion constant** $D = v\ell$.

Exercise 1: dodgem cars

Exercise 1: dodgem cars


- ▶ Dodgem cars travel on average 1 m/s ,

Exercise 1: dodgem cars

- ▶ Dodgem cars travel on average 1 m/s , with $\sigma \sim 2 \text{ m}$.


Exercise 1: dodgem cars

- ▶ Dodgem cars travel on average 1 m/s , with $\sigma \sim 2 \text{ m}$.



Exercise 1: dodgem cars


- ▶ Dodgem cars travel on average 1 m/s , with $\sigma \sim 2 \text{ m}$.



- ▶ You and 4 of friends are colliding randomly on a square arena 5 m in width.

Exercise 1: dodgem cars


- ▶ Dodgem cars travel on average 1 m/s , with $\sigma \sim 2 \text{ m}$.



- ▶ You and 4 of friends are colliding randomly on a square arena 5 m in width.
- ▶ 1. What is the mean free path λ ?

Exercise 1: dodgem cars


- ▶ Dodgem cars travel on average 1 m/s , with $\sigma \sim 2 \text{ m}$.



- ▶ You and 4 of friends are colliding randomly on a square arena 5 m in width.
- ▶
 1. What is the **mean free path** λ ?
 2. Roughly **how long** does it take to bounce from the center to the edge of the arena?

Exercise 1: dodgem cars


- ▶ Dodgem cars travel on average 1 m/s , with $\sigma \sim 2 \text{ m}$.



- ▶ You and 4 of friends are colliding randomly on a square arena 5 m in width.
- ▶
 1. What is the **mean free path** λ ?
 2. Roughly **how long** does it take to bounce from the center to the edge of the arena? Use $t = d^2/D = d^2/\ell v$.

Exercise 1: dodgem cars

- ▶ Dodgem cars travel on average 1 m/s , with $\sigma \sim 2 \text{ m}$.



- ▶ You and 4 of friends are colliding randomly on a square arena 5 m in width.
- ▶
 1. What is the **mean free path** λ ?
 2. Roughly **how long** does it take to bounce from the center to the edge of the arena? Use $t = d^2/D = d^2/\ell v$.

Exercise 1: dodgem cars (solution)

Exercise 1: dodgem cars (solution)

- ▶ There are 5 cars in a 25 m^2 area, so $n = 0.2 \text{ m}^{-2}$.

Exercise 1: dodgem cars (solution)

- ▶ There are 5 cars in a 25 m^2 area, so $n = 0.2 \text{ m}^{-2}$.
The cross-section is $\sigma = 2 \text{ m}$, and hence the mfp is

Exercise 1: dodgem cars (solution)

- ▶ There are 5 cars in a 25 m^2 area, so $n = 0.2 \text{ m}^{-2}$.
The cross-section is $\sigma = 2 \text{ m}$, and hence the mfp is

$$\lambda = \frac{1}{\sigma n} = \frac{1}{2 \times 0.2} \text{ m} = 2.5 \text{ m}.$$

Exercise 1: dodgem cars (solution)

- ▶ There are 5 cars in a 25 m^2 area, so $n = 0.2 \text{ m}^{-2}$.
The cross-section is $\sigma = 2 \text{ m}$, and hence the mfp is

$$\lambda = \frac{1}{\sigma n} = \frac{1}{2 \times 0.2} \text{ m} = 2.5 \text{ m}.$$

- ▶ Using the hint,

Exercise 1: dodgem cars (solution)

- ▶ There are 5 cars in a 25 m^2 area, so $n = 0.2 \text{ m}^{-2}$.
The cross-section is $\sigma = 2 \text{ m}$, and hence the mfp is

$$\lambda = \frac{1}{\sigma n} = \frac{1}{2 \times 0.2} \text{ m} = 2.5 \text{ m}.$$

- ▶ Using the hint,

$$t = \frac{d^2}{\ell v} = \frac{25}{2.5 \times 1} \text{ s} = 10 \text{ s}.$$

Exercise 1: dodgem cars (solution)

- ▶ There are 5 cars in a 25 m^2 area, so $n = 0.2 \text{ m}^{-2}$.
The cross-section is $\sigma = 2 \text{ m}$, and hence the mfp is

$$\lambda = \frac{1}{\sigma n} = \frac{1}{2 \times 0.2} \text{ m} = 2.5 \text{ m}.$$

- ▶ Using the hint,

$$t = \frac{d^2}{\ell v} = \frac{25}{2.5 \times 1} \text{ s} = 10 \text{ s}.$$

- ▶ In reality, you're trying to hit each other.

Exercise 1: dodgem cars (solution)

- ▶ There are 5 cars in a 25 m^2 area, so $n = 0.2 \text{ m}^{-2}$.
The cross-section is $\sigma = 2 \text{ m}$, and hence the mfp is

$$\lambda = \frac{1}{\sigma n} = \frac{1}{2 \times 0.2} \text{ m} = 2.5 \text{ m}.$$

- ▶ Using the hint,

$$t = \frac{d^2}{\ell v} = \frac{25}{2.5 \times 1} \text{ s} = 10 \text{ s}.$$

- ▶ In reality, you're trying to hit each other. So this is a good model only if, e.g.

Exercise 1: dodgem cars (solution)

- ▶ There are 5 cars in a 25 m^2 area, so $n = 0.2 \text{ m}^{-2}$.
The cross-section is $\sigma = 2 \text{ m}$, and hence the mfp is

$$\lambda = \frac{1}{\sigma n} = \frac{1}{2 \times 0.2} \text{ m} = 2.5 \text{ m}.$$

- ▶ Using the hint,

$$t = \frac{d^2}{\ell v} = \frac{25}{2.5 \times 1} \text{ s} = 10 \text{ s}.$$

- ▶ In reality, you're trying to hit each other. So this is a good model only if, e.g. **you fall asleep!**

Review: ideal gas law

Review: ideal gas law

- ▶ To connect to atomic motion, we need to learn about the ideal gas law.

Review: ideal gas law


- ▶ To connect to atomic motion, we need to learn about the ideal gas law. (This may be review for many of you.)

Review: ideal gas law

- ▶ To connect to atomic motion, we need to learn about the ideal gas law. (This may be review for many of you.)
- ▶ Imagine a balloon \mathcal{N} gas particles.


Review: ideal gas law

- ▶ To connect to atomic motion, we need to learn about the ideal gas law. (This may be review for many of you.)
- ▶ Imagine a balloon \mathcal{N} gas particles.



Review: ideal gas law


- ▶ To connect to atomic motion, we need to learn about the **ideal gas law**. (This may be review for many of you.)
- ▶ Imagine a balloon \mathcal{N} **gas particles**.



- ▶ The gas is hot (**temperature \mathcal{T}**),

Review: ideal gas law


- ▶ To connect to atomic motion, we need to learn about the ideal gas law. (This may be review for many of you.)
- ▶ Imagine a balloon \mathcal{N} gas particles.



- ▶ The gas is hot (temperature \mathcal{T}), takes up space (volume V),

Review: ideal gas law

- ▶ To connect to atomic motion, we need to learn about the **ideal gas law**. (This may be review for many of you.)
- ▶ Imagine a balloon \mathcal{N} **gas particles**.



- ▶ The gas is hot (**temperature \mathcal{T}**), takes up space (**volume V**), and presses on the balloon (**pressure P**).

Derivation of ideal gas law

Derivation of ideal gas law

- ▶ The ideal gas law states that these properties are related:

Derivation of ideal gas law

- ▶ The ideal gas law states that these properties are related:

$$PV = k_B \mathcal{N} \mathcal{T}.$$

Derivation of ideal gas law

- ▶ The ideal gas law states that these properties are related:

$$PV = k_B \mathcal{N} \mathcal{T}.$$

- ▶ Here, $k_B = 1.38 \times 10^{-23}$ J/K is Boltzmann's constant.

Derivation of ideal gas law

- ▶ The **ideal gas law** states that these properties are related:

$$PV = k_B \mathcal{N} \mathcal{T}.$$

- ▶ Here, $k_B = 1.38 \times 10^{-23}$ J/K is **Boltzmann's constant**.
- ▶ We can **“derive”** this from dimensional analysis!

Derivation of ideal gas law

- ▶ The **ideal gas law** states that these properties are related:

$$PV = k_B \mathcal{N} \mathcal{T}.$$

- ▶ Here, $k_B = 1.38 \times 10^{-23}$ J/K is **Boltzmann's constant**.
- ▶ We can **“derive”** this from dimensional analysis! But we **need more** than *MLT* (mass, length, time).

Derivation of ideal gas law

- ▶ The **ideal gas law** states that these properties are related:

$$PV = k_B \mathcal{N} T.$$

- ▶ Here, $k_B = 1.38 \times 10^{-23}$ J/K is **Boltzmann's constant**.
- ▶ We can **“derive”** this from dimensional analysis! But we **need more** than *MLT* (mass, length, time).
- ▶ In addition to length L and time T , use the following:

Derivation of ideal gas law

- ▶ The **ideal gas law** states that these properties are related:

$$PV = k_B \mathcal{N} \mathcal{T}.$$

- ▶ Here, $k_B = 1.38 \times 10^{-23}$ J/K is **Boltzmann's constant**.
- ▶ We can “**derive**” this from dimensional analysis! But we **need more** than MLT (mass, length, time).
- ▶ In addition to length L and time T , use the following:
 1. **energy** \mathcal{E} (instead of M);

Derivation of ideal gas law

- ▶ The **ideal gas law** states that these properties are related:

$$PV = k_B \mathcal{N} \mathcal{T}.$$

- ▶ Here, $k_B = 1.38 \times 10^{-23}$ J/K is **Boltzmann's constant**.
- ▶ We can “**derive**” this from dimensional analysis! But we **need more** than MLT (mass, length, time).
- ▶ In addition to length L and time T , use the following:
 1. **energy** \mathcal{E} (instead of M);
 2. **temperature** Θ ;

Derivation of ideal gas law

- ▶ The **ideal gas law** states that these properties are related:

$$PV = k_B \mathcal{N} T.$$

- ▶ Here, $k_B = 1.38 \times 10^{-23}$ J/K is **Boltzmann's constant**.
- ▶ We can “**derive**” this from dimensional analysis! But we **need more** than MLT (mass, length, time).
- ▶ In addition to length L and time T , use the following:
 1. **energy** \mathcal{E} (instead of M);
 2. **temperature** Θ ;
 3. and **particle number** Ξ .

Derivation of ideal gas law

- ▶ The **ideal gas law** states that these properties are related:

$$PV = k_B \mathcal{N} T.$$

- ▶ Here, $k_B = 1.38 \times 10^{-23}$ J/K is **Boltzmann's constant**.
- ▶ We can **“derive”** this from dimensional analysis! But we **need more** than MLT (mass, length, time).
- ▶ In addition to length L and time T , use the following:
 1. **energy** \mathcal{E} (instead of M);
 2. **temperature** Θ ;
 3. and **particle number** Ξ .
- ▶ Ξ is for stuff growing with particle number,

Derivation of ideal gas law

- ▶ The **ideal gas law** states that these properties are related:

$$PV = k_B \mathcal{N} T.$$

- ▶ Here, $k_B = 1.38 \times 10^{-23}$ J/K is **Boltzmann's constant**.
- ▶ We can “**derive**” this from dimensional analysis! But we **need more** than *MLT* (mass, length, time).
- ▶ In addition to length L and time T , use the following:
 1. **energy** \mathcal{E} (instead of M);
 2. **temperature** Θ ;
 3. and **particle number** Ξ .
- ▶ Ξ is for stuff growing with particle number, e.g. $[\mathcal{N}] = \Xi$.

Exercise 2: ideal gas

Exercise 2: ideal gas

(a) Show that P and V have dimension

Exercise 2: ideal gas

(a) Show that P and V have dimension

$$[V] = \Xi L^3, \quad [P] = \frac{\mathcal{E}}{L^3}.$$

Exercise 2: ideal gas

(a) Show that P and V have dimension

$$[V] = \Xi L^3, \quad [P] = \frac{\mathcal{E}}{L^3}.$$

(b) From $k_B = 1.38 \times 10^{-23} \text{ J/K}$, deduce $[k_B] = \mathcal{E}/\Theta$.

Exercise 2: ideal gas

- (a) Show that P and V have dimension

$$[V] = \Xi L^3, \quad [P] = \frac{\mathcal{E}}{L^3}.$$

- (b) From $k_B = 1.38 \times 10^{-23} \text{ J/K}$, deduce $[k_B] = \mathcal{E}/\Theta$.

- (c) Conclude that

Exercise 2: ideal gas

- (a) Show that P and V have dimension

$$[V] = \Xi L^3, \quad [P] = \frac{\mathcal{E}}{L^3}.$$

- (b) From $k_B = 1.38 \times 10^{-23} \text{ J/K}$, deduce $[k_B] = \mathcal{E}/\Theta$.

- (c) Conclude that

$$[PV] = [\mathcal{N}k_B\mathcal{T}] = \Xi\mathcal{E}.$$

Exercise 2: ideal gas

- (a) Show that P and V have dimension

$$[V] = \Xi L^3, \quad [P] = \frac{\mathcal{E}}{L^3}.$$

- (b) From $k_B = 1.38 \times 10^{-23} \text{ J/K}$, deduce $[k_B] = \mathcal{E}/\Theta$.

- (c) Conclude that

$$[PV] = [\mathcal{N}k_B\mathcal{T}] = \Xi\mathcal{E}.$$

- With more care, you can show $PV = \mathcal{N}k_B\mathcal{T}$

Exercise 2: ideal gas

- (a) Show that P and V have dimension

$$[V] = \Xi L^3, \quad [P] = \frac{\mathcal{E}}{L^3}.$$

- (b) From $k_B = 1.38 \times 10^{-23} \text{ J/K}$, deduce $[k_B] = \mathcal{E}/\Theta$.

- (c) Conclude that

$$[PV] = [\mathcal{N}k_B\mathcal{T}] = \Xi\mathcal{E}.$$

- With more care, you can show $PV = \mathcal{N}k_B\mathcal{T}$ is the **only dimensionally consistent relation** between all these things.

Exercise 2: ideal gas (solution)

Exercise 2: ideal gas (solution)

(a) Usually, V has units $[V] = L^3$.

Exercise 2: ideal gas (solution)

- (a) Usually, V has units $[V] = L^3$. But the volume of a gas grows with particle number, so $[V] = \Xi L^3$.

Exercise 2: ideal gas (solution)

- (a) Usually, V has units $[V] = L^3$. But the volume of a gas grows with particle number, so $[V] = \Xi L^3$.
- ▶ As for pressure, using work ($Fd = W$)

Exercise 2: ideal gas (solution)

(a) Usually, V has units $[V] = L^3$. But the volume of a gas grows with particle number, so $[V] = \Xi L^3$.

- ▶ As for pressure, using work ($Fd = W$)

$$P = \frac{F}{A} = \frac{W}{Ad} \implies [P] = \frac{[W]}{[Ad]} = \frac{\mathcal{E}}{L^3}.$$

Exercise 2: ideal gas (solution)

(a) Usually, V has units $[V] = L^3$. But the volume of a gas grows with particle number, so $[V] = \Xi L^3$.

► As for pressure, using work ($Fd = W$)

$$P = \frac{F}{A} = \frac{W}{Ad} \implies [P] = \frac{[W]}{[Ad]} = \frac{\mathcal{E}}{L^3}.$$

(b) We have

Exercise 2: ideal gas (solution)

(a) Usually, V has units $[V] = L^3$. But the volume of a gas grows with particle number, so $[V] = \Xi L^3$.

► As for pressure, using work ($Fd = W$)

$$P = \frac{F}{A} = \frac{W}{Ad} \implies [P] = \frac{[W]}{[Ad]} = \frac{\mathcal{E}}{L^3}.$$

(b) We have

$$[k_B = 1.38 \times 10^{-23} \text{ J/K}] = [\text{J}]/[\text{K}] = \mathcal{E}/\Theta.$$

Exercise 2: ideal gas (solution)

(a) Usually, V has units $[V] = L^3$. But the volume of a gas grows with particle number, so $[V] = \Xi L^3$.

► As for pressure, using work ($Fd = W$)

$$P = \frac{F}{A} = \frac{W}{Ad} \implies [P] = \frac{[W]}{[Ad]} = \frac{\mathcal{E}}{L^3}.$$

(b) We have

$$[k_B = 1.38 \times 10^{-23} \text{ J/K}] = [\text{J}]/[\text{K}] = \mathcal{E}/\Theta.$$

(c) From part (a), $[PV] = (\Xi L^3)(\mathcal{E}/L^3) = \Xi \mathcal{E}$.

Exercise 2: ideal gas (solution)

- (a) Usually, V has units $[V] = L^3$. But the volume of a gas grows with particle number, so $[V] = \Xi L^3$.

- As for pressure, using work ($Fd = W$)

$$P = \frac{F}{A} = \frac{W}{Ad} \implies [P] = \frac{[W]}{[Ad]} = \frac{\mathcal{E}}{L^3}.$$

- (b) We have

$$[k_B = 1.38 \times 10^{-23} \text{ J/K}] = [\text{J}]/[\text{K}] = \mathcal{E}/\Theta.$$

- (c) From part (a), $[PV] = (\Xi L^3)(\mathcal{E}/L^3) = \Xi \mathcal{E}$.
From part (b), $[k_B \mathcal{N} \mathcal{T}] = (\mathcal{E}/\Theta)[\mathcal{N}][\mathcal{T}] = \Xi \mathcal{E}$.

Brownian motion

Lucretius, Brown, Einstein

Lucretius, Brown, Einstein

- ▶ A little history!

Lucretius, Brown, Einstein

- ▶ A little history! In 60 BC, Roman philosopher Lucretius observed the zigzag motion of dust motes.

Lucretius, Brown, Einstein

- ▶ A little history! In 60 BC, Roman philosopher Lucretius observed the zigzag motion of dust motes. He correctly attributed it to collisions with tiny invisible particles.

Lucretius, Brown, Einstein

- ▶ A little history! In 60 BC, Roman philosopher Lucretius observed the zigzag motion of dust motes. He correctly attributed it to collisions with tiny invisible particles.
- ▶ In 1827, botanist Robert Brown saw pollen jiggle under a microscope.

Lucretius, Brown, Einstein

- ▶ A little history! In 60 BC, Roman philosopher Lucretius observed the **zigzag motion of dust motes**. He correctly attributed it to **collisions with tiny invisible particles**.
- ▶ In 1827, botanist Robert Brown saw **pollen jiggle** under a microscope. Unlike Lucretius, he couldn't explain it!

Lucretius, Brown, Einstein

- ▶ A little history! In 60 BC, Roman philosopher Lucretius observed the **zigzag motion of dust motes**. He correctly attributed it to **collisions with tiny invisible particles**.
- ▶ In 1827, botanist Robert Brown saw **pollen jiggle** under a microscope. Unlike Lucretius, he couldn't explain it!
- ▶ Most 19th century scientists were **skeptical of atoms**.

Lucretius, Brown, Einstein

- ▶ A little history! In 60 BC, Roman philosopher Lucretius observed the **zigzag motion of dust motes**. He correctly attributed it to **collisions with tiny invisible particles**.
- ▶ In 1827, botanist Robert Brown saw **pollen jiggle** under a microscope. Unlike Lucretius, he couldn't explain it!
- ▶ Most 19th century scientists were **skeptical of atoms**.
- ▶ In 1905, a 26-year old Swiss patent clerk finished a PhD on Brownian motion,

Lucretius, Brown, Einstein

- ▶ A little history! In 60 BC, Roman philosopher Lucretius observed the **zigzag motion of dust motes**. He correctly attributed it to **collisions with tiny invisible particles**.
- ▶ In 1827, botanist Robert Brown saw **pollen jiggle** under a microscope. Unlike Lucretius, he couldn't explain it!
- ▶ Most 19th century scientists were **skeptical of atoms**.
- ▶ In 1905, a 26-year old Swiss patent clerk finished a PhD on Brownian motion, expanding on Lucretius' idea to account for the jiggling grains.

Lucretius, Brown, Einstein

- ▶ A little history! In 60 BC, Roman philosopher Lucretius observed the **zigzag motion of dust motes**. He correctly attributed it to **collisions with tiny invisible particles**.
- ▶ In 1827, botanist Robert Brown saw **pollen jiggle** under a microscope. Unlike Lucretius, he couldn't explain it!
- ▶ Most 19th century scientists were **skeptical of atoms**.
- ▶ In 1905, a 26-year old Swiss patent clerk finished a PhD on Brownian motion, expanding on Lucretius' idea to account for the jiggling grains. That clerk: **Einstein!**

The pollen polka

The pollen polka

- ▶ We will reproduce one of the main results of Einstein's PhD thesis using cheap guesswork, i.e. **hacking**.

The pollen polka


- ▶ We will reproduce one of the main results of Einstein's PhD thesis using cheap guesswork, i.e. **hacking**.
- ▶ Pour a viscous fluid into a container,

The pollen polka

- ▶ We will reproduce one of the main results of Einstein's PhD thesis using cheap guesswork, i.e. **hacking**.
- ▶ Pour a viscous fluid into a container, then plonk a few spherical pollen grains into it, as below:


The pollen polka

- ▶ We will reproduce one of the main results of Einstein's PhD thesis using cheap guesswork, i.e. **hacking**.
- ▶ Pour a viscous fluid into a container, then plonk a few spherical pollen grains into it, as below:



The pollen polka


- ▶ We will reproduce one of the main results of Einstein's PhD thesis using cheap guesswork, i.e. **hacking**.
- ▶ Pour a viscous fluid into a container, then plonk a few spherical pollen grains into it, as below:



- ▶ The pollen (pink) will start **jiggling around** as it collides with fluid molecules (green),

The pollen polka

- ▶ We will reproduce one of the main results of Einstein's PhD thesis using cheap guesswork, i.e. **hacking**.
- ▶ Pour a viscous fluid into a container, then plonk a few spherical pollen grains into it, as below:



- ▶ The pollen (pink) will start **jiggling around** as it collides with fluid molecules (green), executing a **random walk**.

Brownian motion: mean free path

Brownian motion: mean free path

- ▶ Let's put it all together to see how far the pollen jitters.

Brownian motion: mean free path

- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ !

Brownian motion: mean free path

- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ ! The pollen is much larger than the molecules.

Brownian motion: mean free path

- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ ! The pollen is much larger than the molecules. If it has radius R ,

Brownian motion: mean free path


- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ ! The pollen is much larger than the molecules. If it has radius R , it has cross-section $\sigma =$

Brownian motion: mean free path

- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ ! The pollen is much larger than the molecules. If it has radius R , it has cross-section $\sigma = \pi R^2$.


Brownian motion: mean free path

- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ ! The pollen is much larger than the molecules. If it has radius R , it has cross-section $\sigma = \pi R^2$.



Brownian motion: mean free path


- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ ! The pollen is much larger than the molecules. If it has radius R , it has cross-section $\sigma = \pi R^2$.



- ▶ Assume the fluid obeys the ideal gas law, $PV = k_B \mathcal{N} \mathcal{T}$.

Brownian motion: mean free path


- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ ! The pollen is much larger than the molecules. If it has radius R , it has cross-section $\sigma = \pi R^2$.



- ▶ Assume the fluid obeys the ideal gas law, $PV = k_B \mathcal{N} \mathcal{T}$. Since density $n = \mathcal{N}/V$,

Brownian motion: mean free path


- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ ! The pollen is much larger than the molecules. If it has radius R , it has cross-section $\sigma = \pi R^2$.



- ▶ Assume the fluid obeys the ideal gas law, $PV = k_B \mathcal{N} \mathcal{T}$. Since density $n = \mathcal{N}/V$, the mfp λ is

Brownian motion: mean free path

- ▶ Let's put it all together to see how far the pollen jitters. This is measured by the diffusion coefficient $D = \lambda v$.
- ▶ First, λ ! The pollen is much larger than the molecules. If it has radius R , it has cross-section $\sigma = \pi R^2$.



- ▶ Assume the fluid obeys the ideal gas law, $PV = k_B \mathcal{N} \mathcal{T}$. Since density $n = \mathcal{N}/V$, the mfp λ is

$$\lambda = \frac{1}{n\sigma} = \frac{V}{\mathcal{N}\pi R^2} = \frac{V}{\mathcal{N}\pi R^2} \times \frac{k_B \mathcal{N} \mathcal{T}}{PV} = \frac{k_B \mathcal{T}}{\pi P R^2}.$$

Brownian motion: terminal velocity

Brownian motion: terminal velocity

- ▶ What about the speed v ?

Brownian motion: terminal velocity

- ▶ What about the speed v ? A reasonable guess is **terminal velocity**, achieved when **weight balances drag force**.

Brownian motion: terminal velocity

- ▶ What about the speed v ? A reasonable guess is **terminal velocity**, achieved when **weight balances drag force**.
- ▶ From **Stokes' law**, if the fluid has **viscosity η** ,

Brownian motion: terminal velocity


- ▶ What about the speed v ? A reasonable guess is **terminal velocity**, achieved when **weight balances drag force**.
- ▶ From **Stokes' law**, if the fluid has **viscosity η** ,

$$mg = 6\pi\eta v_{\text{term}} R \quad \implies \quad v_{\text{term}} = \frac{mg}{6\pi\eta R}.$$

Brownian motion: terminal velocity

- ▶ What about the speed v ? A reasonable guess is **terminal velocity**, achieved when **weight balances drag force**.
- ▶ From **Stokes' law**, if the fluid has **viscosity η** ,

$$mg = 6\pi\eta v_{\text{term}} R \quad \implies \quad v_{\text{term}} = \frac{mg}{6\pi\eta R}.$$




$$F_{\text{drag}} = 6\pi\eta v R$$
$$W = mg$$

Brownian motion: terminal velocity

- ▶ What about the speed v ? A reasonable guess is **terminal velocity**, achieved when **weight balances drag force**.
- ▶ From **Stokes' law**, if the fluid has **viscosity η** ,

$$mg = 6\pi\eta v_{\text{term}} R \quad \implies \quad v_{\text{term}} = \frac{mg}{6\pi\eta R}.$$




- ▶ Combining with our expression for λ ,

Brownian motion: terminal velocity

- ▶ What about the speed v ? A reasonable guess is **terminal velocity**, achieved when **weight balances drag force**.
- ▶ From **Stokes' law**, if the fluid has **viscosity η** ,

$$mg = 6\pi\eta v_{\text{term}} R \quad \implies \quad v_{\text{term}} = \frac{mg}{6\pi\eta R}.$$




- ▶ Combining with our expression for λ , we get

Brownian motion: terminal velocity

- ▶ What about the speed v ? A reasonable guess is **terminal velocity**, achieved when **weight balances drag force**.
- ▶ From **Stokes' law**, if the fluid has **viscosity η** ,

$$mg = 6\pi\eta v_{\text{term}} R \quad \Longrightarrow \quad v_{\text{term}} = \frac{mg}{6\pi\eta R}.$$



- ▶ Combining with our expression for λ , we get

$$D = \lambda v_{\text{term}} = \frac{k_B \mathcal{T}}{\pi P R^2} \cdot \frac{mg}{6\pi\eta R}.$$

Brownian motion: magic trick!

Brownian motion: magic trick!

- ▶ A magic trick:

Brownian motion: magic trick!

- ▶ A magic trick: suppose the pollen settles at a height

Brownian motion: magic trick!

- ▶ A magic trick: suppose the pollen settles at a height where pressure balances weight,

Brownian motion: magic trick!


- ▶ A magic trick: suppose the pollen settles at a height where pressure balances weight, or $mg = PA$

Brownian motion: magic trick!

- ▶ A magic trick: suppose the pollen settles at a height where pressure balances weight, or $mg = PA = \pi PR^2$.


Brownian motion: magic trick!

- ▶ A magic trick: suppose the pollen settles at a height where pressure balances weight, or $mg = PA = \pi PR^2$.



Brownian motion: magic trick!


- ▶ A magic trick: suppose the pollen settles at a height where pressure balances weight, or $mg = PA = \pi PR^2$.



- ▶ Plugging this into D gives the Stokes-Einstein relation:

Brownian motion: magic trick!

- ▶ A magic trick: suppose the pollen settles at a height where pressure balances weight, or $mg = PA = \pi PR^2$.




- ▶ Plugging this into D gives the Stokes-Einstein relation:

$$D = \frac{k_B \mathcal{T}}{\pi P R^2} \cdot \frac{mg}{6\pi\eta R} =$$

Brownian motion: magic trick!

- ▶ A magic trick: suppose the pollen settles at a height where pressure balances weight, or $mg = PA = \pi PR^2$.




- ▶ Plugging this into D gives the Stokes-Einstein relation:

$$D = \frac{k_B \mathcal{T}}{\pi P R^2} \cdot \frac{mg}{6\pi\eta R} = \frac{k_B \mathcal{T}}{mg} \cdot \frac{mg}{6\pi\eta R} =$$

Brownian motion: magic trick!

- ▶ A magic trick: suppose the pollen settles at a height where pressure balances weight, or $mg = PA = \pi PR^2$.




- ▶ Plugging this into D gives the Stokes-Einstein relation:

$$D = \frac{k_B \mathcal{T}}{\pi P R^2} \cdot \frac{mg}{6\pi\eta R} = \frac{k_B \mathcal{T}}{mg} \cdot \frac{mg}{6\pi\eta R} = \frac{k_B \mathcal{T}}{6\pi\eta R},$$

Brownian motion: magic trick!

- ▶ A magic trick: suppose the pollen settles at a height where pressure balances weight, or $mg = PA = \pi PR^2$.



- ▶ Plugging this into D gives the Stokes-Einstein relation:

$$D = \frac{k_B \mathcal{T}}{\pi P R^2} \cdot \frac{mg}{6\pi\eta R} = \frac{k_B \mathcal{T}}{mg} \cdot \frac{mg}{6\pi\eta R} = \frac{k_B \mathcal{T}}{6\pi\eta R},$$

one of the main results of Einstein's PhD thesis!

Brownian motion: comments

Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

$$d \sim \sqrt{Dt},$$

Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

$$d \sim \sqrt{Dt}, \quad D = \frac{k_B \mathcal{T}}{6\pi\eta R}.$$

Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

$$d \sim \sqrt{Dt}, \quad D = \frac{k_B \mathcal{T}}{6\pi\eta R}.$$

- ▶ In 1908, Jean Perrin experimentally confirmed Einstein's predictions,

Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

$$d \sim \sqrt{Dt}, \quad D = \frac{k_B \mathcal{T}}{6\pi\eta R}.$$

- ▶ In 1908, Jean Perrin experimentally confirmed Einstein's predictions, hence the existence of atoms.

Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

$$d \sim \sqrt{Dt}, \quad D = \frac{k_B \mathcal{T}}{6\pi\eta R}.$$


- ▶ In 1908, Jean Perrin experimentally confirmed Einstein's predictions, hence the existence of atoms. Another Nobel!

Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

$$d \sim \sqrt{Dt}, \quad D = \frac{k_B \mathcal{T}}{6\pi\eta R}.$$

- ▶ In 1908, Jean Perrin experimentally confirmed Einstein's predictions, hence the existence of atoms. Another Nobel!




Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

$$d \sim \sqrt{Dt}, \quad D = \frac{k_B \mathcal{T}}{6\pi\eta R}.$$

- ▶ In 1908, Jean Perrin experimentally confirmed Einstein's predictions, hence the existence of atoms. Another Nobel!




- ▶ Grains don't settle at a fixed height,

Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

$$d \sim \sqrt{Dt}, \quad D = \frac{k_B \mathcal{T}}{6\pi\eta R}.$$

- ▶ In 1908, Jean Perrin experimentally confirmed Einstein's predictions, hence the existence of atoms. Another Nobel!




- ▶ Grains don't settle at a fixed height, but exist in "dynamic equilibrium".

Brownian motion: comments

- ▶ Thus, a pollen grain wanders a distance d in time t ,

$$d \sim \sqrt{Dt}, \quad D = \frac{k_B \mathcal{T}}{6\pi\eta R}.$$

- ▶ In 1908, Jean Perrin experimentally confirmed Einstein's predictions, hence the existence of atoms. Another Nobel!



- ▶ Grains don't settle at a fixed height, but exist in “dynamic equilibrium”. We used a cheeky hacker shortcut!

Exercise 3: Avogadro's constant I

Exercise 3: Avogadro's constant I

- ▶ You may have seen the **chem version** of the ideal gas law:

Exercise 3: Avogadro's constant I

- ▶ You may have seen the **chem version** of the ideal gas law:

$$PV = \mathcal{N}_{\text{mol}} \mathcal{R} \mathcal{T},$$

Exercise 3: Avogadro's constant I

- ▶ You may have seen the **chem version** of the ideal gas law:

$$PV = \mathcal{N}_{\text{mol}} \mathcal{R} \mathcal{T},$$

where $\mathcal{R} = 8.3 \text{ J/K mol}$ is the **ideal gas constant**.

Exercise 3: Avogadro's constant I

- ▶ You may have seen the **chem version** of the ideal gas law:

$$PV = \mathcal{N}_{\text{mol}} \mathcal{R} \mathcal{T},$$

where $\mathcal{R} = 8.3 \text{ J/K mol}$ is the **ideal gas constant**.

- (a) Recall that one mol is N_A particles, where N_A is **Avogadro's constant**.

Exercise 3: Avogadro's constant I

- ▶ You may have seen the **chem version** of the ideal gas law:

$$PV = \mathcal{N}_{\text{mol}} \mathcal{R} \mathcal{T},$$

where $\mathcal{R} = 8.3 \text{ J/K mol}$ is the **ideal gas constant**.

- (a) Recall that one mol is N_A particles, where N_A is **Avogadro's constant**. Show that $N_A = \mathcal{R}/k_B$.

Exercise 3: Avogadro's constant I

- ▶ You may have seen the **chem version** of the ideal gas law:

$$PV = \mathcal{N}_{\text{mol}} \mathcal{R} \mathcal{T},$$

where $\mathcal{R} = 8.3 \text{ J/K mol}$ is the **ideal gas constant**.

- (a) Recall that one mol is N_A particles, where N_A is **Avogadro's constant**. Show that $N_A = \mathcal{R}/k_B$.
- (b) In 1905, \mathcal{R} was known but N_A was not.

Exercise 3: Avogadro's constant I

- ▶ You may have seen the **chem version** of the ideal gas law:

$$PV = \mathcal{N}_{\text{mol}} \mathcal{R} \mathcal{T},$$

where $\mathcal{R} = 8.3 \text{ J/K mol}$ is the **ideal gas constant**.

- (a) Recall that one mol is N_A particles, where N_A is **Avogadro's constant**. Show that $N_A = \mathcal{R}/k_B$.
- (b) In 1905, \mathcal{R} was known but N_A was not. Argue that

Exercise 3: Avogadro's constant I

- ▶ You may have seen the **chem version** of the ideal gas law:

$$PV = \mathcal{N}_{\text{mol}} \mathcal{R} \mathcal{T},$$

where $\mathcal{R} = 8.3 \text{ J/K mol}$ is the **ideal gas constant**.

- (a) Recall that one mol is N_A particles, where N_A is **Avogadro's constant**. Show that $N_A = \mathcal{R}/k_B$.
- (b) In 1905, \mathcal{R} was known but N_A was not. Argue that

$$N_A \sim \frac{t}{d^2} \cdot \frac{\mathcal{R} \mathcal{T}}{6\pi\eta R}.$$

Exercise 3: Avogadro's constant I

- ▶ You may have seen the **chem version** of the ideal gas law:

$$PV = \mathcal{N}_{\text{mol}} \mathcal{R} \mathcal{T},$$

where $\mathcal{R} = 8.3 \text{ J/K mol}$ is the **ideal gas constant**.

- (a) Recall that one mol is N_A particles, where N_A is **Avogadro's constant**. Show that $N_A = \mathcal{R}/k_B$.
- (b) In 1905, \mathcal{R} was known but N_A was not. Argue that

$$N_A \sim \frac{t}{d^2} \cdot \frac{\mathcal{R} \mathcal{T}}{6\pi\eta R}.$$

This was one of no fewer than **five** methods Einstein proposed for measuring Avogadro's constant!

Exercise 3: Avogadro's constant I (solution)

Exercise 3: Avogadro's constant I (solution)

(a) We equate the physics and chem version to get:

Exercise 3: Avogadro's constant I (solution)

(a) We equate the physics and chem version to get:

$$\begin{aligned}\mathcal{N}k_B\mathcal{T} &= PV = \mathcal{N}_{\text{mol}}\mathcal{R}\mathcal{T} \\ \implies \mathcal{N}k_B &= \mathcal{N}_{\text{mol}}\mathcal{R}.\end{aligned}$$

Exercise 3: Avogadro's constant I (solution)

(a) We equate the physics and chem version to get:

$$\begin{aligned}\mathcal{N}k_B\mathcal{T} &= PV = \mathcal{N}_{\text{mol}}\mathcal{R}\mathcal{T} \\ \implies \mathcal{N}k_B &= \mathcal{N}_{\text{mol}}\mathcal{R}.\end{aligned}$$

Since $\mathcal{N}_{\text{mol}}N_A = \mathcal{N}$,

Exercise 3: Avogadro's constant I (solution)

(a) We equate the physics and chem version to get:

$$\begin{aligned}\mathcal{N}k_B\mathcal{T} &= PV = \mathcal{N}_{\text{mol}}\mathcal{R}\mathcal{T} \\ \implies \mathcal{N}k_B &= \mathcal{N}_{\text{mol}}\mathcal{R}.\end{aligned}$$

Since $\mathcal{N}_{\text{mol}}N_A = \mathcal{N}$, we find

$$N_Ak_B = \mathcal{R}.$$

Exercise 3: Avogadro's constant I (solution)

(a) We equate the physics and chem version to get:

$$\begin{aligned}\mathcal{N}k_B\mathcal{T} &= PV = \mathcal{N}_{\text{mol}}\mathcal{R}\mathcal{T} \\ \implies \mathcal{N}k_B &= \mathcal{N}_{\text{mol}}\mathcal{R}.\end{aligned}$$

Since $\mathcal{N}_{\text{mol}}N_A = \mathcal{N}$, we find

$$N_Ak_B = \mathcal{R}.$$

(b) From the Stokes-Einstein relation,

Exercise 3: Avogadro's constant I (solution)

(a) We equate the physics and chem version to get:

$$\begin{aligned}\mathcal{N}k_B\mathcal{T} &= PV = \mathcal{N}_{\text{mol}}\mathcal{R}\mathcal{T} \\ \implies \mathcal{N}k_B &= \mathcal{N}_{\text{mol}}\mathcal{R}.\end{aligned}$$

Since $\mathcal{N}_{\text{mol}}N_A = \mathcal{N}$, we find

$$N_Ak_B = \mathcal{R}.$$

(b) From the Stokes-Einstein relation,

$$\frac{k_B\mathcal{T}}{6\pi\eta R} \sim \frac{d^2}{t} \implies$$

Exercise 3: Avogadro's constant I (solution)

(a) We equate the physics and chem version to get:

$$\begin{aligned}\mathcal{N}k_B\mathcal{T} &= PV = \mathcal{N}_{\text{mol}}\mathcal{R}\mathcal{T} \\ \implies \mathcal{N}k_B &= \mathcal{N}_{\text{mol}}\mathcal{R}.\end{aligned}$$

Since $\mathcal{N}_{\text{mol}}N_A = \mathcal{N}$, we find

$$N_A k_B = \mathcal{R}.$$

(b) From the Stokes-Einstein relation,

$$\frac{k_B\mathcal{T}}{6\pi\eta R} \sim \frac{d^2}{t} \quad \implies \quad N_A \sim \frac{t}{d^2} \cdot \frac{\mathcal{R}\mathcal{T}}{6\pi\eta R}.$$


Exercise 3: Avogadro's constant II

Exercise 3: Avogadro's constant II

- ▶ Below, we show some of Perrin's data ($R = 0.5 \mu\text{m}$):


Exercise 3: Avogadro's constant II

- Below, we show some of Perrin's data ($R = 0.5 \mu\text{m}$):



Exercise 3: Avogadro's constant II


- ▶ Below, we show some of Perrin's data ($R = 0.5 \mu\text{m}$):



- ▶ Observations are made every 30 s,

Exercise 3: Avogadro's constant II


- ▶ Below, we show some of Perrin's data ($R = 0.5 \mu\text{m}$):



- ▶ Observations are made every 30 s, and lines ruled every $3 \mu\text{m}$.

Exercise 3: Avogadro's constant II


- Below, we show some of Perrin's data ($R = 0.5 \mu\text{m}$):



- Observations are made every 30 s, and lines ruled every $3 \mu\text{m}$. The water had $T = 290 \text{ K}$ and $\eta = 0.011 \text{ kg/m s}$.

Exercise 3: Avogadro's constant II

- Below, we show some of Perrin's data ($R = 0.5 \mu\text{m}$):



- Observations are made every 30 s, and lines ruled every $3 \mu\text{m}$. The water had $T = 290 \text{ K}$ and $\eta = 0.011 \text{ kg/m s}$.
- (c) Using this data, find N_A .

Exercise 3: Avogadro's constant II (solution)

Exercise 3: Avogadro's constant II (solution)

(c) We have 20 points, spread over 5 divisions or so.

Exercise 3: Avogadro's constant II (solution)

- (c) We have 20 points, spread over 5 divisions or so.
Thus, $t = 30 \times 20$ s, and $d \sim 5 \times 3 \mu\text{m}$.

Exercise 3: Avogadro's constant II (solution)

- (c) We have 20 points, spread over 5 divisions or so.
Thus, $t = 30 \times 20$ s, and $d \sim 5 \times 3 \mu\text{m}$.
- ▶ We have $\mathcal{T} = 290$ K, $R = 0.5 \mu\text{ m}$, $\eta = 0.0011$ kg/m s.

Exercise 3: Avogadro's constant II (solution)

- (c) We have 20 points, spread over 5 divisions or so.
Thus, $t = 30 \times 20$ s, and $d \sim 5 \times 3 \mu\text{m}$.
- ▶ We have $\mathcal{T} = 290$ K, $R = 0.5 \mu\text{ m}$, $\eta = 0.0011$ kg/m s.
Plugging into (b) and using SI units everywhere,

Exercise 3: Avogadro's constant II (solution)

- (c) We have 20 points, spread over 5 divisions or so.
Thus, $t = 30 \times 20$ s, and $d \sim 5 \times 3 \mu\text{m}$.
- We have $\mathcal{T} = 290$ K, $R = 0.5 \mu\text{ m}$, $\eta = 0.0011$ kg/m s.
Plugging into (b) and using SI units everywhere,

$$\begin{aligned} N_A &\sim \frac{t}{d^2} \cdot \frac{\mathcal{RT}}{6\pi\eta R} \\ &= \frac{30 \times 20}{(5 \times 3 \times 10^{-6})^2} \frac{8.3 \cdot 290}{6\pi \cdot 0.0011 \cdot (0.5 \times 10^{-6})} \\ &\approx 6.2 \times 10^{23}. \end{aligned}$$

Exercise 3: Avogadro's constant II (solution)

- (c) We have 20 points, spread over 5 divisions or so.
Thus, $t = 30 \times 20$ s, and $d \sim 5 \times 3 \mu\text{m}$.
- We have $\mathcal{T} = 290$ K, $R = 0.5 \mu\text{ m}$, $\eta = 0.0011$ kg/m s.
Plugging into (b) and using SI units everywhere,

$$\begin{aligned} N_A &\sim \frac{t}{d^2} \cdot \frac{\mathcal{RT}}{6\pi\eta R} \\ &= \frac{30 \times 20}{(5 \times 3 \times 10^{-6})^2} \frac{8.3 \cdot 290}{6\pi \cdot 0.0011 \cdot (0.5 \times 10^{-6})} \\ &\approx 6.2 \times 10^{23}. \end{aligned}$$

The modern value is $N_A = 6.022 \times 10^{23}$.

Exercise 3: Avogadro's constant II (solution)

- (c) We have 20 points, spread over 5 divisions or so.
Thus, $t = 30 \times 20$ s, and $d \sim 5 \times 3 \mu\text{m}$.
- We have $\mathcal{T} = 290$ K, $R = 0.5 \mu\text{ m}$, $\eta = 0.0011$ kg/m s.
Plugging into (b) and using SI units everywhere,

$$\begin{aligned} N_A &\sim \frac{t}{d^2} \cdot \frac{\mathcal{RT}}{6\pi\eta R} \\ &= \frac{30 \times 20}{(5 \times 3 \times 10^{-6})^2} \frac{8.3 \cdot 290}{6\pi \cdot 0.0011 \cdot (0.5 \times 10^{-6})} \\ &\approx 6.2 \times 10^{23}. \end{aligned}$$

The modern value is $N_A = 6.022 \times 10^{23}$. Sweet!

Exercise 3: Avogadro's constant III

Exercise 3: Avogadro's constant III

- ▶ Of course, the conventional definition of N_A is

Exercise 3: Avogadro's constant III

- ▶ Of course, the conventional definition of N_A is the number of carbon atoms in 12 g of carbon-12.

Exercise 3: Avogadro's constant III

- ▶ Of course, the conventional definition of N_A is the number of carbon atoms in 12 g of carbon-12.
- (d) From $N_A \approx 6 \times 10^{23}$, estimate a carbon-12 atom's mass.

Exercise 3: Avogadro's constant III


- ▶ Of course, the conventional definition of N_A is the number of carbon atoms in 12 g of carbon-12.
- (d) From $N_A \approx 6 \times 10^{23}$, estimate a carbon-12 atom's mass.
 - ▶ Most of the atom's mass is concentrated in its nucleus,

Exercise 3: Avogadro's constant III

- ▶ Of course, the conventional definition of N_A is the number of carbon atoms in 12 g of carbon-12.
- (d) From $N_A \approx 6 \times 10^{23}$, estimate a **carbon-12 atom's mass**.
- ▶ Most of the atom's mass is concentrated in its nucleus, made of (roughly) equally weighted protons and neutrons.


Exercise 3: Avogadro's constant III

- ▶ Of course, the conventional definition of N_A is the number of carbon atoms in 12 g of carbon-12.
- (d) From $N_A \approx 6 \times 10^{23}$, estimate a **carbon-12 atom's mass**.
- ▶ Most of the atom's mass is concentrated in its nucleus, made of (roughly) equally weighted protons and neutrons.



Exercise 3: Avogadro's constant III

- ▶ Of course, the conventional definition of N_A is the number of carbon atoms in 12 g of carbon-12.
- (d) From $N_A \approx 6 \times 10^{23}$, estimate a **carbon-12 atom's mass**.
 - ▶ Most of the atom's mass is concentrated in its nucleus, made of (roughly) equally weighted protons and neutrons.



- (e) What is the approximate **mass of a nucleon**?

Exercise 3: Avogadro's constant III (solution)

Exercise 3: Avogadro's constant III (solution)

(d) The atom mass m_C is the total mass divided by N_A :

Exercise 3: Avogadro's constant III (solution)

(d) The atom mass m_C is the total mass divided by N_A :

$$m_C = \frac{12 \text{ g}}{N_A} \approx 2 \times 10^{-23} \text{ g} = 2 \times 10^{-26} \text{ kg}.$$

Exercise 3: Avogadro's constant III (solution)

(d) The atom mass m_C is the total mass divided by N_A :

$$m_C = \frac{12 \text{ g}}{N_A} \approx 2 \times 10^{-23} \text{ g} = 2 \times 10^{-26} \text{ kg}.$$

(e) We have $m_C \approx 12m_{\text{nucleon}}$,

Exercise 3: Avogadro's constant III (solution)

(d) The atom mass m_C is the total mass divided by N_A :

$$m_C = \frac{12 \text{ g}}{N_A} \approx 2 \times 10^{-23} \text{ g} = 2 \times 10^{-26} \text{ kg}.$$

(e) We have $m_C \approx 12m_{\text{nucleon}}$, and hence:

Exercise 3: Avogadro's constant III (solution)

(d) The atom mass m_C is the total mass divided by N_A :

$$m_C = \frac{12 \text{ g}}{N_A} \approx 2 \times 10^{-23} \text{ g} = 2 \times 10^{-26} \text{ kg}.$$

(e) We have $m_C \approx 12m_{\text{nucleon}}$, and hence:

$$m_{\text{nucleon}} \approx \frac{m_C}{12} \approx 1.7 \times 10^{-27} \text{ kg}.$$

Questions?

Questions?

Thanks everyone, you've been grand.

Questions?

Thanks everyone, you've been grand.

Go forth and hack physics!

Questions?

Thanks everyone, you've been grand.

Go forth and hack physics!

