

Inflation in scalar-tensor gravity

David Wakeham
School of Physics, University of Melbourne

Supervisor
Prof. Raymond Volkas

Nov 09, 2016




COEPP
ARC Centre of Excellence for
Particle Physics at the Terascale



Australian Government
Australian Research Council

Big Bang cosmology


- **Cosmological Principle:** at large scales, universe is the same everywhere and in all directions.
- Three possibilities: uniform sphere, saddle or plane.
- Geometries scale with time. Track with scale factor $a(t)$.



- Scaling is driven by energy and pressure of large-scale stuff.

Troubles with tuning


- Flatness problem: indications we live in flat space.
- This requires fine tuning of amount of matter in universe.
- Horizon problem: the CMB is highly uniform.



- But CMB splits into $\sim 10^4$ patches which never interacted!
Necessitates fine tuning of patch conditions.

Inflation I



- Inflation is burst of **accelerating expansion** after Big Bang.
- Rapid growth smooths out **lumps**, creating **flat, uniform** space.



- Tiny **quantum lumps** later become **large-scale structure**.

Inflation II


- To solve horizon problem, just **grow light cones**:



- Light cone width $\sim \int 1/\dot{a}$. Usually, **$1/\dot{a}$ increases with time**, and fixed by observation at time CMB forms.
- To grow cones, add big contribution early on. Simple way: **increase $1/\dot{a}$ as we go back in time**.
- Since $\partial_{-t}\dot{a} = \ddot{a}/a^2$, this is just **accelerating expansion!**

The scalar roller coaster


- A scalar ϕ with $V > T$ can cause inflation. Called **inflaton**.
- If $V \gg T$, have **slow-roll inflation** governed by shape of V .
- Slow roll on **flat patches**, ends on inclines.



- A successful model must
 - increase scale factor by $\sim e^{60}$ (60 e-folds);
 - predict right **power law distribution** for quantum lumps.

Higgs inflation

- Standard model has a unique scalar: **Higgs boson**.
- The Higgs gives mass to leptons, quarks, and weak force carriers W^\pm and Z via **spontaneous symmetry breaking**.




- Sadly, **can't drive inflation!** Mexican hat isn't flat enough.
- Bezrukov and Shaposhnikov [2008] coupled Higgs **directly** to gravity to get viable inflation. Called **scalar-tensor gravity**.

Scalar-tensor gravity

- Simplest approach to scalar-tensor gravity is **action principle**.
- Modify **curvature term** in Lagrangian for general relativity:

$$\mathcal{L}_{\text{GR}} \supset \kappa \cdot \sqrt{-g} \mathcal{R} \quad \longrightarrow \quad \mathcal{L}_{\text{ST}} \supset f(\phi) \cdot \sqrt{-g} \mathcal{R}.$$

- Can think of f as **warping space** on top of curvature.



- If $\phi \rightarrow v$, recover general relativity for $f(v) = \kappa$.

Rescaling and frames

- By locally rescaling metric, get new field χ governed by GR:

$$g_{\mu\nu} \rightarrow \tilde{g}_{\mu\nu} = \Omega(f)^2 g_{\mu\nu}, \quad \phi \rightarrow \chi.$$




Figure: Rescaling metric = variable magnification.


- Choosing f cleverly and rescaling allows χ to viably slow-roll.
- Metric $g_{\mu\nu}$ is called **Jordan frame**, $\tilde{g}_{\mu\nu}$ **Einstein frame**.
- Dictionary** connects them but **not physically equivalent**.

de Sitter inflation

- If Jordan \neq Einstein, motivates **inflation in Jordan frame**. Consider arbitrary coupling f and potential V .
- Simplest case is **de Sitter evolution**. Exponential scale factor:

$$a(t) = e^{Ht}, \quad H \equiv \dot{a}/a = \text{const}, \quad (\text{note } \ddot{a} > 0).$$

- Scalar-tensor equations imply **quadratic for $\dot{\phi}$** . Obtain **consistent patches** (real $\dot{\phi}$) for discriminant $\Delta \geq 0$.



- This **constrains coupling f and potential V** .

Perturbing de Sitter

- Earlier, inflation defined as accelerating expansion.
Equivalently, slow change in Hubble parameter $H \equiv \dot{a}/a$.
- Suggests perturbation expansion around constant H :

$$\begin{aligned}H(t) &= H_0 + \lambda H_1(t) + \dots \\ \phi(t) &= \phi_0(t) + \lambda \phi_1(t) + \dots.\end{aligned}$$

In other words, H_0 , ϕ_0 terms represent de Sitter inflation.

- Plug expansion into scalar-tensor equations and get homogeneous matrix DE for first-order terms:

$$\frac{d}{dt} \begin{bmatrix} \phi_1 \\ H_1 \end{bmatrix} = M(t) \begin{bmatrix} \phi_1 \\ H_1 \end{bmatrix}.$$

- Still have consistency constraint at zeroth order.

Jordan frame slow-roll

- That is classical story, but what about quantum lumps?
Requires QFT on curved, scalar-tensor background...
- For normal slow-roll, need parameters ϵ_1 and ϵ_2 to be small and approximately constant to get power-law behaviour:

$$\epsilon_1 = \frac{\dot{H}}{H^2}, \quad \epsilon_2 = \frac{\ddot{\phi}}{H\dot{\phi}}.$$


- In Jordan frame, two more slow-roll parameters ϵ_3 and ϵ_4 :

$$\epsilon_3 = \frac{\dot{f}}{2Hf}, \quad \epsilon_4 = \frac{\dot{\alpha}}{2H\alpha}, \quad \text{where } \alpha \equiv 1 + 3(f')^2/f.$$


- Distribution determined by linear combination of ϵ_i .

Toy models


- Explored scalar-tensor inflation for various **toy models**:
 - Higgs inflation;
 - linearly coupled massive scalar;
 - coupling driven inflation.



(a) Higgs inflation.



(b) Massive scalar.



(c) Gaussian coupling.

- Massive scalar can **oscillate resonantly**, like **reheating**.
- By tuning parameters, usually **possible to get 60 e-folds**.
- “Bottom up” approach does not give **Jordan frame slow-roll**.

Top down approach

- Try imposing slow-roll constraints on ϵ_3, ϵ_4 directly:

$$\frac{\dot{f}}{2Hf} = \epsilon_3^{(0)} + \dots, \quad \frac{\dot{\alpha}}{2H\alpha} = \epsilon_4^{(0)} + \dots.$$

$\epsilon_3^{(0)}$ and $\epsilon_4^{(0)}$ are small constants, “...” higher order terms.

- Leads to explicit form $f(\phi_0(t))$, coupling as function of time.
- Use this to calculate ϵ_2 . Find ϵ_2 is also small and roughly constant, with some suggestive cancellations.
- In principle, can work backwards to determine $f(\phi)$ and $V(\phi)$, and hence ϵ_1 and first-order dynamics. Work in progress!

Summary

- Inflation solves tuning problems with Big Bang model.
- Scalar-tensor gravity gives wider range of inflaton candidates.
- If Jordan \neq Einstein, motivates inflation in Jordan frame.
Perturbing around de Sitter inflation is one approach.
- Toy models had interesting dynamics, but realistic inflation requires top down approach.
- Ultimate goal: richer interplay of cosmology/particle physics.

Supplementary slides

FRW metric and Friedmann equations

- Cosmological principle implies **FRW metric**:

$$ds^2 = -dt^2 + a^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right].$$

Geometry given by $k \in \{0, \pm 1\}$: $k = 1$ is sphere, $k = -1$ is saddle, and $k = 0$ is flat space.


- **Scale factor** $a(t)$ controls expansion of space.
- Responds to **energy** and **pressure** via **Friedmann equations**:

$$\begin{aligned} \left(\frac{\dot{a}}{a} \right)^2 &= \frac{8\pi G}{3}\rho - \frac{k}{a^2} \\ \frac{\ddot{a}}{a} &= -\frac{4\pi G}{3}(\rho + 3P). \end{aligned}$$

Scalar-tensor cosmology

- Scalar-tensor action gives equivalent of Einstein's equation.
- Plug in FRW to get equivalent of Friedmann equations:

$$H^2 + H\dot{f} \propto \rho, \quad H\dot{f} - 2\dot{H} - \ddot{f} \propto (\rho + P).$$



- Density and pressure related to ϕ , V by

$$\rho = T + V = \frac{1}{2}\dot{\phi}^2 + V, \quad P = \frac{1}{2}\dot{\phi}^2 - V.$$

Inflationary spectra

- Lumps are **momentum modes** for inhomogeneous corrections.
- Two types: **scalar** and **tensor**. Spectra obey **power-law**:

$$\Delta_s^2 = A_s k^{n_s - 1}, \quad \Delta_t^2 = A_t k^{n_t}.$$

- Key observables** are $n_s \simeq 0.97$ and $r \equiv A_t/A_s \lesssim 0.11$.




Figure: Life cycle of a momentum mode.

Jordan frame spectra

- Rough idea for spectral calculations:
 - ① add inhomogeneous correction, $\phi(t, \mathbf{x}) = \bar{\phi}(t) + \delta\phi(t, \mathbf{x})$;
 - ② expand inflaton $S[\phi, \dot{\phi}]$ action to second order, $S^{(2)}$;
 - ③ set corresponding variation to zero, $\delta S^{(2)} = 0$;
 - ④ quantise Mukhanov-Sasaki equation in Fourier space;
 - ⑤ evolve to horizon crossing.
- Calculation in Jordan frame uses rescaling dictionary:

$$\mathcal{J} \rightarrow \mathcal{E} \rightarrow \mathcal{J}.$$

- Jordan frame spectra:

$$n_s \simeq 1 + 2(2\epsilon_1 - \epsilon_2 + \epsilon_3 - \epsilon_4)$$

$$n_t \simeq 2(\epsilon_1 - \epsilon_3)$$

$$r \simeq 4\pi \left(-\frac{\epsilon_1}{16\pi fG} + 3\epsilon_3^2 \right).$$

Higgs inflation in Einstein frame

- Bezrukov and Shaposhnikov [2008] set $\phi = \text{Higgs}$ and

$$f(\phi) = \frac{1}{2} \left[M_{\text{pl}}^2 + \xi (\phi^2 - v^2) \right].$$

- In Einstein frame, making $\xi \sim 50000\sqrt{\lambda}$ large flattens potential enough to satisfy constraints!
- Get $n_s \simeq 0.97$ and $r \simeq 0.003$, consistent with observation.




Figure: Flattening outer brim of Mexican hat.