

Continuous Positive Airway Pressure (CPAP) Therapy For Obstructive Sleep Apnea (OSA)

240.4 Links in PDF documents are not guaranteed to work. To follow a web link, please use the MCD Website.

Tracking Information

Publication Number

100-3

Manual Section Number

240.4

Manual Section Title

Continuous Positive Airway Pressure (CPAP) Therapy For Obstructive Sleep Apnea (OSA)

Version Number

3

Effective Date of this Version

03/13/2008

Implementation Date

08/04/2008

Description Information

Benefit Category

Durable Medical Equipment

Please Note: This may not be an exhaustive list of all applicable Medicare benefit categories for this item or service.

Item/Service Description

A. General

Continuous Positive Airway Pressure (CPAP) is a non-invasive technique for providing single levels of air pressure from a flow generator, via a nose mask, through the nares. The purpose is to prevent the collapse of the oropharyngeal walls and the obstruction of airflow during sleep, which occurs in obstructive sleep apnea (OSA).

The apnea hypopnea index (AHI) is equal to the average number of episodes of apnea and hypopnea per hour. The respiratory disturbance index (RDI) is equal to the average number of respiratory disturbances per hour.

Apnea is defined as a cessation of airflow for at least 10 seconds. Hypopnea is defined as an abnormal respiratory event lasting at least 10 seconds with at least a 30% reduction in thoracoabdominal movement or airflow as compared to baseline, and with at least a 4% oxygen desaturation.

The AHI and/or RDI may be measured by polysomnography (PSG) in a facility-based sleep study laboratory, or by a Type II home sleep test (HST) monitor, a Type III HST monitor, or a Type IV HST monitor measuring at least 3 channels.

Indications and Limitations of Coverage

B. Nationally Covered Indications

Effective for claims with dates of service on and after March 13, 2008, the Centers for Medicare & Medicaid Services (CMS) determines that CPAP therapy when used in adult patients with OSA is considered reasonable and necessary under the following situations:

1. The use of CPAP is covered under Medicare when used in adult patients with OSA. Coverage of CPAP is initially limited to a 12-week period to identify beneficiaries diagnosed with OSA as subsequently described who benefit from CPAP. CPAP is subsequently covered only for those beneficiaries diagnosed with OSA who benefit from CPAP during this 12-week period.
2. The provider of CPAP must conduct education of the beneficiary prior to the use of the CPAP device to ensure that the beneficiary has been educated in the proper use of the device. A caregiver, for example a family member, may be compensatory, if consistently available in the beneficiary's home and willing and able to safely operate the CPAP device.
3. A positive diagnosis of OSA for the coverage of CPAP must include a clinical evaluation and a positive:
 - a. attended PSG performed in a sleep laboratory; or
 - b. unattended HST with a Type II home sleep monitoring device; or
 - c. unattended HST with a Type III home sleep monitoring device; or
 - d. unattended HST with a Type IV home sleep monitoring device that measures at least 3 channels.
4. The sleep test must have been previously ordered by the beneficiary's treating physician and furnished under appropriate physician supervision.
5. An initial 12-week period of CPAP is covered in adult patients with OSA if either of the following criterion using the AHI or RDI are met:
 - a. AHI or RDI greater than or equal to 15 events per hour, or
 - b. AHI or RDI greater than or equal to 5 events and less than or equal to 14 events per hour with documented symptoms of excessive daytime sleepiness, impaired cognition, mood disorders or insomnia, or documented hypertension, ischemic heart disease, or history of stroke.

6. The AHI or RDI is calculated on the average number of events of per hour. If the AHI or RDI is calculated based on less than 2 hours of continuous recorded sleep, the total number of recorded events to calculate the AHI or RDI during sleep testing must be at a minimum the number of events that would have been required in a 2-hour period.
7. Apnea is defined as a cessation of airflow for at least 10 seconds. Hypopnea is defined as an abnormal respiratory event lasting at least 10 seconds with at least a 30% reduction in thoracoabdominal movement or airflow as compared to baseline, and with at least a 4% oxygen desaturation.
8. Coverage with Evidence Development (CED): Medicare provides the following limited coverage for CPAP in adult beneficiaries who do not qualify for CPAP coverage based on criteria 1-7 above. A clinical study seeking Medicare payment for CPAP provided to a beneficiary who is an enrolled subject in that study must address one or more of the following questions
 - a. In Medicare-aged subjects with clinically identified risk factors for OSA, how does the diagnostic accuracy of a clinical trial of CPAP compare with PSG and Type II, III & IV HST in identifying subjects with OSA who will respond to CPAP?
 - b. In Medicare-aged subjects with clinically identified risk factors for OSA who have not undergone confirmatory testing with PSG or Type II, III & IV HST, does CPAP cause clinically meaningful harm?
 - c. The study must meet the following additional standards:
 - d. The principal purpose of the research study is to test whether a particular intervention potentially improves the participants' health outcomes.
 - e. The research study is well-supported by available scientific and medical information or it is intended to clarify or establish the health outcomes of interventions already in common clinical use.
 - f. The research study does not unjustifiably duplicate existing studies.
 - g. The research study design is appropriate to answer the research question being asked in the study.
 - h. The research study is sponsored by an organization or individual capable of executing the proposed study successfully.
 - i. The research study is in compliance with all applicable Federal regulations concerning the protection of human subjects found at 45 CFR Part 46. If a study is Food and Drug Administration-regulated, it also must be in compliance with 21 CFR Parts 50 and 56.
 - j. All aspects of the research study are conducted according to the appropriate standards of scientific integrity.
 - k. The research study has a written protocol that clearly addresses, or incorporates by reference, the Medicare standards.
 - l. The clinical research study is not designed to exclusively test toxicity or disease pathophysiology in healthy individuals. Trials of all medical technologies measuring therapeutic outcomes as one of the objectives meet this standard only if the disease or condition being studied is life-threatening as defined in 21 CFR § 312.81(a) and the patient has no other viable treatment options.

- m. The clinical research study is registered on the ClinicalTrials.gov Web site by the principal sponsor/investigator prior to the enrollment of the first study subject.
- n. The research study protocol specifies the method and timing of public release of all pre-specified outcomes to be measured, including release of outcomes if outcomes are negative or study is terminated early. The results must be made public within 24 months of the end of data collection. If a report is planned for publication in a peer-reviewed journal, then that initial release may be an abstract that meets the requirements of the International Committee of Medical Journal Editors. However, a full report of the outcomes must be made public no later than 3 years after the end of data collection.
- o. The research study protocol must explicitly discuss subpopulations affected by the treatment under investigation, particularly traditionally underrepresented groups in clinical studies, how the inclusion and exclusion criteria affect enrollment of these populations, and a plan for the retention and reporting of said populations in the trial. If the inclusion and exclusion criteria are expected to have a negative effect on the recruitment or retention of underrepresented populations, the protocol must discuss why these criteria are necessary.
- p. The research study protocol explicitly discusses how the results are or are not expected to be generalizable to the Medicare population to infer whether Medicare patients may benefit from the intervention. Separate discussions in the protocol may be necessary for populations eligible for Medicare due to age, disability, or Medicaid eligibility.

C. Nationally Non-covered Indications

Effective for claims with dates of services on and after March 13, 2008, other diagnostic tests for the diagnosis of OSA, other than those noted above for prescribing CPAP, are not sufficient for the coverage of CPAP.

D. Other

N/A

(This NCD last reviewed March 2008.)

Cross Reference

See Section 280.1, [Durable Medical Equipment Reference List](#) 
[CED page](#) 

Transmittal Information

Transmittal Number

96

Coverage Transmittal Link

<https://www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/Downloads/R96NCD.pdf> 

Revision History

10/2008 - NOTE: Transmittal 96, dated October 15, 2008 rescinds and replaces Transmittal 94, dated August 29, 2008. The only change is a clarification to the verbiage under the Policy section of the Business Requirements at I.B.3. All other information remains the same. Additionally, this clarification does not in any way constitute revisions to the corresponding NCD at 240.4 of Pub 100-03. ([TN 96](#)) (CR6048)

08/2008 - NOTE: Transmittal 94, dated August 29, 2008 rescinds and replaces Transmittal 91, dated July 25, 2008. The only change is a clarification to the verbiage under the Policy section of the Business Requirements at I.B.3. All other information remains the same. Additionally, this clarification does not in any way constitute revisions to the corresponding NCD at 240.4 of Pub 100-03. ([TN 94](#)) (CR6048)

07/2008 - After reconsideration, coverage for CPAP therapy for OSA is expanded to include a positive diagnosis of OSA made using a home sleep test under specified criteria. Effective date: 03/13/2008. Implementation date: 08/04/2008. ([TN 91](#)) (CR6048)

05/2005 - Found insufficient evidence to conclude that unattended portable multi-channel sleep study testing is reasonable and necessary in diagnosis of OSA for CPAP therapy, and these tests remain non-covered for this purpose. Effective date 04/04/2005. Implementation date 06/06/2005. ([TN 35](#)) (CR 3843)

12/2001 - Expanded coverage for use in obstructive sleep apnea. Effective and implementation dates 04/01/2002. ([TN 150](#)) (CR 1949)

06/1987 - Revised indications for treatment to include patients for whom surgery is a likely alternative to CPAP, and deleted documentation requirements. Effective date 01/12/1987. (TN 15)

12/1986 - Provided coverage when used in adult patients with moderate or severe obstructive sleep apnea who have failed to obtain relief from other non-invasive therapies. Effective date 01/12/1987. (TN 12)

National Coverage Analyses (NCAs)

This NCD has been or is currently being reviewed under the National Coverage Determination process. The following are existing associations with NCAs, from the National Coverage Analyses database.

- [Original Consideration for Continuous Positive Airway Pressure \(CPAP\) Therapy for Obstructive Sleep Apnea \(OSA\) \(CAG-00093N\)](#)
- [First reconsideration for Continuous Positive Airway Pressure \(CPAP\) Therapy for Obstructive Sleep Apnea \(OSA\) \(CAG-00093R\)](#)
- [Second reconsideration for Continuous Positive Airway Pressure \(CPAP\) Therapy for Obstructive Sleep Apnea \(OSA\) \(CAG-00093R2\)](#)

Additional Information

Other Versions

Title	Version	Effective Between	
Continuous Positive Airway Pressure (CPAP) Therapy For Obstructive Sleep Apnea (OSA)	3	03/13/2008 - N/A	You are here
Continuous Positive Airway Pressure (CPAP) Therapy For Obstructive Sleep Apnea (OSA)	2	04/04/2005 - 03/13/2008	View
Continuous Positive Airway Pressure (CPAP)	1	04/01/2002 - 04/04/2005	View