

CDSCO AI REGULATORY REPORT

Generated on: 2026-04-18 16:50:02.701564

FILE 1

Document Type: {'type': 'serious adverse event report', 'confidence': 0.386}

Completeness: 1.0

Severity: serious adverse event

Summary (Original)

Ofatumumab was granted regular approval in 2007, on the basis of superior PFS. Patients in the Hx-CD20-406 trial frequently had a history of severe infections. Infusional toxicity was common, manifesting as fever, dyspnea, and rash despite premedication. Myocardial infarction or angina was noted in four patients within two days of a dose of ofatumumAB. FDA recommended that monthly disease status assessments for progression occur during the first year for all patients enrolled into OMB110911. In a population with unmet need, an observed response rate where the lower bound of the 95% CI for the ORR was at least 25% would be of interest. Some of the regimens in Table 3 are based on randomized phase 3 trials that have not been submitted to FDA in support of an NDA or BLA. Inclusion of these regimens does not indicate that they are approved or recommended by FDA. The applicant's analyses of OS in patients alive at week 12 are not informative and potentially misleading. Because these analyses a a patient's OS a week 12 is not informative, these analyses could be misleading. Some studies were submitted as synopses or interim reports. Only studies 402 and 406 are included in the efficacy and primary safety analyses. The NCIWG criteria (Cheson 1996) were used to define response in study 406. Responders were defined as those with a complete response (CR) or nodular partial response (nPR) Non-responders were those with stable disease (SD) or progressive disease (PD)

Summary (Anonymized)

Ofatumumab is only being considered for CLL patients without adequate available therapy. Post-marketing studies are underway to further characterize the clinical benefit(s) of ofatumumbe (see section 1.4). This reviewer believes that based on the approval of other drugs for the treatment of CLL (and cancer in general), that ofatumomab has a tolerable risk-benefit profile for the intended population. The primary endpoint of study 406 was [NAME] Committee (IRC)-assessed ORR at 24 weeks. A total of 69 patients withdrew during the treatment period. 85 patients completed treatment and were withdrawn during follow-up. Infections (including infectious deaths) occurred frequent. There were some inconsistencies applied to multiple occurrences of tingling in areas of the face. The ORRs across the body weight subgroups (either dichotomized or trichotomized) were fairly consistent. The trend in the DR group of decreasing DOR for increasing body weight was not seen in the dichotomous group. The study Hx-CD20-406 was the only study to evaluate ofatumumab at the doses and schedule for which the applicant is seeking approval. It was the first study to assess the safety and effectiveness of the drug at the dose and schedule it was being evaluated for. It is the first to evaluate the drug's safety and efficacy at a higher dose and at a lower dose. The drug was not approved when the double refractory (refractory to fludarabine and alemtuzumab) CLL patient population was deemed by FDA as having a

Summary Metrics

Compression Ratio: 0.004

Reduction: 99.6%

FILE 2

Document Type: {'type': 'serious adverse event report', 'confidence': 0.386}
Completeness: 1.0
Severity: serious adverse event

Summary (Original)

Ofatumumab was granted regular approval in 2007, on the basis of superior PFS. Patients in the Hx-CD20-406 trial frequently had a history of severe infections. Infusional toxicity was common, manifesting as fever, dyspnea, and rash despite premedication. Myocardial infarction or angina was noted in four patients within two days of a dose of ofatumumAB. FDA recommended that monthly disease status assessments for progression occur during the first year for all patients enrolled into OMB110911. In a population with unmet need, an observed response rate where the lower bound of the 95% CI for the ORR was at least 25% would be of interest. Some of the regimens in Table 3 are based on randomized phase 3 trials that have not been submitted to FDA in support of an NDA or BLA. Inclusion of these regimens does not indicate that they are approved or recommended by FDA. The applicant's analyses of OS in patients alive at week 12 are not informative and potentially misleading. Because these analyses a a patient's OS a week 12 is not informative, these analyses could be misleading. Some studies were submitted as synopses or interim reports. Only studies 402 and 406 are included in the efficacy and primary safety analyses. The NCIWG criteria (Cheson 1996) were used to define response in study 406. Responders were defined as those with a complete response (CR) or nodular partial response (nPR) Non-responders were those with stable disease (SD) or progressive disease (PD)

Summary (Anonymized)

Ofatumumab is only being considered for CLL patients without adequate available therapy. Post-marketing studies are underway to further characterize the clinical benefit(s) of ofatumumbe (see section 1.4). This reviewer believes that based on the approval of other drugs for the treatment of CLL (and cancer in general), that ofatumomab has a tolerable risk-benefit profile for the intended population. The primary endpoint of study 406 was [NAME] Committee (IRC)-assessed ORR at 24 weeks. A total of 69 patients withdrew during the treatment period. 85 patients completed treatment and were withdrawn during follow-up. Infections (including infectious deaths) occurred frequent. There were some inconsistencies applied to multiple occurrences of tingling in areas of the face. The ORRs across the body weight subgroups (either dichotomized or trichotomized) were fairly consistent. The trend in the DR group of decreasing DOR for increasing body weight was not seen in the dichotomous group. The study Hx-CD20-406 was the only study to evaluate ofatumumab at the doses and schedule for which the applicant is seeking approval. It was the first study to assess the safety and effectiveness of the drug at the dose and schedule it was being evaluated for. It is the first to evaluate the drug's safety and efficacy at a higher dose and at a lower dose. The drug was not approved when the double refractory (refractory to fludarabine and alemtuzumab) CLL patient population was deemed by FDA as having a

Summary Metrics

Compression Ratio: 0.004
Reduction: 99.6%

GLOBAL ANALYSIS

Duplicate: True
Similarity Score: 1.0

Confidence: High

Comparison Metrics

Change Ratio: 0.045

Number of Changes: 217