

LinLog und LinDisCats

Max Wiegand

March 5, 2024

1 Introduction

what am I even writing about?

2 LinLog Preliminaries

2.1 Motivation/Intuition

feels kinda rambling... combine intuition with syntax definition below

Classical (and intuitionistic) logic deals with the propagation of stable truth values. If one has a true sentence A and an implication $A \Rightarrow B$, then B follows while A remains true. However, real-life implications are often causal and modify their premises. They cannot therefore be iterated arbitrarily. For example if A describes the ownership of 1€ and B owning a chocolate bar, an implication $A \multimap B$ (to be formally introduced later) would describe the process of buying such a chocolate bar for 1€, losing the 1€ in the process.

While such dealings with resources can of course be modeled in classical logic, it is easier done in the resource-sensitive *linear logic*, first described by Girard in 1987 [Gir87].

Here, we have two conjunctions simultaneously \otimes ("times" or "tensor") and $\&$ ("with"), which describe the availability of resources:

Suppose C is the ownership of a cookie and it costs also 1€ (i.e. we have $A \multimap C$). Then $B \otimes C$ states that one owns both a chocolate bar and a cookie. The implication $A \multimap B \otimes C$ is not possible, as it would mean that you are buying both, cookie and chocolate bar at the same time, for just 1€ total. However, from $A \multimap B$ and $A \multimap C$ we get $A \otimes A \multimap B \otimes C$, i.e. the process of buying both for 2€.

On the other hand, $B \& C$ states that one has a choice between either B or C (imagine a token). From the implications $A \multimap B$ and $A \multimap C$ we get the implication $A \multimap B \& C$, i.e. the process of buying a token to be exchanged for a chocolate bar or a cookie at a later time (with the choice lying with oneself). While this may seem like a disjunction, both implications $B \& C \multimap B$ and $B \& C \multimap C$ (exchanging the token for either product) are provable from $B \& C$, although not simultaneously.

Dually, we have two disjunctions \wp ("par") and \oplus ("plus"):

Suppose now that B and C are the ownership of a figurine of Pikachu or Mew respectively. Then $B \oplus C$ may be the ownership of a Kinder Egg containing either figurine. This means when buying that egg ($A \multimap B \oplus C$) we do not know which one we will get.

Our second disjunction, dual to \otimes , can be understood by linear implication and the linear negation (denoted as $(\cdot)^\perp$): Under the interpretation of ownership the linear negation is no interpreted as the absence of ownership but as negative ownership, i.e. debt. That means the negation of owning 1€, A , is owing someone 1€, A^\perp . With the par operator we can now write the linear implication $A \multimap B$ symmetrically as $A^\perp \wp B$.

(explicit interpretation?! shared pool between people containing Pika and Mew -> you have to get rid of one to use the other?)

In order to regain our stable truths known from classical logic, we need to employ two unitary connectives ! ("of course" or "bang") and ? ("why not"). The bang operator informs us that there is an infinite amount of a resource: The statement $!A$ translates into the ownership of an amount of money that is large enough for us to ignore resource sensitivity. Imagine for example a billionaire buying a Pokemon figurine instead of a social media site: His amount of money will not be noticeably smaller after buying the figurine. We can informally say $!A = (1 \& A) \otimes (1 \& A) \otimes \dots$ and therefore view classical logic as some sort of limit of linear logic, just as classical mechanics is a limit of quantum mechanics and the theory of relativity.

interpretation of ?wn.

interpretation of constants

2.2 Syntax

Quellen, Quellen, Quellen!!!

section needs serious reformatting...

main source: IntroLinLog

definition: "proof"; "cut elimination"

MAYBE to be added: \forall and \exists. Do we want predicate logic?

Definition 2.1 (Formula) *Formulas are defined inductively from atomic formulas and four constants 1 , \perp , \top and 0 :*

- *Every constant is a formula.*
- *Every atomic formula is a formula.*
- *If A is a formula, so are A^\perp , $!A$ and $?A$.*
- *If A and B are formula, so are $A \otimes B$, $A \wp B$, $A \& B$ and $A \oplus B$.*

Let Γ, Δ etc. be arbitrary, finite sequents of formulas (e.g.: $\Gamma = p_1, \dots, p_n$) and A and B formulas.

As we will later consider fragments without negation, we shall define linear logic with a two-sided calculus.

Definition 2.2 (Proof (tree)) A proof tree (often just proof) is a rooted tree with sequents of the form $\Gamma \vdash \Delta$ as edges and the following sequent rules as vertices (except for the root).

upper
vertex of
identity?

Structural rules We only have the exchange rule as a structural, missing the (general) weakening and contraction rules known from classical logic:

$$\frac{\Gamma, A, B, \Gamma' \vdash \Delta}{\Gamma, B, A, \Gamma' \vdash \Delta} \text{ex.L} \quad \frac{\Gamma \vdash \Delta, A, B, \Delta'}{\Gamma \vdash \Delta, B, A, \Delta'} \text{ex.R}$$

Identity rules We have the identity and negation rules:

$$\frac{}{A \vdash A} \text{id} \quad \frac{\Gamma_1 \vdash \Delta_1, A, \Delta'_1 \quad \Gamma_2, A, \Gamma'_2 \vdash \Delta_2}{\Gamma_2, \Gamma_1, \Gamma'_2 \vdash \Delta_1, \Delta_2, \Delta'_1} \text{Cut}$$

$$\frac{\Gamma \vdash A, \Delta}{\Gamma, A^\perp \vdash \Delta} \text{neg.L} \quad \frac{\Gamma, A \vdash \Delta}{\Gamma \vdash A^\perp, \Delta} \text{neg.R}$$

move negation to other place?

permanente
Lösung
für empty
axiom
finden

As already mentioned, the classical conjunction \wedge and disjunction \vee as well as their respective units split into two respectively. These can be classified as multiplicative and additive connectives.

Multiplicatives The calculus rules for the *multiplicative* conjunction \otimes , disjunction \wp ("par") and their units 1 and \perp ("bottom") are as follows:

$$\frac{\Gamma, A, B, \Gamma' \vdash \Delta}{\Gamma, A \otimes B, \Gamma' \vdash \Delta} \otimes_L \quad \frac{\Gamma \vdash \Delta, A \quad \Gamma' \vdash B, \Delta'}{\Gamma, \Gamma' \vdash \Delta, A \otimes B, \Delta'} \otimes_R$$

$$\frac{\Gamma, A \vdash \Delta \quad B, \Gamma' \vdash \Delta'}{\Gamma, A \wp B, \Gamma' \vdash \Delta, \Delta'} \wp_L \quad \frac{\Gamma \vdash \Delta, A, B, \Delta'}{\Gamma \vdash \Delta, A \wp B, \Delta'} \wp_R$$

$$\frac{\Gamma \vdash \Delta}{\Gamma, 1 \vdash \Delta} 1_L \quad \frac{}{\vdash 1} 1_R$$

$$\frac{}{\perp \vdash} \perp_L \quad \frac{\Gamma \vdash \Delta}{\Gamma \vdash \perp, \Delta} \perp_R$$

Remark 2.3 The rules \otimes_L and \wp_R imply, that the commas are to be read as \otimes on the left-hand side and as \wp on the the right-hand side. That means $A, B \vdash C, D$ is provable iff $A \otimes B \vdash C \wp D$ is provable.

PROOF: We only have to show that $A, B \vdash C, D$ follows from $A \otimes B \vdash C \wp D$ as the other direction is just our introduction rule:

$$\frac{\frac{\frac{}{A \vdash A} \text{id} \quad \frac{}{B \vdash B} \text{id}}{A, B \vdash A \otimes B} \otimes_R \quad \frac{A \otimes B \vdash C \wp D}{A, B \vdash C \wp D} \text{Cut} \quad \frac{\frac{}{C \vdash C} \quad \frac{}{D \vdash D}}{C \wp D \vdash C, D} \wp_L}{A, B \vdash C, D} \quad \square$$

proofsymbol
:/

Additives The calculus rules for the *additive* conjunction $\&$ and disjunction \oplus ("plus") are as follows:

$$\begin{array}{c} \frac{\Gamma, A \vdash \Delta}{\Gamma, A \& B \vdash \Delta} \&_{L1} \quad \frac{\Gamma, B \vdash \Delta}{\Gamma, A \& B \vdash \Delta} \&_{L2} \\ \frac{\Gamma \vdash \Delta, A \quad \Gamma \vdash \Delta, B}{\Gamma \vdash \Delta, A \& B} \&_R \\ \\ \frac{\Gamma, A \vdash \Delta \quad \Gamma, B \vdash \Delta}{\Gamma, A \oplus B \vdash \Delta} \oplus_L \\ \frac{\Gamma \vdash A, \Delta}{\Gamma \vdash A \oplus B, \Delta} \oplus_{R1} \quad \frac{\Gamma \vdash B, \Delta}{\Gamma \vdash A \oplus B, \Delta} \oplus_{R2} \\ \\ \frac{}{\Gamma, 0 \vdash \Delta} 0_L \quad \frac{}{\Gamma \vdash \top, \Delta} \top_R \end{array}$$

Notice the difference between $\&_R$ and \otimes_R (and dually between \oplus_L and \wp_L): while for \otimes_R the contexts Γ etc. are arbitrary and get combined in the conclusion, $\&_R$ requires the contexts to be equal. In classical logic these rules can be shown to be equivalent using its additional structural rules.

Furthermore, note that the unit introduction rules also introduce arbitrary contexts, and that each unit only has one introduction rule as opposed to the multiplicative units.

Remark 2.4 Similarly to the multiplicative connectives above, the additive connectives have a invertibility statement: $\Gamma \vdash A \& B$ is provable iff $\Gamma \vdash A$ and $\Gamma \vdash B$ are provable. Dually, $A \oplus B \vdash \Delta$ iff $A \vdash \Delta$ and $B \vdash \Delta$.

remark
unnötig?

PROOF: As with the multiplicative statement, one direction suffices:

$$\frac{\Gamma \vdash A \& B \quad \frac{}{A \vdash A}}{A \& B \vdash A} \quad \square$$

$\Gamma \vdash B$ follows the same way.

Definition 2.5 We call two formulas A and B (linearly) equivalent iff $A \vdash B$ and $B \vdash A$ are provable, and write $A \equiv B$.

Remark 2.6 The names "multiplicative" and "additive" are motivated by the following relations:

$$\begin{aligned} A \otimes (B \oplus C) &\equiv (A \otimes B) \oplus (A \otimes C) \\ A \wp (B \& C) &\equiv (A \wp B) \& (A \wp C) \end{aligned}$$

The similarities to basic arithmetic don't end there:

$$A \otimes 0 \equiv 0, \quad A \wp \top \equiv \top$$

PROOF: We only show $A \otimes (B \oplus C) \equiv (A \otimes B) \oplus (A \otimes C)$.

- $A \otimes (B \oplus C) \vdash (A \otimes B) \oplus (A \otimes C)$:

$$\frac{\frac{\frac{A \vdash A}{A, B \vdash A \otimes B} \quad \frac{B \vdash B}{A, B \vdash A \otimes B}}{A, B \vdash (A \otimes B) \oplus (A \otimes C)} \quad \frac{\frac{A \vdash A}{A, C \vdash A \otimes C} \quad \frac{C \vdash C}{A, C \vdash A \otimes C}}{A, C \vdash (A \otimes B) \oplus (A \otimes C)} \\ \frac{A, B \oplus C \vdash (A \otimes B) \oplus (A \otimes C)}{A \otimes (B \oplus C) \vdash (A \otimes B) \oplus (A \otimes C)}$$

- $(A \otimes B) \oplus (A \otimes C) \vdash A \otimes (B \oplus C)$:

$$\frac{\frac{\frac{A \vdash A}{A, B \vdash A \otimes (B \oplus C)} \quad \frac{B \vdash B}{B \vdash B \oplus C}}{A \otimes B \vdash A \otimes (B \oplus C)} \quad \frac{\frac{A \vdash A}{A, C \vdash A \otimes (B \oplus C)} \quad \frac{C \vdash C}{C \vdash B \oplus C}}{A \otimes C \vdash A \otimes (B \oplus C)} \\ \frac{A \otimes B \vdash A \otimes (B \oplus C) \quad A \otimes C \vdash A \otimes (B \oplus C)}{(A \otimes B) \oplus (A \otimes C) \vdash A \otimes (B \oplus C)}$$

The proof for $A \wp (B \& C) \equiv (A \wp B) \& (A \wp C)$ is similar, the proof for the equations of the constants a trivial application of their introduction rules. \square

Theorem 2.7 We have the following equivalencies for the linear negation:

- For the constants:

$$\begin{aligned} 1^\perp &\equiv \perp & \perp^\perp &\equiv 1 \\ \top^\perp &\equiv 0 & 0^\perp &\equiv \top \end{aligned}$$

- The negation is involutory: $(A^\perp)^\perp \equiv A$

- The De Morgan equations hold:

$$\begin{aligned} (A \otimes B)^\perp &\equiv A^\perp \wp B^\perp & (A \wp B)^\perp &\equiv A^\perp \otimes B^\perp \\ (A \& B)^\perp &\equiv A^\perp \oplus B^\perp & (A \oplus B)^\perp &\equiv A^\perp \& B^\perp \end{aligned}$$

PROOF: We shall only prove parts as the rest is trivial.

- $1^\perp \vdash \perp$:

$$\frac{\frac{\frac{}{\vdash 1} 1_R}{\vdash 1, \perp} \perp_R}{1^\perp \vdash \perp} \text{neg.L}$$

- $(A^\perp)^\perp \vdash A$:

$$\frac{\frac{\overline{A \vdash A}}{\vdash A^\perp, A}}{(A^\perp)^\perp \vdash A}$$

- $(A \otimes B)^\perp \vdash A^\perp \wp B^\perp$:

$$\frac{\frac{\frac{\overline{A \vdash A}}{\vdash A^\perp, A} \quad \frac{\overline{B \vdash B}}{\vdash B^\perp, B}}{\vdash A \otimes B, A^\perp, B^\perp}}{(A \otimes B)^\perp \vdash A^\perp, B^\perp} \quad \square$$

$$(A \otimes B)^\perp \vdash A^\perp \wp B^\perp$$

proofsymbol

With the negation our calculus rules become therefore quite redundant and we can restrict them on the rules for the right-hand side by translating any two-sided sequent $A_1, \dots, A_n \vdash B_1, \dots, B_n$ into a one-sided sequent $\vdash A_1^\perp, \dots, A_n^\perp, B_1, \dots, B_n$. However, these redundancies are necessary when dealing with a negation-free fragment of LL as we will later do.

Definition 2.8 (Syntactical implication and equivalence) *We define linear implication with the par-operator:*

$$A \multimap B := A^\perp \wp B$$

We further define (syntactical) linear equivalence:

$$A \circ\!\!\multimap B := (A \multimap B) \& (B \multimap A)$$

It is easily seen that $\vdash A \multimap B$ iff $A \vdash B$ and that $\vdash A \circ\!\!\multimap B$ iff $A \equiv B$.

Exponentials The modality connectives reintroduce stable truths and with them the the weakening and contraction rules known from classical logic:

$$\begin{array}{ll} \frac{\Gamma \vdash \Delta}{\Gamma, !A \vdash \Delta} !_W \text{ (weakening)} & \frac{\Gamma, A \vdash \Delta}{\Gamma, !A \vdash \Delta} !_D \text{ (dereliction)} \\ \frac{\Gamma, !A, !A \vdash \Delta}{\Gamma, !A \vdash \Delta} !_C \text{ (contraction)} & \frac{! \Gamma \vdash ? \Delta, A}{! \Gamma \vdash ? \Delta, !A} !_R \\ \frac{\Gamma \vdash \Delta}{\Gamma \vdash \Delta, ?A} ?_W \text{ (weakening)} & \frac{\Gamma \vdash \Delta, A}{\Gamma \vdash \Delta, ?A} ?_D \text{ (dereliction)} \\ \frac{\Gamma \vdash \Delta, ?A, ?A}{\Gamma \vdash \Delta, ?A} ?_C \text{ (contraction)} & \frac{! \Gamma, A \vdash ? \Delta}{! \Gamma, ?A \vdash ? \Delta} ?_L \end{array}$$

$$(!A)^\perp \equiv ?(A^\perp) \quad (?A)^\perp \equiv !(A^\perp)$$

Here, the context $! \Gamma$ is given by applying the $!$ -modality to every formula of Γ , i.e. $! \Gamma = !q, !p, \dots$ for $\Gamma = q, p, \dots$; same for $? \Delta$.

Remark 2.9 These modalities are called exponentials because of the following relation:

$$!(A \& B) \equiv !A \otimes !B, \quad ?(A \oplus B) \equiv ?A \wp ?B$$

PROOF: We will only prove the second equivalence. The first is acquired by duality.

- $?A \wp ?B \vdash ?(A \oplus B)$:

$$\frac{\frac{\frac{\overline{A \vdash A}}{A \vdash A \oplus B} \oplus_R}{A \vdash ?(A \oplus B)} ?_D \quad \frac{\frac{\frac{\overline{B \vdash B}}{B \vdash A \oplus B} \oplus_R}{B \vdash, ?(A \oplus B)} ?_D}{?B \vdash ?(A \oplus B)} ?_L}{(?A) \wp (?B) \vdash ?(A \oplus B), ?(A \oplus B)} \wp}{?A \wp ?B \vdash ?(A \oplus B)} ?_D$$

- $?(A \oplus B) \vdash ?A \wp ?B$:

$$\frac{\frac{\frac{\overline{A \vdash A}}{A \vdash ?A} ?_D}{A \vdash ?A, ?B} ?_W \quad \frac{\frac{\frac{\overline{B \vdash B}}{B \vdash ?B}}{B \vdash ?B, ?A} ?_W}{A \oplus B \vdash ?A, ?B} \oplus_L}{?(A \oplus B) \vdash ?A, ?B} ?_L}{?(A \oplus B) \vdash ?A \wp ?B} \wp_R$$

□

proofsymbol
なんで?:(

2.3 Cut elimination

add "spoiler" to categoru theoru

As is the case with most, if not all, logical system, the Cut rule of our calculus is admissible. That means for any proof π of a sequent $\Gamma \vdash \Delta$ we can find a *cut-free* proof π' of that same sequent. We will prove this for the multiplicative fragment by simplifying the proof due to Braüner [Bra96, Appendix B]. Baüner's proof includes the exponentials which require additional care because of the contraction rule. The general strategy consists of either, moving a cut upwards in a proof or replacing it with cuts of simpler formulas. To do this, we need two definitions:

Definition 2.10 (Degree) The degree $\partial(A)$ of a formula A is defined inductively:

- $\partial(A) = 1$, for A atomic or constant
- $\partial(A^\perp) = \partial(A)$
- $\partial(A \otimes B) = \partial(A \wp B) = \max\{\partial(A), \partial(B)\} + 1$
- $\partial(!A) = \partial(?A) = \partial(A) + 1$

The degree of a cut is the degree of the cut formula. The cut formula is called principal formula. The degree $\partial(\pi)$ of a proof π is the supremum of the degrees of the cuts in the proof.

Definition 2.11 (Height) The height $h(\pi)$ of a proof π is just its height as a rooted tree:

- $h(\pi) = 1$ when π is an instance of a rule with zero premises, i.e. an axiom.

- $h(\pi) = h(\tau) + 1$, for $\pi = \frac{\tau}{\Gamma \vdash \Delta}$
- $h(\pi) = \max\{h(\tau), h(\tau')\} + 1$, for $\pi = \frac{\tau}{\Gamma \vdash \Delta} \frac{\tau'}{\Gamma \vdash \Delta}$

With these, we can now eliminate cuts in a proof:

Lemma 2.12 *Let τ be a proof consisting of two subproofs τ_1 and τ_2 of strictly lower degree, i.e.:*

$$\tau = \frac{\frac{\tau_1}{\vdots} \quad \frac{\tau_2}{\vdots}}{\frac{\Gamma_1 \vdash \Delta_1, A, \Delta'_1 \quad \Gamma_2, A, \Gamma'_2 \vdash \Delta_2}{\Gamma_2, \Gamma_1, \Gamma'_2 \vdash \Delta_1, \Delta_2, \Delta'_1}}$$

Then, we can find a proof τ' of that same sequent $\Gamma_2, \Gamma_1, \Gamma'_2 \vdash \Delta_1, \Delta_2, \Delta'_1$ with $\partial(\tau') < \partial(\tau)$.

tabularx format

PROOF: Induction on $h(\tau_1) + h(\tau_2)$. The immediate subproofs of τ_i are denoted by π_j . We perform transformations case by case:

- If one of the subproofs is an axiom, we just take the remaining one, resulting in a proof with lower degree:

$$\begin{array}{ccc} \frac{\frac{A \vdash A}{\vdots} \quad \frac{\tau_2}{\vdots}}{A, \Gamma \vdash \Delta} & \rightsquigarrow & \frac{\tau_2}{\vdots} \\ \frac{\tau_1}{\vdots} & & A, \Gamma \vdash \Delta \\ \frac{\Gamma \vdash \Delta, A \quad A \vdash A}{\Gamma \vdash \Delta, A} & \rightsquigarrow & \frac{\tau_1}{\vdots} \\ & & \Gamma \vdash \Delta, A \end{array}$$

- τ_1 does not introduce the principal formula in its last rule. Depending on the subcase, we perform one of the following transformations, pushing τ_2 upwards, and use the induction hypothesis on the resulting subproof.

Problem: Resulting proofs for the \mathfrak{A}_L -cases are illegal under current formulation of the \mathfrak{A}_L -rule. (Also: they're kinda the same?) Possible solutions:

- 1: add exchanges to transformation (easy; ugly)
- 2: find generalized \mathfrak{A}_L -rule (more difficult; possibly uglier)

To be added: \otimes_R -cases.

Maybe combine \mathfrak{A}_L - and \otimes_R -cases into one? (Requires generalized formulation of the rules)

$$\begin{array}{c}
\frac{\frac{\pi_1}{\Gamma_1, A \vdash \Delta_1, C, \Delta'_1} \quad \frac{\pi_2}{B, \Gamma_2 \vdash \Delta_2} \mathfrak{A}_L \quad \frac{\tau_2}{\Gamma_3, C, \Gamma'_3 \vdash \Delta_3}}{\Gamma_1, A \mathfrak{A} B, \Gamma_2 \vdash \Delta_1, C, \Delta'_1, \Delta_2} \mathfrak{A}_L \quad \frac{\tau_2}{\Gamma_3, C, \Gamma'_3 \vdash \Delta_3} \rightsquigarrow \frac{\frac{\pi_1}{\Gamma_1, A \vdash \Delta_1, C, \Delta'_1} \quad \frac{\tau_2}{\Gamma_3, C, \Gamma'_3 \vdash \Delta_3}}{\Gamma_3, \Gamma_1, A, \Gamma'_3 \vdash \Delta_1, \Delta_3, \Delta'_1} \quad \frac{\pi_2}{B, \Gamma_2 \vdash \Delta_2} \\
\frac{\pi_1}{\Gamma_1, A \vdash \Delta_1} \quad \frac{\pi_2}{B, \Gamma_2 \vdash \Delta_2, C, \Delta'_2} \mathfrak{A}_L \quad \frac{\tau_2}{\Gamma_3, C, \Gamma'_3 \vdash \Delta_3} \rightsquigarrow \frac{\frac{\pi_1}{\Gamma_1, A \vdash \Delta_1} \quad \frac{\pi_2}{B, \Gamma_2 \vdash \Delta_2, C, \Delta'_2} \quad \frac{\tau_2}{\Gamma_3, C, \Gamma'_3 \vdash \Delta_3}}{\Gamma_3, \Gamma_1, A \mathfrak{A} B, \Gamma_2, \Gamma'_3 \vdash \Delta_1, \Delta_2, \Delta_3, \Delta'_2} \\
\frac{\pi}{\Psi_1 \vdash \Psi_2, A, \Psi'_2} \quad \frac{\tau_2}{\Gamma, A, \Gamma' \vdash \Delta} \rightsquigarrow \frac{\frac{\pi}{\Psi_1 \vdash \Psi_2, A, \Psi'_2} \quad \frac{\tau_2}{\Gamma, A, \Gamma' \vdash \Delta}}{\Gamma, \Psi_1, \Gamma' \vdash \Psi_2, \Delta, \Psi'_2} r \\
\frac{\Psi_1 \vdash \Psi_2, A, \Psi'_2}{\Phi_1 \vdash \Phi_2, A, \Phi'_2} r \quad \frac{\tau_2}{\Gamma, A, \Gamma' \vdash \Delta} \rightsquigarrow \frac{\frac{\Psi_1 \vdash \Psi_2, A, \Psi'_2}{\Phi_1 \vdash \Phi_2, A, \Phi'_2} r \quad \frac{\tau_2}{\Gamma, A, \Gamma' \vdash \Delta}}{\Gamma, \Phi_1, \Gamma' \vdash \Phi_2, \Delta, \Phi'_2} r
\end{array}$$

Here, r is one of the following rules: $\otimes_L, \mathfrak{A}_R, 1_L, \perp_R, \text{ex.R}$ or ex.L . Note that the exchange rule might be performed on A and a neighboring formula B , thus possibly requiring multiple applications of the exchange rule in the transformed proof. This is denoted by the double line.

- τ_2 does not introduce the principal formula in its last rule. Dually to the case above, we perform one of the following transformations, pushing τ_1 upwards, and use the induction hypothesis on the resulting subproof.

To be added: \mathfrak{A}_L - and \otimes_R -cases (same problems as above)

$$\frac{\frac{\tau_1}{\vdots} \quad \frac{\pi}{\Psi_1, A, \Psi'_1 \vdash \Psi_2} \quad \frac{\Phi_1, A, \Phi'_1 \vdash \Phi_2}{\Phi_1, \Gamma, \Phi'_1 \vdash \Delta, \Phi_2, \Delta'} r}{\Gamma \vdash \Delta, A, \Delta'} \rightsquigarrow \frac{\frac{\tau_1}{\vdots} \quad \frac{\pi}{\Psi_1, A, \Psi'_1 \vdash \Psi_2}}{\Gamma \vdash \Delta, A, \Delta'} \quad \frac{\Phi_1, A, \Phi'_1 \vdash \Phi_2}{\Phi_1, \Gamma, \Phi'_1 \vdash \Delta, \Phi_2, \Delta'} r$$

Again, r is one of the following rules: $\otimes_L, \mathfrak{A}_R, 1_L, \perp_R, \text{ex.R}$ or ex.L .

- Finally, if both τ_1 and τ_2 introduce the principle formula with their last rule, we perform one the following transformations resulting in a proof of lower degree:

$$\begin{array}{c}
\frac{\frac{\pi_1}{\vdots} \quad \frac{\pi_2}{\vdots} \quad \frac{\pi_3}{\vdots}}{\Gamma_1 \vdash \Delta_1, A \quad \Gamma_2 \vdash B, \Delta_2 \quad \Gamma_3, A, B, \Gamma'_3 \vdash \Delta_3} \rightsquigarrow \frac{\frac{\pi_1}{\vdots} \quad \frac{\pi_2}{\vdots} \quad \frac{\pi_3}{\vdots}}{\Gamma_1 \vdash \Delta_1, A \quad \Gamma_2 \vdash B, \Delta_2 \quad A, B, \Gamma_3 \vdash \Delta_3} \\
\frac{\Gamma_1, \Gamma_2 \vdash \Delta_1, A \otimes B, \Delta_2}{\Gamma_3, \Gamma_1, \Gamma_2, \Gamma'_3 \vdash \Delta_1, \Delta_3, \Delta_2} \rightsquigarrow \frac{\Gamma_1, \Gamma_2, \Gamma_3 \vdash \Delta_1, \Delta_3, \Delta_2}{\Gamma_3, \Gamma_1, \Gamma_2, \Gamma'_3 \vdash \Delta_1, \Delta_3, \Delta_2} \\
\frac{\frac{\pi_1}{\vdots} \quad \frac{\pi_2}{\vdots} \quad \frac{\pi_3}{\vdots}}{\Gamma_1 \vdash \Delta_1, A, B, \Delta'_1 \quad \Gamma_2, A \vdash \Delta_2 \quad B, \Gamma_3 \vdash \Delta_3} \rightsquigarrow \frac{\frac{\pi_1}{\vdots} \quad \frac{\pi_2}{\vdots} \quad \frac{\pi_3}{\vdots}}{\Gamma_1 \vdash \Delta_1, A, B, \Delta'_1 \quad \Gamma_2, A \vdash \Delta_2 \quad B, \Gamma_3 \vdash \Delta_3} \\
\frac{\Gamma_1 \vdash \Delta_1, A \mathfrak{A} B, \Delta'_1}{\Gamma_2, \Gamma_1, \Gamma_3 \vdash \Delta_1, \Delta_2, \Delta_3, \Delta'_1} \rightsquigarrow \frac{\Gamma_1 \vdash \Delta_1, A, B, \Delta'_1 \quad \Gamma_2, A \vdash \Delta_2 \quad B, \Gamma_3 \vdash \Delta_3}{\Gamma_1, \Gamma_2, \Gamma_3 \vdash \Delta_1, \Delta_2, \Delta_3, \Delta'_1} \\
\frac{\pi}{\vdots} \rightsquigarrow \frac{\pi}{\vdots} \\
\frac{\vdash 1}{\Gamma \vdash \Delta} \rightsquigarrow \frac{\pi}{\vdots} \\
\frac{\vdash 1}{\Gamma \vdash \Delta} \rightsquigarrow \frac{\pi}{\vdots} \\
\frac{\vdash 1}{\Gamma \vdash \Delta} \rightsquigarrow \frac{\pi}{\vdots}
\end{array}$$

testing for proofsymbol

proofsymbol

□

Lemma 2.13

2.3.1 Commuting cuts

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma_1 \vdash \Delta_1, A, \Delta'_1} \quad \frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_2, A, \Gamma'_2 \vdash \Delta_2, B, \Delta'_2} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, B, \Gamma'_3 \vdash \Delta_3}}{\Gamma_3, \Gamma_2, A, \Gamma'_2 \vdash \Delta_2, \Delta_3, \Delta'_2} \text{Cut}}{\Gamma_3, \Gamma_2, \Gamma_1, \Gamma'_2, \Gamma'_3 \vdash \Delta_1, \Delta_2, \Delta_3, \Delta'_2, \Delta'_1} \text{Cut} \quad \longleftrightarrow \quad \frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma_1 \vdash \Delta_1, A, \Delta'_1} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_2, A, \Gamma'_2 \vdash \Delta_2, B, \Delta'_2}}{\Gamma_2, \Gamma_1, \Gamma'_2 \vdash \Delta_2, B, \Delta'_2} \text{Cut} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, B, \Gamma'_3 \vdash \Delta_3}}{\Gamma_3, \Gamma_2, \Gamma_1, \Gamma'_2, \Gamma'_3 \vdash \Delta_1, \Delta_2, \Delta_3, \Delta'_2, \Delta'_1} \text{Cut}$$

put that stuff back into a normal center instead of tabularx?

The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma_1 \vdash \Delta_1, A, \Delta'_1} \quad \frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_2 \vdash \Delta_2, B, \Delta'_2} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, A, B, \Gamma'_3 \vdash \Delta_3}}{\Gamma_3, A, \Gamma_2, \Gamma'_3 \vdash \Delta_2, \Delta_3, \Delta'_2} \text{Cut}}{\Gamma_3, \Gamma_1, \Gamma_2, \Gamma'_3 \vdash \Delta_1, \Delta_2, \Delta_3, \Delta'_2, \Delta'_1} \text{Cut}$$

is transformed into

$$\frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_2 \vdash \Delta_2, B, \Delta'_2} \quad \frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma_1 \vdash \Delta_1, A, \Delta'_1} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, A, B, \Gamma'_3 \vdash \Delta_3}}{\Gamma_3, \Gamma_1, B, \Gamma'_3 \vdash \Delta_1, \Delta_3, \Delta'_1} \text{Cut}}{\Gamma_3, \Gamma_1, \Gamma_2, \Gamma'_3 \vdash \Delta_2, \Delta_1, \Delta_3, \Delta'_1, \Delta'_2} \text{Cut}$$

and vice versa

think about the deltas...: aufteilen in zwei Regeln mit einseitigen Deltas für A und B oder allgemein halten und viele exchanges drunter klatschen?

2.3.2 expansion steps

why are they called η -expansion steps?

$$\begin{array}{ccc} \frac{}{A \otimes B \vdash A \otimes B} \text{id} & \rightsquigarrow & \frac{\frac{}{A \vdash A} \text{id} \quad \frac{}{B \vdash B} \text{id}}{\frac{A, B \vdash A \otimes B}{A \otimes B \vdash A \otimes B} \otimes_L} \otimes_R \\[10pt] \frac{}{A \wp B \vdash A \wp B} \text{id} & \rightsquigarrow & \frac{\frac{}{A \vdash A} \text{id} \quad \frac{}{B \vdash B} \text{id}}{\frac{A \wp B \vdash A, B}{A \wp B \vdash A \wp B} \wp_R} \wp_L \\[10pt] \frac{}{1 \vdash 1} \text{id} & \rightsquigarrow & \frac{}{\frac{\vdash 1}{1 \vdash 1} 1_L} 1_R \\[10pt] \frac{}{\perp \vdash \perp} \text{id} & \rightsquigarrow & \frac{}{\frac{\perp \vdash}{\perp \vdash \perp} \perp_R} \perp_L \end{array}$$

format
(page
break)

2.3.3 The axiom steps

gone to 2.12

2.3.4 The exchange steps

that's just changing the contexts Γ_2 and Γ'_2

$$\begin{array}{ccc}
 \frac{\frac{\pi_1}{\vdots} \quad \frac{\frac{\pi_2}{\vdots} \quad \frac{\Gamma_2, A, B, \Gamma'_2 \vdash \Delta_2}{\Gamma_2, B, A, \Gamma'_2 \vdash \Delta_2}}{\Gamma_1 \vdash \Delta_1, A, \Delta'_1} \quad \frac{\Gamma_2, A, B, \Gamma'_2 \vdash \Delta_2}{\Gamma_2, B, A, \Gamma'_2 \vdash \Delta_2}}{\Gamma_2, B, \Gamma_1, \Gamma'_2 \vdash \Delta_1, \Delta_2, \Delta'_1} \text{ex.L} & \rightsquigarrow & \frac{\frac{\pi_1}{\vdots} \quad \frac{\Gamma_1 \vdash \Delta_1, A, \Delta'_1}{\Gamma_2, \Gamma_1, B, \Gamma'_2 \vdash \Delta_1, \Delta_2, \Delta'_1}}{\Gamma_2, B, \Gamma_1, \Gamma'_2 \vdash \Delta_1, \Delta_2, \Delta'_1} \text{ex.L} \\
 \\
 \frac{\frac{\pi_1}{\vdots} \quad \frac{\frac{\pi_2}{\vdots} \quad \frac{A, B, \Gamma_2 \vdash \Delta_2}{B, A, \Gamma_2 \vdash \Delta_2}}{\Gamma_1 \vdash \Delta_1, B} \quad \frac{A, B, \Gamma_2 \vdash \Delta_2}{B, A, \Gamma_2 \vdash \Delta_2}}{\Gamma_1, A, \Gamma_2 \vdash \Delta_1, \Delta_2} \text{ex.L} & \rightsquigarrow & \frac{\frac{\pi_1}{\vdots} \quad \frac{\Gamma_1 \vdash \Delta_1, B}{A, \Gamma_1, \Gamma_2 \vdash \Delta_1, \Delta_2}}{\Gamma_1, A, \Gamma_2 \vdash \Delta_1, \Delta_2} \text{ex.L}
 \end{array}$$

The double line signals multiple applications of the rule.

2.3.5 Principal formulae

gone to 2.12

2.3.6 Secondary conclusion

gone to 2.12

2.3.7 Secondary hypothesis

3 Categorical Preliminaries

3.1 Categories

sources for basic definitions. (nlab proly doesn't suffice)

Definition 3.1 (Category) A Category \mathfrak{C} consists of the following data:

- A class $\text{Obj}(\mathfrak{C})$ of objects.
- For every pair of objects $A, B \in \text{Obj}(\mathfrak{C})$ there is a class $\text{Hom}(A, B)$ of morphisms $f : A \rightarrow B$ from A to B .

- *Morphisms compose:* For $f \in \text{Hom}(A, B)$ and $g \in \text{Hom}(B, C)$ there is a morphism $g \circ f \in \text{Hom}(A, C)$. That composition is associative:

$$h \circ (g \circ f) = (h \circ g) \circ f$$

We will write gf for $g \circ f$ when appropriate.

- For every object A there is an identity morphism $\text{id}_A \in \text{Hom}(A, A)$:

$$f \circ \text{id}_A = f, \quad \text{id}_B \circ f = f$$

for $f \in \text{Hom}(A, B)$

If the hom-classes are sets, we call the category *locally small*. If the object class is a set, we call the category *small*. Otherwise, we call the category *large*.

If we have a category \mathfrak{C} , we call its opposite category $\mathfrak{C}^{\text{opp}}$ the category with the following structure:

- $\text{Obj}(\mathfrak{C}^{\text{opp}}) = \text{Obj}(\mathfrak{C})$
- $\forall A, B \in \text{Obj}(\mathfrak{C}) : \text{Hom}_{\mathfrak{C}^{\text{opp}}}(A, B) = \text{Hom}_{\mathfrak{C}}(B, A)$

extra definition for opposite structure?

Definition 3.2 (Functor) A (covariant) functor $F : \mathfrak{C} \rightarrow \mathfrak{D}$ between two categories \mathfrak{C} and \mathfrak{D} is a function mapping each object $A \in \text{Obj}(\mathfrak{C})$ to an object $F(A) \in \text{Obj}(\mathfrak{D})$ and each morphism $f \in \text{Hom}(A, B)$ to a morphism $F(f) \in \text{Hom}(F(A), F(B))$ such that identity and composition are preserved:

$$F(\text{id}_A) = \text{id}_{F(A)}, \quad F(g \circ f) = F(g) \circ F(f)$$

A functor $F : \mathfrak{C}^{\text{opp}} \rightarrow \mathfrak{D}$ is called *contravariant* on \mathfrak{C} .

We will drop the parentheses when appropriate.

Definition 3.3 (Natural transformation) A natural transformation $\tau : F \rightarrow G$ between two functors $F, G : \mathfrak{C} \rightarrow \mathfrak{D}$ is family of morphisms in \mathfrak{D} :

$$\tau = \{\tau_A : FA \rightarrow GA \mid A \in \text{Obj}(\mathfrak{C})\}$$

such that $\tau_B F(f) = G(f) \tau_A$ for all $f : A \rightarrow B \in \text{Morph}(\mathfrak{C})$, i.e. the following diagram commutes:

$$\begin{array}{ccc} FA & \xrightarrow{Ff} & FB \\ \downarrow \tau_A & & \downarrow \tau_B \\ GA & \xrightarrow{Gf} & GB \end{array}$$

If τ_A is an isomorphism for all $A \in \text{Obj}(\mathfrak{C})$, we call τ a *natural isomorphism*. We will often represent a natural transformation by a single one of its members and also drop the index when appropriate.

definition: equivalence of categories

Definition 3.4 (Adjunction) *Let \mathfrak{C} and \mathfrak{D} be categories. We call $F : \mathfrak{C} \rightarrow \mathfrak{D}$ the left adjoint of $G : \mathfrak{D} \rightarrow \mathfrak{C}$ and G the right adjoint of F if there is an isomorphism that is natural in $A \in \mathfrak{C}$ and $B \in \mathfrak{D}$:*

source:
Brandenburg

$$\text{Hom}_{\mathfrak{D}}(FA, B) \cong \text{Hom}_{\mathfrak{C}}(A, GB)$$

Equivalently, we call F the left adjoint of G if there are natural transformations

$$\begin{aligned} \eta : \text{id}_{\mathfrak{C}} &\rightarrow G \circ F \\ \varepsilon : F \circ G &\rightarrow \text{id}_{\mathfrak{D}} \end{aligned}$$

fulfilling the following condition:

$$\begin{aligned} \text{id}_{FA} &= \varepsilon(FA) \circ F\eta(A) \\ \text{id}_{GB} &= G\varepsilon(B) \circ \eta(GB) \end{aligned}$$

i.e. the following diagrams commute:

$$\begin{array}{ccc} & F \circ G \circ F & \\ F \bullet \eta \nearrow & & \searrow \varepsilon \bullet F \\ F \text{id}_{\mathfrak{C}} & \xlongequal{\text{id}} & \text{id}_{\mathfrak{C}} \circ F \end{array} \quad \begin{array}{ccc} & G \circ F \circ G & \\ \eta \bullet G \nearrow & & \searrow G \bullet \varepsilon \\ \text{id}_{\mathfrak{C}} \circ G & \xlongequal{\text{id}} & G \circ \text{id}_{\mathfrak{D}} \end{array}$$

maybe
index
Schreib-
weise?

cut triangle diagrams or define •

We write $F \dashv G$.

Definition 3.5 (Monoidal category) A monoidal category $(\mathfrak{C}, \otimes, \mathbb{1}, \alpha, \lambda, \rho)$ is a category \mathfrak{C} with the following additional data:

- a functor $\otimes : \mathfrak{C} \times \mathfrak{C} \rightarrow \mathfrak{C}$, called monoidal or tensor product;
- an object $\mathbb{1} \in \text{Obj}(\mathfrak{C})$, called unit object;
- a natural isomorphism $\alpha : A \otimes (B \otimes C) \rightarrow (A \otimes B) \otimes C$, called associator;
- a natural isomorphism $\lambda : \mathbb{1} \otimes A \rightarrow A$, called left unitor;
- a natural isomorphism $\rho : A \otimes \mathbb{1} \rightarrow A$, called right unitor;

such that the following diagrams commute:

- the pentagon diagram:

$$\begin{array}{ccccc} & & A \otimes (B \otimes (C \otimes D)) & & \\ & \swarrow \text{id} \otimes \alpha & & \searrow \alpha & \\ A \otimes ((B \otimes C) \otimes D) & & & & (A \otimes B) \otimes (C \otimes D) \\ \downarrow \alpha & & & & \downarrow \alpha \\ (A \otimes (B \otimes C)) \otimes D & \xrightarrow{\alpha \otimes \text{id}} & & & ((A \otimes B) \otimes C) \otimes D \end{array}$$

- the unit diagram:

$$\begin{array}{ccc}
 (A \otimes \mathbb{1}) \otimes B & \xrightarrow{\alpha} & A \otimes (\mathbb{1} \otimes B) \\
 \searrow \rho \otimes \text{id} & & \swarrow \text{id} \otimes \lambda \\
 & A \otimes B &
 \end{array}$$

A symmetric monoidal category has an additional natural isomorphism:

$$s_{A,B} : A \otimes B \rightarrow B \otimes A$$

satisfying the following conditions:

- the hexagon law:

$$\begin{array}{ccccc}
 A \otimes (B \otimes C) & \xrightarrow{\alpha} & (A \otimes B) \otimes C & \xrightarrow{s_{A \otimes B, C}} & C \otimes (A \otimes B) \\
 \downarrow \text{id} \otimes s_{B, C} & & & & \downarrow \alpha \\
 A \otimes (C \otimes B) & \xrightarrow{\alpha} & (A \otimes C) \otimes B & \xrightarrow{s_{A, C \otimes B}} & (C \otimes A) \otimes B
 \end{array}$$

- the inverse law: $s_{A,B}^{-1} = s_{B,A}$
- the unit law: $\lambda = \rho s$

$$\begin{array}{ccc}
 \mathbb{1} \otimes A & \xrightarrow{s_{\mathbb{1}, A}} & A \otimes \mathbb{1} \\
 \searrow \lambda & & \swarrow \rho \\
 & A &
 \end{array}$$

Diagramme verschönern

fix associators throughout the document (direction)

Definition 3.6 (Closed monoidal category) A monoidal category \mathfrak{C} is called right closed, resp. left closed, iff there is a functor $- \multimap - : \mathfrak{C}^{\text{opp}} \times \mathfrak{C} \rightarrow \mathfrak{C}$, resp. $- \multimap - : \mathfrak{C} \times \mathfrak{C}^{\text{opp}} \rightarrow \mathfrak{C}$, such that there is a natural isomorphism $\text{Hom}(A \otimes B, C) \cong \text{Hom}(B, A \multimap C)$, resp. $\text{Hom}(A \otimes B, C) \cong \text{Hom}(A, C \multimap B)$, i.e. iff the functor $- \otimes B$, resp. $A \otimes -$, has a right adjoint. A monoidal \mathfrak{C} is called biclosed iff it is both left and right closed.

lengthy,
ram-
bling...
muss el-
elegant
gehen

definitely mixed up the right and left homs there

If \mathfrak{C} is symmetric, these functors coincide and we call the category closed.

The functors $- \multimap - : \mathfrak{C}^{\text{opp}} \times \mathfrak{C} \rightarrow \mathfrak{C}$ and $- \multimap - : \mathfrak{C} \times \mathfrak{C}^{\text{opp}} \rightarrow \mathfrak{C}$ are called internal hom functors.

following definitions prollly unnötig:

Definition 3.7 (Enriched category) Let $(\mathfrak{M}, \mathbb{1}, \alpha, \lambda, \rho)$ be a monoidal category. An \mathfrak{M} -enriched category \mathfrak{C} consists of the following data:

- A class $\text{Obj}(\mathfrak{C})$ of objects;

- For every pair of objects $(A, B) \in \text{Obj}(\mathfrak{C}) \times \text{Obj}(\mathfrak{C})$ an object $\mathfrak{C}(A, B) \in \text{Obj}(\mathfrak{M})$ called hom object;
- For every triple of objects $(A, B, C) \in \text{Obj}(\mathfrak{C}) \times \text{Obj}(\mathfrak{C}) \times \text{Obj}(\mathfrak{C})$ a morphism $\text{Morph}(\mathfrak{M}) \ni \circ_{A,B,C} : \mathfrak{C}(B, C) \otimes \mathfrak{C}(A, B) \rightarrow \mathfrak{C}(A, C)$ called composition morphism;
- For every object $A \in \mathfrak{C}$ a morphism $\text{id}_A : \mathbb{1} \rightarrow \mathfrak{C}(A, A)$;

such that the following diagrams commute:

$$\begin{array}{ccc}
(\mathfrak{C}(C, D) \otimes \mathfrak{C}(B, C)) \otimes \mathfrak{C}(A, B) & \xrightarrow{\alpha} & \mathfrak{C}(C, D) \otimes (\mathfrak{C}(B, C) \otimes \mathfrak{C}(A, B)) \\
\downarrow \circ_{B,C,D} \otimes \text{id}_{\mathfrak{C}(A,B)} & & \downarrow \text{id}_{\mathfrak{C}(C,D)} \otimes \circ_{A,B,C} \\
\mathfrak{C}(D, B) \otimes \mathfrak{C}(A, B) & & \mathfrak{C}(C, D) \otimes \mathfrak{C}(A, C) \\
& \searrow \circ_{A,B,D} \quad \swarrow \circ_{A,C,D} & \\
& \mathfrak{C}(A, D) &
\end{array}$$

$$\begin{array}{ccccc}
\mathbb{1} \otimes \mathfrak{C}(A, B) & \xrightarrow{\lambda} & \mathfrak{C}(A, B) & \xleftarrow{\rho} & \mathfrak{C}(A, B) \otimes \mathbb{1} \\
\downarrow \text{id}_B \otimes \text{id}_{\mathfrak{C}(A,B)} & & \nearrow \circ_{A,B,B} & & \downarrow \text{id}_{\mathfrak{C}(A,B)} \otimes \text{id}_A \\
& & \mathfrak{C}(B, B) \otimes \mathfrak{C}(A, B) & & \mathfrak{C}(A, B) \otimes \mathfrak{C}(A, A) \\
& & \nwarrow \circ_{A,A,B} & & \nwarrow \circ_{A,A,B}
\end{array}$$

Definition 3.8 (Enriched functors) Let \mathfrak{C} and \mathfrak{D} be \mathfrak{M} -enriched categories. An enriched functor $F : \mathfrak{C} \rightarrow \mathfrak{D}$ consists of:

- a function $F_0 : \text{Obj}(\mathfrak{C}) \rightarrow \text{Obj}(\mathfrak{D})$ between the objects
- a family of morphisms in \mathfrak{M} : $F_{A,B} : \mathfrak{C}(A, B) \rightarrow \mathfrak{D}(F_0 A, F_0 B)$

such that the following diagrams commute:

$$\begin{array}{ccc}
\mathfrak{C}(B, C) \otimes \mathfrak{C}(A, B) & \xrightarrow{\circ} & \mathfrak{C}(A, C) \\
\downarrow F_{B,C} \otimes F_{A,B} & & \downarrow F_{A,C} \\
\mathfrak{D}(F_0 B, F_0 C) \otimes \mathfrak{D}(F_0 A, F_0 B) & \xrightarrow{\circ} & \mathfrak{D}(F_0 A, F_0 C)
\end{array}$$

$$\begin{array}{ccc}
& \mathbb{1} & \\
\text{id}_A \swarrow & & \searrow \text{id}_{F_0 A} \\
\mathfrak{C}(A, A) & \xrightarrow{F_{A,A}} & \mathfrak{D}(F_0 A, F_0 A)
\end{array}$$

3.2 LinDisCats

Definition 3.9 (Linearly distributive category) A linearly distributive category $(\mathfrak{C}, \otimes, 1, \mathfrak{A}, \perp)$ is a category \mathfrak{C} consisting of:

- a monoidal category $(\mathfrak{C}, \otimes, 1, \alpha_{\otimes}, \lambda_{\otimes}, \rho_{\otimes})$, with \otimes called "tensor";
- a monoidal category $(\mathfrak{C}, \mathfrak{A}, \perp, \alpha_{\mathfrak{A}}, \lambda_{\mathfrak{A}}, \rho_{\mathfrak{A}})$, with \mathfrak{A} called "par";

- two natural transformations called left and right linear distributors respectively:

$$\partial_L : A \otimes (B \wp C) \rightarrow (A \otimes B) \wp C$$

$$\partial_R : (A \wp B) \otimes C \rightarrow A \wp (B \otimes C)$$

satisfying the following coherence conditions:

- coherence between the distributors and unitors:

$$\begin{array}{ccc}
1 \otimes (A \wp B) & \xrightarrow{\partial_L} & (1 \otimes A) \wp B \\
\searrow \lambda_{\otimes} & & \swarrow \lambda_{\otimes} \wp \text{id} \\
& A \wp B & \\
(A \wp B) \otimes 1 & \xrightarrow{\partial_R} & A \wp (B \otimes 1) \\
\searrow \rho_{\otimes} & & \swarrow \text{id} \otimes \rho_{\otimes} \\
& A \wp B & \\
\\
A \otimes (B \wp \perp) & \xrightarrow{\partial_L} & (A \otimes B) \wp \perp \\
\searrow \text{id} \otimes \rho_{\wp} & & \swarrow \rho_{\wp} \\
& A \otimes B & \\
(\perp \wp A) \otimes B & \xrightarrow{\partial_R} & \perp \wp (A \otimes B) \\
\searrow \lambda_{\wp} \otimes \text{id} & & \swarrow \lambda_{\wp} \\
& A \otimes B &
\end{array}$$

- coherence between distributors and associators:

$$\begin{array}{ccc}
& (A \otimes B) \otimes (C \wp D) & \\
\swarrow \alpha_{\otimes} & & \searrow \partial_L \\
A \otimes (B \otimes (C \wp D)) & & ((A \otimes B) \otimes C) \wp D \\
\downarrow \text{id} \otimes \partial_L & & \downarrow \alpha_{\otimes} \wp \text{id} \\
A \otimes ((B \otimes C) \wp D) & \xrightarrow{\partial_L} & (A \otimes (B \otimes C)) \wp D \\
\\
& A \otimes (B \wp (C \wp D)) & \\
\swarrow \text{id} \otimes \alpha_{\wp} & & \searrow \partial_L \\
A \otimes ((B \wp C) \wp D) & & (A \otimes B) \wp (C \wp D) \\
\downarrow \partial_L & & \downarrow \alpha_{\wp} \\
(A \otimes (B \wp C)) \wp D & \xrightarrow{\partial_L \wp \text{id}} & ((A \otimes B) \wp C) \wp D \\
\\
& (A \wp B) \otimes (C \otimes D) & \\
\swarrow \alpha_{\otimes} & & \searrow \partial_R \\
((A \wp B) \otimes C) \otimes D & & A \wp (B \otimes (C \otimes D)) \\
\downarrow \partial_R \otimes \text{id} & & \downarrow \text{id} \wp \alpha_{\otimes} \\
(A \wp (B \otimes C)) \otimes D & \xrightarrow{\partial_R} & A \wp ((B \otimes C) \otimes D)
\end{array}$$

Diagramme
besser
alignen

$$\begin{array}{ccc}
& ((A \wp B) \wp C) \otimes D & \\
\swarrow \alpha_{\wp} \otimes \text{id} & & \searrow \partial_R \\
(A \wp (B \wp C)) \otimes D & & (A \wp B) \wp (C \otimes D) \\
\downarrow \partial_R & & \downarrow \alpha_{\wp} \\
A \wp ((B \wp C) \otimes D) & \xrightarrow{\text{id} \wp \partial_R} & A \wp (B \wp (C \otimes D))
\end{array}$$

- coherence between the distributors:

$$\begin{array}{ccc}
& (A \wp B) \otimes (C \wp D) & \\
\swarrow \partial_R & & \searrow \partial_L \\
((A \wp B) \otimes C) \wp D & & A \wp (B \otimes (C \wp D)) \\
\downarrow \partial_R \wp \text{id} & & \downarrow \text{id} \wp \partial_L \\
(A \wp (B \otimes C)) \wp D & \xleftarrow{\alpha_{\wp}} & A \wp ((B \otimes C) \wp D)
\end{array}$$

obiges Diagramm rumdrehen

\parr \id sieht kacke aus...

$$\begin{array}{ccc}
& A \otimes ((B \wp C) \otimes D) & \\
\swarrow \alpha_{\otimes} & & \searrow \text{id} \otimes \partial_R \\
(A \otimes (B \wp C)) \otimes D & & A \otimes (B \wp (C \otimes D)) \\
\downarrow \partial_L \otimes \text{id} & & \downarrow \partial_L \\
((A \otimes B) \wp C) \otimes D & \xrightarrow{\partial_R} & (A \otimes B) \wp (C \otimes D)
\end{array}$$

itemize anpassen?

3.3 *-aut. cats

structure an Mellies, CatSemOfLinLog, anpassen; equivalence Srinivasan and Barr definitions

Definition 3.10 (Dual object) Let \mathfrak{C} be a linearly distributive category and $A, A^* \in \text{Obj}(\mathfrak{C})$, then we call A^* left dual (or left linearly adjoint) to A , if there are morphisms $\tau : 1 \rightarrow A^* \wp A$ and $\gamma : A \otimes A^* \rightarrow \perp$, called unit and counit resp., such that the following diagrams commute:

$$\begin{array}{ccc}
A^* & \xlongequal{\text{id}} & A^* \\
\rho_{\wp} \uparrow & & \downarrow \lambda_{\otimes}^{-1} \\
A^* \wp \perp & & 1 \otimes A^* \\
\text{id} \wp \gamma \uparrow & & \downarrow \tau \otimes \text{id} \\
A^* \wp (A \otimes A^*) & \xleftarrow{\partial_R} & (A^* \wp A) \otimes A^*
\end{array}
\quad
\begin{array}{ccc}
A & \xlongequal{\text{id}} & A \\
\lambda_{\wp} \uparrow & & \downarrow \rho_{\otimes}^{-1} \\
\perp \wp A & & A \otimes 1 \\
\gamma \wp \text{id} \uparrow & & \downarrow \text{id} \otimes \tau \\
(A^* \wp A) \otimes A^* & \xleftarrow{\partial_L} & A^* \wp (A \otimes A^*)
\end{array}$$

das geht schöner

We write $(\tau, \gamma) : A^* \dashv A$ and also call A right dual to A^* .

mathabx incompatible with amssymb, stix absolutely broken; repalce dashV with dashv + some index?

Lemma 3.11 1. In an LDC: if $A^* \dashv A$ and $A' \dashv A$, then A^* and A' are isomorphic. We will from now on only talk about the dual object, when equality up to isomorphism is sufficient.

2. In a symmetric LDC: $(\tau, \gamma) : A^* \dashv A \iff (\tau s_{\mathcal{A}}, s_{\mathcal{A}} \gamma) : A \dashv A^*$

proof

[Sri23] Lemma 2.9 (iii)?

Definition 3.12 (*-autonomous categories (Srinivasan [Sri23])) An LDC \mathfrak{C} in which for every Object $A \in \mathfrak{C}$ there exists a left and right dual, resp. $(\tau^*, \gamma^*) : A^* \dashv A$ and $(\tau, \gamma^*) : A \dashv A^*$, is called *-autonomous category.

is Srinivasan the origin of this version?

Definition 3.13 (*-autonomous categories (Barr [Bar95], A)) A *-autonomous category is a biclosed monoidal category \mathfrak{C} with a closed functor $(-)^* : \mathfrak{C} \rightarrow \mathfrak{C}^{\text{opp}}$, which is a strong equivalence of categories.

find a way to make letter part of definition numbering?

definition closed functor, strong equivalence. (strong closed functor?)

Definition 3.14 (Dualizing object) Let \mathfrak{C} be a biclosed monoidal category. An object \perp is called a dualizing object if for every $A \in \text{Obj}(\mathfrak{C})$ the natural map $A \rightarrow \perp \multimap (A \multimap \perp)$ gotten by transposing $\text{id} : A \multimap \perp \rightarrow A \multimap \perp$ twice is an isomorphism.

details (Barr, 99)

Definition 3.15 (*-autonomous categories (Barr, B)) A *-autonomous category is a biclosed category with a dualizing object.

main definition, Barr

Definition 3.16 (*-autonomous categories (Barr, C)) A *-autonomous category is a monoidal category \mathfrak{C} equipped with an equivalence $(-)^* : \mathfrak{C} \rightarrow \mathfrak{C}^{\text{opp}}$ such that there is a natural isomorphism

$$\text{Hom}(A, B^*) \rightarrow \text{Hom}(1, (A \oplus B)^*)$$

Definition 3.17 (*-autonomous categories (Barr, D)) A *-autonomous category is a closed category \mathfrak{C} in the sense of Eilenberg and Kelly (1966) together with an equivalence $(-)^* : \mathfrak{C}^{\text{opp}} \rightarrow \mathfrak{C}$ such that

nachschauen

$$A \multimap (B \multimap C) \cong (A \multimap B) \multimap C$$

where $A \multimap B := A^* \multimap B^*$.

unnötig? C impliziert offensichtlich D, von D nach C sind drei Zeilen.

Theorem 3.18 Barr's definitions are equivalent.

PROOF:

to be added ([Bar95])

Theorem 3.19 *Srinivasan's definition is equivalent to Barr's definitions.*

Weihnachtsbaustelle:

PROOF: Let \mathfrak{A} be a symmetric LDC satisfying definition 3.12.

The mapping $A \mapsto A^*$ is an isomorphism $\text{Obj}(\mathfrak{A}) \rightarrow \text{Obj}(\mathfrak{A}^{\text{opp}})$ between objects and defines a Functor by extending it to morphisms with the following diagram:

change definition to diagram?

$$\text{Hom}_{\mathfrak{A}}(A, B) \ni f \mapsto f^* \in \text{Hom}_{\mathfrak{A}^{\text{opp}}}(A^*, B^*) = \text{Hom}_{\mathfrak{A}}(B^*, A^*)$$

$$\begin{aligned} f^* : B^* &\xrightarrow{\lambda_{\otimes}^{-1}} 1 \otimes B^* \xrightarrow{\eta_A \otimes \text{id}} (A^* \mathbin{\mathfrak{A}} A) \otimes B^* \xrightarrow{(\text{id} \mathbin{\mathfrak{A}} f) \otimes \text{id}} (A^* \mathbin{\mathfrak{A}} B) \otimes B^* \\ &\xrightarrow{\partial_R} A^* \mathbin{\mathfrak{A}} (B \otimes B^*) \xrightarrow{\text{id} \mathbin{\mathfrak{A}} \varepsilon_B} A^* \mathbin{\mathfrak{A}} 1 \xrightarrow{\rho_{\mathfrak{A}}} A^* \end{aligned}$$

This mapping of morphisms is also a bijection:

$$\begin{aligned} f &= \lambda_{\mathfrak{A}, B} \circ (\varepsilon_A \mathbin{\mathfrak{A}} \text{id}_B) \circ ((\text{id}_A \otimes f^*) \mathbin{\mathfrak{A}} \text{id}_B) \circ \partial_L \circ (\text{id}_A \otimes \eta_B) \circ \rho_{\otimes, A}^{-1} \\ &= \lambda_{\mathfrak{A}, B} \circ (\varepsilon_A \mathbin{\mathfrak{A}} \text{id}_B) \\ &\quad \circ \left(\left(\text{id}_A (\otimes \rho_{\mathfrak{A}, A}) \circ (\text{id}_{A^*} \mathbin{\mathfrak{A}} \varepsilon_B) \circ (\text{id}_{A^*} \mathbin{\mathfrak{A}} (f \otimes \text{id}_{B^*})) \circ \partial_R \circ (\eta_A \otimes \text{id}) \circ \lambda_{\otimes, B}^{-1} \right) \mathbin{\mathfrak{A}} \text{id}_{B^*} \right) \\ &\quad \circ \partial_L \circ (\text{id}_A \otimes \eta_B) \circ \rho_{\otimes, A}^{-1} \end{aligned}$$

(maybe suppress brackets for stuff like $A \otimes 1 \otimes 1$)

$$\begin{array}{ccccc} A & \xrightarrow{(\text{id} \otimes \eta_B) \circ \rho_{\otimes}^{-1}} & A \otimes (B^* \otimes B) & \xrightarrow{\text{id} \otimes ((\eta_A \otimes \text{id}) \mathbin{\mathfrak{A}} \text{id})} & A \otimes (((A^* \mathbin{\mathfrak{A}} A) \otimes B^*) \mathbin{\mathfrak{A}} B) \\ \downarrow \rho_{\otimes}^{-1} \circ \rho_{\otimes}^{-1} & & & & \downarrow \text{id} \otimes (((\text{id} \mathbin{\mathfrak{A}} f) \otimes \text{id}) \mathbin{\mathfrak{A}} \text{id}) \\ (A \otimes 1) \otimes 1 & & & & A \otimes (((A^* \mathbin{\mathfrak{A}} B) \otimes B^*) \mathbin{\mathfrak{A}} B) \\ \downarrow \text{id} \otimes \eta_A \otimes \eta_B & & & & \downarrow \text{id} \otimes (\partial_R \mathbin{\mathfrak{A}} \text{id}) \\ (A \otimes (A^* \mathbin{\mathfrak{A}} A)) \otimes (B^* \mathbin{\mathfrak{A}} B) & & & & A \otimes ((A^* \mathbin{\mathfrak{A}} (B \otimes B^*)) \mathbin{\mathfrak{A}} B) \\ \downarrow \partial_L \otimes \text{id} & & & & \downarrow \text{id} \otimes (\partial_R \mathbin{\mathfrak{A}} \text{id}) \\ ((A \otimes A^*) \mathbin{\mathfrak{A}} A) \otimes (B^* \mathbin{\mathfrak{A}} B) & & & & A \otimes ((A^* \mathbin{\mathfrak{A}} (B \otimes B^*)) \mathbin{\mathfrak{A}} B) \\ \downarrow (\lambda_{\mathfrak{A}} \otimes \text{id}) \circ ((\varepsilon_A \mathbin{\mathfrak{A}} f) \otimes \text{id}) & & & & \downarrow \text{id} \otimes (\rho_{\mathfrak{A}} \circ (\text{id} \mathbin{\mathfrak{A}} \varepsilon_B \mathbin{\mathfrak{A}} \text{id})) \\ B \otimes (B^* \mathbin{\mathfrak{A}} B) & \xrightarrow{\lambda_{\mathfrak{A}} \circ \varepsilon_B \circ \partial_L} & B & \xleftarrow{\lambda_{\mathfrak{A}} \circ \varepsilon_B \circ \partial_L} & A \otimes (A^* \mathbin{\mathfrak{A}} B) \end{array}$$

f

does this commute?

$$\begin{array}{c}
 A \xrightarrow{\rho_{\otimes}^{-1}} A \otimes 1 \xrightarrow{\text{id} \otimes \eta_B} A \otimes (B^* \wp B) \xrightarrow{\partial_L} (A \otimes B^*) \wp B \\
 \xrightarrow{(\text{id} \otimes f^*) \wp \text{id}} (A \otimes A^*) \wp B \xrightarrow{\varepsilon_A \wp \text{id}} \perp \wp B \xrightarrow{\lambda_{\wp}} B
 \end{array}
 \quad \square$$

is this really f?

non-symmetric case to be added (should already be in there, never used symmetry)

what did I need Barr's definition for, again?

4 CatSem of LinLog

Cats as LinLog semantics and LinLog syntax as cat

add refs

References

- [Bar95] Michael Barr. “Nonsymmetric *-autonomous categories”. In: *Theoretical Computer Science* 139.1 (1995), pp. 115–130. ISSN: 0304-3975. DOI: [https://doi.org/10.1016/0304-3975\(94\)00089-2](https://doi.org/10.1016/0304-3975(94)00089-2). URL: <https://www.sciencedirect.com/science/article/pii/0304397594000892>.
- [Bra96] Torben Braüner. “Introduction to Linear Logic”. In: (1996).
- [Gir87] Jean-Yves Girard. “Linear logic”. In: *Theoretical Computer Science* 50.1 (1987), pp. 1–101. ISSN: 0304-3975. DOI: [https://doi.org/10.1016/0304-3975\(87\)90045-4](https://doi.org/10.1016/0304-3975(87)90045-4). URL: <https://www.sciencedirect.com/science/article/pii/0304397587900454>.
- [Sri23] Priyaa Srinivasan. “Dagger linear logic and categorical quantum mechanics”. PhD thesis. Mar. 2023.

Todo list

■ what am I even writing about?	1
■ feels kinda rambling... combine intuition with syntax definition below	1
■ (explicit interpretation?! shared pool between people containing Pika and Mew -> you have to get rid of one to use the other?)	2
■ interpretation of ?wn.	2
■ interpretation of constants	2
■ Quellen, Quellen, Quellen!!!	2
■ section needs serious reformatting...	2
■ main source: IntroLinLog	2
■ definition: "proof"; "cut elimination"	2
■ MAYBE to be added: \forall and \exists. Do we want predicate logic?	2
■ upper vertex of identity?	3
■ permanente Lösung für empty axiom finden	3
■ move negation to other place?	3
■ proofsymbols /	4
■ remark unnötig?	4
■ proofsymbols	6
■ proofsymbols なんですか :(.	7
■ add "spoiler" to categoru theoru	7
■ tabularx format	8
■ Problem: Resulting proofs for the \mathfrak{A}_L -cases are illegal under current formulation of the \mathfrak{A}_L -rule. (Also: they're kinda the same?) Possible solutions: 1: add exchanges to transformation (easy; ugly) 2: find generalized \mathfrak{A}_L -rule (more difficult; possibly uglier)	8
■ To be added: \otimes_R -cases. Maybe combine \mathfrak{A}_L - and \otimes_R -cases into one? (Requires generalized formulation of the rules)	8
■ To be added: \mathfrak{A}_L - and \otimes_R -cases (same problems as above)	9
■ proofsymbols	9
■ put that stuff back into a normal center instead of tabularx?	10
■ think about the deltas...: aufteilen in zwei Regeln mit einseitigen Deltas für A und B oder allgemein halten und viele exchanges drunter klatschen?	10
■ why are they called η -expansion steps?	10
■ format (page break)	10

gone to 2.12	11
that's just changing the contexts Γ_2 and Γ'_2	11
gone to 2.12	11
gone to 2.12	11
sources for basic definitions. (nlab proly doesn't suffice)	11
extra definition for opposite structure?	12
definition: equivalence of categories	13
source: Brandenburg	13
maybe index Schreibweise?	13
cut triangle diagrams or define \bullet	13
Diagramme verschönern	14
fix associators throughout the document (direction)	14
lengthy, rambling... muss eleganter gehen	14
definitely mixed up the right and left homs there	14
following definitions proly unnötig:	14
Diagramme besser alignen	16
obiges Diagramm rumdrehen	17
$\backslash\text{parr}$ $\backslash\text{id}$ sieht kacke aus...	17
itemize anpassen?	17
structure an Mellies, CatSemOfLinLog, anpassen; equivalence Srinivasan and Barr definitions	17
das geht schöner	17
mathabx incompatible with amssymb, stix absolutely broken; repalce dashV with dashv + some index?	18
proof	18
[Sri23] Lemma 2.9 (iii)?	18
is Srinivasan the origin of this version?	18
find a way to make letter part of definition numbering?	18
definition closed functor, strong equivalence. (strong closed functor?)	18
details (Barr, 99)	18
main definition, Barr	18
nachschauen	18
unnötig? C impliziert offensichtlich D , von D nach C sind drei Zeilen.	18
to be added ([Bar95])	19

■ Weihnachtsbaustelle:	19
■ change definition to diagram?	19
■ (maybe suppress brackets for stuff like $A \otimes 1 \otimes 1$)	19
■ does this commute?	19
■ is this really f?	20
■ non-symmetric case to be added (should already be in there, never used symmetry)	20
■ what did I need Barr's definition for, again?	20
■ Cats as LinLog semantics and LinLog syntax as cat	20
■ add refs	20