Typeset using LaTeX default style in AASTeX63

Draft version September 25, 2020

Residuals Summary

Henry A. Prager[1]

[1] *Department of Physics, New Mexico Tech, Socorro, NM 87801*

(Dated: September 25, 2020)

ABSTRACT

1. Q

henry.prager@student.nmt.edu

1.1. *M0*

1.1.1. *First Residual*

$$\text{Residual Fit}: \log \dot{M}_{\text{calc}} - \log \dot{M}_{\text{obsv}} = -12.745618196936379 + 5.8076081856640105 \log P \tag{1}$$




Figure 1. The results of plotting the first residuals versus L, M, P, and Q.

$$\text{Resid. vs Parameter Fit}: \log \dot{M}_{\text{calc}} - \log \dot{M}_{\text{obsv}} = -12.745618196936379 + 5.8076081856640105 \log P \tag{2}$$

$$\text{Resid. vs Parameter Fit}: \log \dot{M}_{\text{calc}} - \log \dot{M}_{\text{obsv}} = -12.174389758004052 + 3.451383142617217 \log L \tag{3}$$

$$\text{Resid. vs Parameter Fit}: \log \dot{M}_{\text{calc}} - \log \dot{M}_{\text{obsv}} = 1.2054855398116522 + 5.648211407977223 \log Q \tag{4}$$

$$\text{Resid. vs Parameter Fit}: \log \dot{M}_{\text{calc}} - \log \dot{M}_{\text{obsv}} = -0.01858475071591059 + 1.5256809613105102 \log M \tag{5}$$

1.1.2. *First Residual vs. New Residual*




Figure 2. The results of plotting the first residuals versus the new residuals.

1.1.3. *New Residual*

1.2. *M1*

1.3. *C0*

1.4. *C1*


2. QX

2.1. *M0*


2.2. *M1*

2.3. *C0*


2.4. *C1*




(a)




(b)




(c)




(d)




(e)



(f)



(g)



(h)

Figure 3. The results of plotting both residuals versus L, M, P, and Q.