

COMBATING THE FRESHWATER CRISIS:



In vivo Studies Investigating the Elongation of the Loop
of Henle


Christy Au-Yeung & Bushra Haque

FRESHWATER CRISIS


What's the problem?

2

1 AVAILABILITY



2 WATER SCARCITY



2.7 billion people

Experience water scarcity
at least one month of the
year

3 WATER DEMAND



Human Population



Global Temperatures




Water Availability

Water demand is expected to increase
by 55% between 2000 and 2050

Background



Dipodomys ordii (kangaroo rat)




Production of hyperosmotic urine
from elongated loop of Henle

- Kangaroo rats are able to consume seawater while maintaining healthy conditions
- Maximum urine electrolyte concentration reaching ~1200 mN (more than 2x sea water levels!)

Background


4

- Long hairpin-like structure within each nephron
- Comprised of three sections
 1. Descending limb
 2. Thin ascending limb
 3. Thick ascending limb
- Concentrates urine through osmotic gradients
 - Length determines size of gradient




Experimental Setup

5




In silico studies

- Search & identification of Loop of Henle development genes using Gene Ontology (GO) in MGI database
- Use of comparative genomics to assess genetic composition of Wnt7b between kangaroo rat *and* laboratory mouse
 - Prior to mutational analyses



Genome Editing: CRISPR-Cas9

- Cas9 – Endonuclease that initiates double-stranded DNA breaks
- gRNA – Complementary to the *Wnt7b* to add/remove genomic regions
- Homology directed repair (HDR)



In vitro studies

- Characterize molecular and cellular consequences
- Human murine-derived renal epithelial and kidney cells of 1-month old M. musculus
 - Cell viability
 - qPCR to determine mutation
- Organ cultures
 - Compare length of Loop of Henle
 - Laminin and Tamm-Horsfall Protein immunostaining





Image of LoH in organ culture immunostained with laminin (green) and Tamm-Horsfall Protein (red)

In vivo studies

- Murine models containing genetic mutation will be nourished with diluted salt water
- **Primary objective:** Determine if models can consume salt water
- **Secondary objective:** Determine any adverse health effects


Analyze various factors

- Length of loop of Henle: antibodies against AQP1 and CIC-K
- Blood samples: measure serum sodium levels and osmolarity
- Other health impacts: salt poisoning, weight changes



Future Applications

- Advances in genetic engineering
- Saltwater-tolerant generation of humans with elongated Loop of Henle
 - More sustainable than desalination technologies
 - Sustainable lifestyle for future generations



- All in efforts to conserve the limited amounts of freshwater present on Earth
- Drastic measures must be taken now to mitigate unfortunate outcomes

Thank You for Listening
QUESTIONS?

References

Smedley, T. Is the world running out of fresh water?

<https://www.bbc.com/future/article/20170412-is-the-world-running-out-of-fresh-water> (accessed Mar 1, 2020).

Roser, M.; Ortiz-Ospina, E.; Ritchie, H. Life Expectancy. Our World in Data 2013.

Hydrofinity. Why Desalination is Not the Answer to the World's Water Issues

<http://www.hydrofinity.com/blog/why-desalination-is-not-the-answer-to-the-worlds-water-issues> (accessed Mar 1, 2020).

Maximum physiological concentration of sodium in human urine | Journal of Applied Physiology

<https://journals.physiology.org/doi/abs/10.1152/jappl.1959.14.6.1036> (accessed Mar 1, 2020).

Evaporative Water Loss in Desert Rodents in Their Natural Habitat - Schmidt-Nielsen - 1950 - Ecology - Wiley Online Library

<https://esajournals.onlinelibrary.wiley.com/doi/10.2307/1931362> (accessed Mar 1, 2020).

Sands, J. M.; Layton, H. E. The Physiology of Urinary Concentration: An Update. Semin Nephrol 2009, 29 (3), 178–195.

<https://doi.org/10.1016/j.semnephrol.2009.03.008>.

Genetics Home Reference. WNT7B gene <https://ghr.nlm.nih.gov/gene/WNT7B> (accessed Mar 1, 2020).

Yu, J.; Carroll, T. J.; Rajagopal, J.; Kobayashi, A.; Ren, Q.; McMahon, A. P. A Wnt7b-Dependent Pathway Regulates the Orientation of Epithelial Cell Division and Establishes the Cortico-Medullary Axis of the Mammalian Kidney. Development 2009, 136 (1), 161–171.

<https://doi.org/10.1242/dev.022087>.

References (Contd.)

Hall, B.; Cho, A.; Limaye, A.; Cho, K.; Khillan, J.; Kulkarni, A. B. Genome Editing in Mice Using CRISPR/Cas9 Technology. *Curr Protoc Cell Biol* 2018, 81 (1), e57. <https://doi.org/10.1002/cpcb.57>.

Ramsbottom, S. A.; Molinari, E.; Srivastava, S.; Silberman, F.; Henry, C.; Alkanderi, S.; Devlin, L. A.; White, K.; Steel, D. H.; Saunier, S.; et al. Targeted Exon Skipping of a CEP290 Mutation Rescues Joubert Syndrome Phenotypes in Vitro and in a Murine Model. *PNAS* 2018, 115 (49), 12489–12494. <https://doi.org/10.1073/pnas.1809432115>.

Sebinger, D. D. R.; Unbekandt, M.; Ganeva, V. V.; Ofenbauer, A.; Werner, C.; Davies, J. A. A Novel, Low-Volume Method for Organ Culture of Embryonic Kidneys That Allows Development of Cortico-Medullary Anatomical Organization. *PLoS One* 2010, 5 (5). <https://doi.org/10.1371/journal.pone.0010550>.

Chang, C.; Davies, J. A. In Developing Mouse Kidneys, Orientation of Loop of Henle Growth Is Adaptive and Guided by Long-range Cues from Medullary Collecting Ducts. *J Anat* 2019, 235 (2), 262–270. <https://doi.org/10.1111/joa.13012>.

Davidson, A. J. Mouse Kidney Development. In *StemBook*; Harvard Stem Cell Institute: Cambridge (MA), 2008.

Fereidoun, H.; Bahram, A.; Sadraddin, K.; Ahmad, R.; Pouria, H. Determination of Tolerable and Toxic Salt Concentrations in Drinking Water in Rat, Pig, and Rabbit. *Toxicological & Environmental Chemistry* 2008, 90 (6), 1115–1123. <https://doi.org/10.1080/02772240701850707>.