

ERDŐS SIMILARITY FOR DISTANCE SETS: FINITE PATTERNS, A PINNED OBSTRUCTION, AND AN OPEN WINDOW

HARALD SCHILLY

ABSTRACT. For a compact set $E \subset \mathbb{R}^d$, $d \geq 2$, let $\Delta(E) = \{|x - y| : x, y \in E\}$ be its distance set. We ask whether distance sets of sets of large Hausdorff dimension must contain a nontrivial affine copy $x + s\{2^{-n} : n \geq 1\}$ of the dyadic sequence. This is a distance-set version of the Erdős similarity problem. We combine two recent results announced by OpenAI: the Falconer distance theorem in all dimensions, and the dyadic case of the Erdős similarity conjecture. Assuming these, we show the following. (A) If $\dim_{\text{H}} E > d/2$, then $\Delta(E)$ contains affine copies of every finite set, for a positive-measure family of translations at every small scale. (B) There is a compact set $K \subset \mathbb{R}^d$ of positive volume, hence of full dimension, whose distance set from the origin contains no affine copy of the dyadic sequence. The full distance set $\Delta(K)$ does contain one, and the origin is the only point of $\mathbb{R}^d$ ($d \geq 2$) from which the obstruction occurs. Since the Mattila–Sjölin theorem settles the question above dimension $(d + 1)/2$, it remains open exactly in the window $d/2 < \dim_{\text{H}} E \leq (d + 1)/2$. A negative answer there would produce distance sets of positive measure with empty interior. Both theorems are formalized in Lean 4 with Mathlib, with the two inputs as explicit hypotheses. We also re-checked OpenAI’s formal proofs of both inputs with the Lean kernel.

1. INTRODUCTION

1.1. Two problems about large sets. A set $P \subset \mathbb{R}$ is *measure universal* if every measurable $S \subset \mathbb{R}$ of positive Lebesgue measure contains a nontrivial affine copy $x + sP$ ($x \in \mathbb{R}$, $s \neq 0$). Steinhaus’ theorem [16] implies that every finite set is measure universal. Erdős conjectured [3] that no infinite set is. The conjecture is known for slowly decaying sequences [5] and for many structured sets [1, 9]. The geometric sequences $\{q^n\}$, and above all the dyadic sequence

$$\mathcal{D} = \{2^{-n} : n \geq 1\},$$

were long described as the main bottleneck [8]. In September and October 2026, OpenAI released a collection of manuscripts produced by an internal AI model. It includes a proof that $\mathcal{D}$ is not measure universal [12] and an extension to every ratio $q \in (0, 1)$ [14].

Falconer’s distance problem [6] asks for the smallest α such that every compact $E \subset \mathbb{R}^d$ with $\dim_{\text{H}} E > \alpha$ has a distance set

$$\Delta(E) = \{|x - y| : x, y \in E\}$$

of positive Lebesgue measure. Falconer showed that $d/2$ would be optimal and conjectured it. Partial progress includes [18, 4, 7]. The same OpenAI release contains a proof of the full conjecture in every dimension $d \geq 2$ [13]. Mattila and Sjölin [11] showed that $\dim_{\text{H}} E > (d + 1)/2$ even forces $\Delta(E)$ to contain an interval.

Date: October 2026.

2020 Mathematics Subject Classification. 28A80 (primary); 28A75, 42B10, 03B35.

Key words and phrases. Erdős similarity problem, distance sets, Falconer conjecture, Hausdorff dimension, formal verification, Lean.

1.2. The question. The Falconer theorem says that $\Delta(E)$ is large in measure, while the dyadic theorem says that being large in measure is not enough to contain $\mathcal{D}$. Distance sets, however, have additional multi-scale structure: $\Delta(E) \supseteq \Delta(E \cap B)$ for every ball B , so $\Delta(E) \cap [0, r)$ has positive measure for every $r > 0$ as soon as the dimension condition holds locally. This motivates:

Question A (Erdős similarity for distance sets). Let $d \geq 2$ and let $E \subset \mathbb{R}^d$ be compact with $\dim_{\text{H}} E > d/2$. Must $\Delta(E)$ contain a nontrivial affine copy of $\mathcal{D}$?

We could not find this question, or any study of Erdős-type patterns inside distance sets, in the literature; it is not discussed in the recent survey [8].

1.3. Results. Throughout, Theorems A and B are *conditional* on the two inputs stated precisely in Section 2. Theorem 2.1 is the Falconer theorem of [13], and Theorem 2.2 is the dyadic avoidance theorem of [12].

Theorem A (Finite patterns). Assume Theorem 2.1. Let $d \geq 2$, let $E \subset \mathbb{R}^d$ be compact with $\dim_{\text{H}} E > d/2$, and let $F \subset \mathbb{R}$ be finite. There is $\delta > 0$ such that for every $|s| < \delta$,

$$\mathcal{L}^1(\{x \in \mathbb{R} : x + sF \subseteq \Delta(E)\}) > 0.$$

In particular $\Delta(E)$ contains affine copies $x + sF$ with $0 < s < \delta'$ and with $-\delta' < s < 0$, for every $\delta' > 0$.

Theorem B (A pinned obstruction). Assume Theorem 2.2. Let $d \geq 1$ and $\eta \in (0, 1)$. There is a compact set K in the closed unit ball of $\mathbb{R}^d$ with $0 \in K$ and

$$\mathcal{L}^d(K) \geq (1 - \eta)^d \mathcal{L}^d(B(0, 1)) > 0, \quad \text{hence } \dim_{\text{H}} K = d,$$

such that the pinned distance set $\Delta_0(K) = \{|y| : y \in K\}$ contains no nontrivial affine copy of $\mathcal{D}$, whereas the full distance set $\Delta(K)$ contains one. If $d \geq 2$, then for every $p \in \mathbb{R}^d \setminus \{0\}$ the pinned distance set $\Delta_p(K) = \{|p - y| : y \in K\}$ contains a nondegenerate interval, hence an affine copy of $\mathcal{D}$.

So pinned distance sets are not universal hosts for $\mathcal{D}$, even from a pin inside a set of full dimension and nearly full volume in the ball. The obstruction, however, sits at one single point. Theorems A and B together with the Mattila–Sjölin theorem show where Question A is undecided:

Proposition C. Let $d \geq 2$ and let $E \subset \mathbb{R}^d$ be compact. If $\dim_{\text{H}} E > (d+1)/2$, or if $\mathcal{L}^d(E) > 0$, then $\Delta(E)$ contains an affine copy $a + sP$ of every bounded set $P \subset [0, \infty)$; in particular of $\mathcal{D}$. If Question A has a negative answer for some E with $d/2 < \dim_{\text{H}} E \leq (d+1)/2$, then $\Delta(E)$ has positive measure but empty interior.

We are not aware of a general result that guarantees nonempty interior of $\Delta(E)$ below the Mattila–Sjölin threshold $(d+1)/2$ (see [10] for background). Under that reading, a negative answer to Question A in the window would also give new examples of distance sets with positive measure and empty interior. Section 6 lists further questions.

1.4. Formalization and status of the inputs. Theorems A and B are formalized in full in Lean 4 [2] with Mathlib [17]. The two inputs enter as explicit hypotheses whose Lean types are copied verbatim from OpenAI’s formal statements, and the formal proofs use no axioms beyond Lean’s standard ones. The two inputs come from an AI-generated, unrefereed manuscript collection whose authors state that some results “could have issues”. OpenAI also provides Lean proofs of both inputs. We rebuilt the relevant part of that library ourselves and checked

with the Lean kernel that these proofs establish exactly our hypotheses, using only the standard axioms (Section 7). This makes the dependence on the inputs as transparent as we can. It does not replace independent mathematical review of [13, 12].

2. NOTATION AND THE TWO INPUTS

$\mathcal{L}^d$ is Lebesgue measure on $\mathbb{R}^d$, $|\cdot|$ the Euclidean norm, $B(0, 1)$ the open unit ball, and $\dim_{\text{H}}$ Hausdorff dimension. For $E \subset \mathbb{R}^d$ and $p \in \mathbb{R}^d$ we write $\Delta(E) = \{|x - y| : x, y \in E\}$ and $\Delta_p(E) = \{|p - y| : y \in E\}$. An *affine copy* of $P \subset \mathbb{R}$ is a set $x + sP$ with $x \in \mathbb{R}$ and $s \neq 0$.

Theorem 2.1 (Falconer distance theorem [13]). *Let $d \geq 2$ and let $E \subset \mathbb{R}^d$ be compact with $\dim_{\text{H}} E > d/2$. Then $\mathcal{L}^1(\Delta(E)) > 0$.*

Theorem 2.2 (Dyadic avoidance [12]). *For every $\eta \in (0, 1)$ there is a compact set $A \subseteq [0, 1]$ with $\mathcal{L}^1(A) > 1 - \eta$ such that for every $x \in \mathbb{R}$ and every $s \neq 0$ there is $n \geq 1$ with $x + s2^{-n} \notin A$.*

3. FINITE PATTERNS: PROOF OF THEOREM A

The following is the standard Steinhaus-type argument (compare the introduction of [12]). We include it because Theorem A needs the version with a positive-measure set of translations at every small scale.

Lemma 3.1. *Let $S \subset \mathbb{R}$ be measurable with $0 < \mathcal{L}^1(S) < \infty$ and let $F \subset \mathbb{R}$ be finite. There is $\delta > 0$ such that $\mathcal{L}^1(\{x : x + sF \subseteq S\}) > 0$ for all $|s| < \delta$.*

Proof. For $a \in F$ let $T_{a,s} = S - sa$. Translation is continuous in measure: $\mathcal{L}^1(T_{a,s} \triangle S) \rightarrow 0$ as $s \rightarrow 0$, since $\|\mathbf{1}_S(\cdot + h) - \mathbf{1}_S\|_{L^1} \rightarrow 0$ as $h \rightarrow 0$. Choose $\delta > 0$ with $\sum_{a \in F} \mathcal{L}^1(T_{a,s} \triangle S) < \mathcal{L}^1(S)$ for $|s| < \delta$. Since

$$S \subseteq \{x : x + sF \subseteq S\} \cup \bigcup_{a \in F} (S \setminus T_{a,s}),$$

we get $\mathcal{L}^1(S) \leq \mathcal{L}^1(\{x : x + sF \subseteq S\}) + \sum_{a \in F} \mathcal{L}^1(T_{a,s} \triangle S)$, and the claim follows. $\square$

Proof of Theorem A. $\Delta(E)$ is the image of the compact set $E \times E$ under the continuous map $(x, y) \mapsto |x - y|$, so it is compact. In particular it is measurable with finite measure, and $\mathcal{L}^1(\Delta(E)) > 0$ by Theorem 2.1. Lemma 3.1 with $S = \Delta(E)$ gives the first claim. For the second, apply it with $s = \pm \min(\delta, \delta')/2$; a set of positive measure is nonempty. $\square$

Remark 3.2. The set $\{(x, s) : x + sF \subseteq \Delta(E)\}$ is closed, and by Tonelli's theorem it has positive planar measure. Theorem A uses nothing about distance sets beyond compactness and positive measure. Its point is the contrast with Theorem B and Question A: the finite-to-infinite gap of the Erdős similarity problem reappears inside Falconer distance sets.

4. THE PINNED OBSTRUCTION: PROOF OF THEOREM B

Lemma 4.1 (Adding the limit point). *Let $A \subset \mathbb{R}$ contain no nontrivial affine copy of $\mathcal{D}$. Then neither does $A \cup \{0\}$.*

Proof. Suppose $x + s2^{-n} \in A \cup \{0\}$ for all $n \geq 1$, with $s \neq 0$. The points $x + s2^{-m}$, $m \geq 0$, are pairwise distinct. If none of them is 0, then $x + s\mathcal{D} \subseteq A$, a contradiction. Otherwise $x + s2^{-m} = 0$ for exactly one $m \geq 0$. The tail $\{x + s2^{-(m+n)} : n \geq 1\} = x + (s2^{-m})\mathcal{D}$ is then an affine copy of $\mathcal{D}$ avoiding 0, hence contained in A , again a contradiction. $\square$

Lemma 4.2 (Radial sets). *Let $d \geq 1$ and let $A \subseteq [0, 1]$ be compact with $\mathcal{L}^1(A) = c > 0$. Then $K_A = \{y \in \mathbb{R}^d : |y| \in A\}$ is compact, contained in the closed unit ball, and $\mathcal{L}^d(K_A) \geq c^d \mathcal{L}^d(B(0, 1))$.*

Proof. Compactness and boundedness are clear. In polar coordinates,

$$\mathcal{L}^d(K_A) = d \mathcal{L}^d(B(0, 1)) \int_A r^{d-1} dr.$$

Since $r \mapsto r^{d-1}$ is nondecreasing,

$$\begin{aligned} \int_A r^{d-1} dr - \int_0^c r^{d-1} dr &= \int_{A \setminus [0, c]} r^{d-1} dr - \int_{[0, c] \setminus A} r^{d-1} dr \\ &\geq c^{d-1} (\mathcal{L}^1(A \setminus [0, c]) - \mathcal{L}^1([0, c] \setminus A)) = c^{d-1} (\mathcal{L}^1(A) - c) = 0, \end{aligned}$$

so $\int_A r^{d-1} dr \geq c^d/d$. $\square$

Lemma 4.3. *If $K \subset \mathbb{R}^d$ has $\mathcal{L}^d(K) > 0$, then $\dim_H K = d$.*

Proof. On $\mathbb{R}^d$ the d -dimensional Hausdorff measure (normalized as in Mathlib, for the sup-norm) is a positive multiple of $\mathcal{L}^d$, and the Euclidean and sup norms are bi-Lipschitz equivalent. Hence $\mathcal{H}^d(K) > 0$ and $\dim_H K \geq d$. The reverse inequality is trivial. $\square$

Lemma 4.4 (Steinhaus). *If $K \subset \mathbb{R}^d$ ($d \geq 1$) is compact with $\mathcal{L}^d(K) > 0$, then $[0, \varepsilon] \subseteq \Delta(K)$ for some $\varepsilon > 0$. Consequently $\Delta(K) \supseteq \frac{\varepsilon}{2} \mathcal{D}$.*

Proof. By Steinhaus' theorem $K - K \supseteq B(0, \varepsilon)$ for some $\varepsilon > 0$. For $0 \leq r < \varepsilon$ write $re_1 = x - y$ with $x, y \in K$; then $r = |x - y| \in \Delta(K)$. $\square$

Proof of Theorem B. Let A be given by Theorem 2.2 for η , and put $K = \{y : |y| \in A \cup \{0\}\}$. The set $A \cup \{0\} \subseteq [0, 1]$ is compact, so K is compact and lies in the closed unit ball, and $0 \in K$. Since $K \supseteq K_A$ and $\mathcal{L}^1(A) > 1 - \eta$, Lemma 4.2 gives $\mathcal{L}^d(K) \geq (1 - \eta)^d \mathcal{L}^d(B(0, 1)) > 0$, and Lemma 4.3 gives $\dim_H K = d$. Further, $\Delta_0(K) = \{|y| : y \in K\} \subseteq A \cup \{0\}$, which contains no affine copy of $\mathcal{D}$ by Lemma 4.1. Lemma 4.4 shows that $\Delta(K)$ contains one.

For the last sentence, let $d \geq 2$ and $p \neq 0$, and pick $\rho \in A$ with $\rho > 0$ (possible since $\mathcal{L}^1(A) > 0$). The sphere $\{|y| = \rho\} \subseteq K$ is connected and the continuous function $y \mapsto |p - y|$ on it attains the values $|\rho - |p||$ and $\rho + |p|$. Hence $\Delta_p(K)$ contains the interval $[|\rho - |p||, \rho + |p|]$ of length $2 \min(\rho, |p|) > 0$. Any nondegenerate interval $[a, b]$ contains $a + (b - a)\mathcal{D}$. $\square$

Remark 4.5 (Other ratios). The proof uses only two properties of $\mathcal{D}$: the avoidance property, and the fact that tails of $\mathcal{D}$ are affine copies of $\mathcal{D}$. Both hold for $\{q^n : n \geq 1\}$ with $q \in (0, 1)$. So Theorem B holds for every geometric sequence, conditional on the geometric case [14], which is not part of the Lean formalization.

5. ABOVE THE MATTILA–SJÖLIN THRESHOLD: PROOF OF PROPOSITION C

If $\dim_H E > (d + 1)/2$, the Mattila–Sjölin theorem [11] gives an interval $[a, b] \subseteq \Delta(E)$ with $a < b$. If $\mathcal{L}^d(E) > 0$, Lemma 4.4 gives one. For bounded $P \subseteq [0, M]$ with $M > 0$, we have $a + \frac{b-a}{M}P \subseteq [a, b]$. If $\Delta(E)$ contains no affine copy of $\mathcal{D}$, it therefore has empty interior. If moreover $\dim_H E > d/2$, then $\mathcal{L}^1(\Delta(E)) > 0$ by Theorem 2.1. $\square$

6. OPEN QUESTIONS

- (1) Question A in the window $d/2 < \dim_H E \leq (d + 1)/2$. A positive answer would need multi-scale structure of distance sets beyond positive measure. A negative answer would need a construction controlling all distances of a set of dimension $> d/2$ simultaneously.

- (2) *All pins.* Is there a compact $E \subset \mathbb{R}^d$ with $\mathcal{L}^d(E) > 0$ (or merely $\dim_{\text{H}} E > d/2$) such that $\Delta_p(E)$ contains no affine copy of $\mathcal{D}$ for *every* $p \in E$? The complement of such a set must contain, around every point of E , entire spheres whose radii meet every affine copy of $\mathcal{D}$. This suggests a connection with spherical maximal function estimates. Pinned Falconer-type theorems [15, 7] give pins with large pinned distance sets, but say nothing about such patterns.
- (3) The same questions for general infinite sets P , for instance for every P known to be non-universal.

7. FORMALIZATION

The Lean development accompanies this note. It is pinned to Lean 4 v4.34.1 and Mathlib commit d13f23b, the same versions as OpenAI’s library, so the input statements typecheck identically. It consists of four files:

- `Statements.lean`: the propositions `FalconerStatement` and `DyadicAvoidanceStatement`, copied verbatim from OpenAI’s statement files `FalconerAllDimensions.lean` and `DyadicAvoidance.lean`, and the definitions of distance sets and of “contains an affine copy of $\mathcal{D}$ ”.
- `FinitePatterns.lean`: Lemma 3.1 and Theorem A (`finite_pattern_distSet`, `exists_finite_pattern_distSet`).
- `RadialVolume.lean`: Lemma 4.2, including the rearrangement inequality and the polar-coordinate formula.
- `PinnedObstruction.lean`: Lemmas 4.1, 4.3, 4.4 and Theorem B (`pinned_dyadic_obstruction`). The formal statement is:

```
theorem pinned_dyadic_obstruction (hDy : DyadicAvoidanceStatement) {d : ℕ}
  (hd : 1 ≤ d) {η : ℝ} (hη0 : 0 < η) (hη1 : η < 1) :
  ∃ K : Set (EuclideanSpace ℝ (Fin d)),
    IsCompact K ∧ K ⊆ closedBall 0 1 ∧
    (0 : EuclideanSpace ℝ (Fin d)) ∈ K ∧
    ENNReal.ofReal ((1 - η) ^ d) * volume (ball 0 1) ≤ volume K ∧
    0 < volume K ∧ dimH K = d ∧
    ¬ ContainsDyadicCopy (pinnedDistSet 0 K) ∧ ContainsDyadicCopy (distSet K) ∧
    (2 ≤ d → ∀ p, p ≠ 0 → ContainsDyadicCopy (pinnedDistSet p K))
```

Not formalized are the remark on other ratios and Proposition C, which relies on the Mattila–Sjölin theorem (not available in Mathlib).

Re-checking the inputs. A script in the repository copies, from a checkout of `github.com/openai/math` at commit `adc7f12`, the transitive closure of OpenAI’s Lean modules proving the two inputs: 2459 files, depending only on Mathlib. These are built with the same toolchain. A further module then states `theorem : FalconerStatement := OAI.Falconer.falconer_distance_conjecture` and the analogous dyadic statement, and derives unconditional versions of Theorems A and B. The kernel accepts these, and `#print axioms` reports only `propext`, `Classical.choice` and `Quot.sound`. We additionally replayed the resulting environment with Lean’s `leanchecker`. We did not run OpenAI’s `comparator` setup. A formal proof certifies the statement as formalized. Readers should still compare the formal statements with the informal ones.

ACKNOWLEDGMENTS AND USE OF AI

This note builds on results produced by an AI model and released by OpenAI [13, 12]. It was itself prepared with substantial assistance from Claude (Anthropic), an AI system, which helped choose the combination studied here, search the literature, draft the text, and write the

Lean formalization. The author checked the mathematics and the formal statements and takes full responsibility for the content.

REFERENCES

1. Jean Bourgain, *Construction of sets of positive measure not containing an affine image of a given infinite structure*, Israel J. Math. **60** (1987), 333–344.
2. Leonardo de Moura and Sebastian Ullrich, *The Lean 4 theorem prover and programming language*, Automated Deduction – CADE 28, Lecture Notes in Computer Science, vol. 12699, 2021, pp. 625–635.
3. Paul Erdős, *Remarks on some problems in number theory*, Math. Balkanica **4** (1974), 197–202.
4. M. Burak Erdoğan, *A bilinear Fourier extension theorem and applications to the distance set problem*, Int. Math. Res. Not. (2005), no. 23, 1411–1425.
5. Kenneth J. Falconer, *On a problem of Erdős on sequences and measurable sets*, Proc. Amer. Math. Soc. **90** (1984), 77–78.
6. ———, *On the Hausdorff dimensions of distance sets*, Mathematika **32** (1985), 206–212.
7. Larry Guth, Alex Iosevich, Yumeng Ou, and Hong Wang, *On Falconer’s distance set problem in the plane*, Invent. Math. **219** (2020), 779–830.
8. Yeonwook Jung, Chun-Kit Lai, and Yuveshen Moorooogen, *Fifty years of the Erdős similarity conjecture*, arXiv:2412.11062, 2024.
9. Mihail N. Kolountzakis, *Infinite patterns that can be avoided by measure*, Bull. London Math. Soc. **29** (1997), 415–424.
10. Pertti Mattila, *Fourier analysis and Hausdorff dimension*, Cambridge Studies in Advanced Mathematics, vol. 150, Cambridge University Press, 2015.
11. Pertti Mattila and Per Sjölin, *Regularity of distance measures and sets*, Math. Nachr. **204** (1999), 157–162.
12. OpenAI, *The dyadic case of the Erdős similarity conjecture*, OpenAI Math Release preprint, September 25, 2026, <https://github.com/openai/math>, commit `adc7f12`, 2026.
13. ———, *The Falconer distance conjecture in all dimensions*, OpenAI Math Release preprint, September 23, 2026, <https://github.com/openai/math>, commit `adc7f12`, 2026.
14. ———, *The geometric case of the Erdős similarity conjecture*, OpenAI Math Release preprint, October 5, 2026, <https://github.com/openai/math>, commit `adc7f12`, 2026.
15. Yuval Peres and Wilhelm Schlag, *Smoothness of projections, Bernoulli convolutions, and the dimension of exceptions*, Duke Math. J. **102** (2000), 193–251.
16. Hugo Steinhaus, *Sur les distances des points dans les ensembles de mesure positive*, Fund. Math. **1** (1920), 93–104.
17. The mathlib Community, *The Lean mathematical library*, Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), 2020, pp. 367–381.
18. Thomas Wolff, *Decay of circular means of Fourier transforms of measures*, Int. Math. Res. Not. (1999), no. 10, 547–567.

INDEPENDENT RESEARCHER, VIENNA, AUSTRIA
 Email address: harald.schilly@gmail.com