Information For the Use of
Erdosteine (300mg) + Acebrophylline (100mg) Bilayered Tablet

Erdoxol AB Tablet

ERDOXOL AB TABLET

Erdosteine (300 mg) + Acebrophylline (100 mg)

Composition

Each film-coated tablet contains:

- **Erdosteine – 300 mg**
- **Acebrophylline – 100 mg**

Pharmacological Action

Erdosteine is a mucolytic agent, while acebrophylline provides bronchodilator and mucoregulating activity. Together, they help reduce mucus viscosity and improve airway function. Medicines

It functions by:

- **Mucolytic Action:** Erdosteine and its active metabolite help break disulfide bonds within bronchial mucus, reducing mucus viscosity and facilitating expectoration.
- **Mucoregulation:** Helps improve mucociliary clearance and the transport of respiratory secretions.
- **Bronchodilation:** Acebrophylline helps relax bronchial smooth muscle and improve airflow.
- **Anti-inflammatory Support:** Acebrophylline helps modulate inflammatory processes within the respiratory tract.
- **Combined Respiratory Action:** The combination provides mucolytic and bronchodilator effects useful in obstructive respiratory conditions. Medicines

Indications

Erdoxol AB Tablets are indicated for the management of:

- **Chronic bronchitis in patients with Chronic Obstructive Pulmonary Disease (COPD)**
- COPD associated with excessive or thick respiratory secretions
- Respiratory symptoms associated with mucus hypersecretion and airflow limitation, as prescribed by the physician. CDSCO

Dosage and Administration

- **Recommended Dose:** As directed by the physician. A commonly used adult regimen for the individual components is **one tablet twice daily**, but the exact dose and duration should be determined according to the patient's condition and the approved product information.

- **Administration:** Swallow the tablet whole with water. It may be taken with or without food, according to the product instructions.
- **Duration of Use:** Continue treatment for the duration prescribed by the healthcare provider.
- Do not exceed the recommended dose.

Precautions

- **Pregnancy & Lactation:** Safety of erdosteine during pregnancy and breastfeeding has not been adequately established; use only when specifically advised by a healthcare professional. Medicines
- **Liver Impairment:** Use cautiously in patients with hepatic impairment. Erdosteine dosage may need to be restricted in mild liver impairment and is not recommended in severe liver failure. Medicines
- **Kidney Impairment:** Erdosteine is not recommended in patients with severe renal impairment, particularly creatinine clearance below 25 mL/min. Medicines
- **Peptic Ulcer:** Use is contraindicated in patients with active peptic ulcer disease.
- **Cardiovascular Conditions:** Acebrophylline should be used cautiously in patients with significant cardiovascular disease or susceptibility to palpitations.
- **Gastrointestinal Effects:** Nausea, abdominal discomfort or other gastrointestinal symptoms may occur.

Contraindications

- Known hypersensitivity to **erdosteine, acebrophylline, ambroxol, theophylline or other xanthine derivatives**, or any component of the formulation.
- **Active peptic ulcer disease.**
- Severe renal impairment where erdosteine is not recommended.
- Severe hepatic impairment.
- Patients with conditions for which acebrophylline or its components are contraindicated. Medicines

Adverse Reactions

Common:

- Nausea
- Vomiting
- Abdominal discomfort or epigastric pain
- Diarrhoea
- Headache
- Dizziness
- Palpitations or gastrointestinal discomfort

Rare but Serious:

- Hypersensitivity reactions
- Angioedema or severe skin reactions

- Difficulty in breathing
- Significant cardiovascular symptoms

Patients should seek medical attention if severe allergic reactions, breathing difficulty, persistent vomiting or other serious symptoms occur. Medicines

Drug Interactions

- **Antitussives:** Concomitant use with cough-suppressant medicines should be avoided unless specifically advised, as suppression of the cough reflex may lead to accumulation of loosened respiratory secretions. Manuals+
- **Other Xanthine Derivatives:** Caution is advised when used with other methylxanthines because of the potential for additive adverse effects.
- **Bronchodilators:** Concomitant respiratory medicines should be reviewed by the treating physician.
- Inform the physician about all prescription, OTC and herbal medicines being taken.

Storage

- Store in a **cool, dry place**, protected from moisture and excessive heat.
- Store according to the temperature specified on the product packaging.
- Keep the tablets in their original packaging until use.
- Keep out of reach of children.

Note

This information is intended as a **professional pharmaceutical-literature guideline**. **Erdoxol AB should be used under the supervision of a qualified healthcare professional.** The combination of **Acebrophylline 100 mg + Erdosteine 300 mg** has been documented by CDSCO as a bronchodilator and mucolytic combination for chronic bronchitis in patients with COPD. CDSCO

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for